



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 01:35 PM UTC

PDB ID : 9F1Q / pdb_00009f1q
Title : Crystal structure of a DyP-type peroxidase Fireprot variant from *Pseudomonas putida*
Authors : Lorena, C.; Borges, P.T.; Frazao, C.; Martins, L.O.
Deposited on : 2024-04-19
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	4% 91% 9%
1	B	286	4% 90% 8%
1	C	286	8% 91% 9%
1	D	286	8% 89% 10%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dyp-type peroxidase family protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	286	Total	C	N	O	S	0	0	0
			2224	1420	390	409	5			
1	B	285	Total	C	N	O	S	0	0	0
			2219	1417	389	408	5			
1	C	286	Total	C	N	O	S	0	0	0
			2224	1420	390	409	5			
1	D	284	Total	C	N	O	S	0	0	0
			2212	1412	388	407	5			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	LEU	ALA	engineered mutation	UNP Q88HV5
A	25	PHE	SER	engineered mutation	UNP Q88HV5
A	54	VAL	ALA	engineered mutation	UNP Q88HV5
A	91	ASP	GLU	engineered mutation	UNP Q88HV5
A	94	GLU	ASP	engineered mutation	UNP Q88HV5
A	100	ARG	GLN	engineered mutation	UNP Q88HV5
A	110	PHE	LEU	engineered mutation	UNP Q88HV5
A	118	ALA	GLY	engineered mutation	UNP Q88HV5
A	120	ARG	LEU	engineered mutation	UNP Q88HV5
A	125	ARG	HIS	engineered mutation	UNP Q88HV5
A	139	GLY	ASP	engineered mutation	UNP Q88HV5
A	149	LEU	ALA	engineered mutation	UNP Q88HV5
A	155	VAL	ALA	engineered mutation	UNP Q88HV5
A	169	ALA	SER	engineered mutation	UNP Q88HV5
A	174	GLU	ASP	engineered mutation	UNP Q88HV5
A	217	MET	VAL	engineered mutation	UNP Q88HV5
A	218	PRO	SER	engineered mutation	UNP Q88HV5
A	222	GLY	GLN	engineered mutation	UNP Q88HV5
A	232	PHE	LEU	engineered mutation	UNP Q88HV5
A	237	ASP	GLU	engineered mutation	UNP Q88HV5
A	279	LEU	VAL	engineered mutation	UNP Q88HV5

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	9	LEU	ALA	engineered mutation	UNP Q88HV5
B	25	PHE	SER	engineered mutation	UNP Q88HV5
B	54	VAL	ALA	engineered mutation	UNP Q88HV5
B	91	ASP	GLU	engineered mutation	UNP Q88HV5
B	94	GLU	ASP	engineered mutation	UNP Q88HV5
B	100	ARG	GLN	engineered mutation	UNP Q88HV5
B	110	PHE	LEU	engineered mutation	UNP Q88HV5
B	118	ALA	GLY	engineered mutation	UNP Q88HV5
B	120	ARG	LEU	engineered mutation	UNP Q88HV5
B	125	ARG	HIS	engineered mutation	UNP Q88HV5
B	139	GLY	ASP	engineered mutation	UNP Q88HV5
B	149	LEU	ALA	engineered mutation	UNP Q88HV5
B	155	VAL	ALA	engineered mutation	UNP Q88HV5
B	169	ALA	SER	engineered mutation	UNP Q88HV5
B	174	GLU	ASP	engineered mutation	UNP Q88HV5
B	217	MET	VAL	engineered mutation	UNP Q88HV5
B	218	PRO	SER	engineered mutation	UNP Q88HV5
B	222	GLY	GLN	engineered mutation	UNP Q88HV5
B	232	PHE	LEU	engineered mutation	UNP Q88HV5
B	237	ASP	GLU	engineered mutation	UNP Q88HV5
B	279	LEU	VAL	engineered mutation	UNP Q88HV5
C	9	LEU	ALA	engineered mutation	UNP Q88HV5
C	25	PHE	SER	engineered mutation	UNP Q88HV5
C	54	VAL	ALA	engineered mutation	UNP Q88HV5
C	91	ASP	GLU	engineered mutation	UNP Q88HV5
C	94	GLU	ASP	engineered mutation	UNP Q88HV5
C	100	ARG	GLN	engineered mutation	UNP Q88HV5
C	110	PHE	LEU	engineered mutation	UNP Q88HV5
C	118	ALA	GLY	engineered mutation	UNP Q88HV5
C	120	ARG	LEU	engineered mutation	UNP Q88HV5
C	125	ARG	HIS	engineered mutation	UNP Q88HV5
C	139	GLY	ASP	engineered mutation	UNP Q88HV5
C	149	LEU	ALA	engineered mutation	UNP Q88HV5
C	155	VAL	ALA	engineered mutation	UNP Q88HV5
C	169	ALA	SER	engineered mutation	UNP Q88HV5
C	174	GLU	ASP	engineered mutation	UNP Q88HV5
C	217	MET	VAL	engineered mutation	UNP Q88HV5
C	218	PRO	SER	engineered mutation	UNP Q88HV5
C	222	GLY	GLN	engineered mutation	UNP Q88HV5
C	232	PHE	LEU	engineered mutation	UNP Q88HV5
C	237	ASP	GLU	engineered mutation	UNP Q88HV5
C	279	LEU	VAL	engineered mutation	UNP Q88HV5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	9	LEU	ALA	engineered mutation	UNP Q88HV5
D	25	PHE	SER	engineered mutation	UNP Q88HV5
D	54	VAL	ALA	engineered mutation	UNP Q88HV5
D	91	ASP	GLU	engineered mutation	UNP Q88HV5
D	94	GLU	ASP	engineered mutation	UNP Q88HV5
D	100	ARG	GLN	engineered mutation	UNP Q88HV5
D	110	PHE	LEU	engineered mutation	UNP Q88HV5
D	118	ALA	GLY	engineered mutation	UNP Q88HV5
D	120	ARG	LEU	engineered mutation	UNP Q88HV5
D	125	ARG	HIS	engineered mutation	UNP Q88HV5
D	139	GLY	ASP	engineered mutation	UNP Q88HV5
D	149	LEU	ALA	engineered mutation	UNP Q88HV5
D	155	VAL	ALA	engineered mutation	UNP Q88HV5
D	169	ALA	SER	engineered mutation	UNP Q88HV5
D	174	GLU	ASP	engineered mutation	UNP Q88HV5
D	217	MET	VAL	engineered mutation	UNP Q88HV5
D	218	PRO	SER	engineered mutation	UNP Q88HV5
D	222	GLY	GLN	engineered mutation	UNP Q88HV5
D	232	PHE	LEU	engineered mutation	UNP Q88HV5
D	237	ASP	GLU	engineered mutation	UNP Q88HV5
D	279	LEU	VAL	engineered mutation	UNP Q88HV5

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total 6	C 3	O 3	0	0
2	A	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	C	1	Total 6	C 3	O 3	0	0
2	C	1	Total 6	C 3	O 3	0	0
2	D	1	Total 6	C 3	O 3	0	0

-
- The diagram illustrates the chemical structure of Heme (HEM). It features a central iron atom (Fe) coordinated by four nitrogen atoms (N) in a porphyrin-like ring. The ring is substituted with various side chains, including vinyl groups (C3A, C3B, C3C, C3D), methyl groups (C2A, C2B, C2C, C2D), and a long phytol chain (C1A, C1B, C1C, C1D). The central iron atom is also coordinated by a proximal histidine residue (NA) and a distal water molecule (ND). The overall structure is shown in a 3D perspective, with atoms labeled with their respective element symbols and bond types.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Cl 1	0	0
4	C	2	Total 2	Cl 2	0	0
4	D	1	Total 1	Cl 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	114	Total 114	O 114	0	0
5	B	104	Total 104	O 104	0	0
5	C	86	Total 86	O 86	0	0
5	D	88	Total 88	O 88	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dyp-type peroxidase family protein



- Molecule 1: Dyp-type peroxidase family protein




- Molecule 1: Dyp-type peroxidase family protein



- Molecule 1: Dyp-type peroxidase family protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	144.95Å 144.95Å 177.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	59.16 – 2.35 59.16 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.9 (59.16-2.35) 99.9 (59.16-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.00 (at 2.34Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.219 , 0.246 0.219 , 0.246	Depositor DCC
R_{free} test set	1999 reflections (2.23%)	wwPDB-VP
Wilson B-factor (Å ²)	49.2	Xtriage
Anisotropy	0.397	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 36.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.017 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9495	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, CL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.10	0/2282	0.31	0/3101
1	B	0.11	0/2277	0.30	0/3094
1	C	0.12	0/2282	0.33	0/3101
1	D	0.10	0/2269	0.28	0/3083
All	All	0.11	0/9110	0.31	0/12379

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2224	0	2179	13	0
1	B	2219	0	2174	17	0
1	C	2224	0	2179	17	0
1	D	2212	0	2166	16	0
2	A	18	0	24	1	0
2	B	12	0	16	0	0
2	C	12	0	16	0	0
2	D	6	0	8	1	0
3	A	43	0	30	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	43	0	30	4	0
3	C	43	0	30	2	0
3	D	43	0	30	3	0
4	B	1	0	0	0	0
4	C	2	0	0	1	0
4	D	1	0	0	1	0
5	A	114	0	0	3	0
5	B	104	0	0	2	0
5	C	86	0	0	3	0
5	D	88	0	0	1	0
All	All	9495	0	8882	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 71 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:303:CL:CL	5:C:472:HOH:O	2.35	0.81
1:B:6:GLY:O	1:B:88:ARG:NH2	2.21	0.73
1:B:212:MET:HE1	3:B:302:HEM:HAC	1.75	0.68
1:B:5:GLN:OE1	1:B:43:LYS:NZ	2.29	0.66
1:D:12:VAL:O	1:D:120:ARG:NH1	2.33	0.61

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	284/286 (99%)	276 (97%)	7 (2%)	1 (0%)	30 34
1	B	283/286 (99%)	272 (96%)	9 (3%)	2 (1%)	18 20

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	284/286 (99%)	272 (96%)	11 (4%)	1 (0%)	30	34
1	D	282/286 (99%)	273 (97%)	9 (3%)	0	100	100
All	All	1133/1144 (99%)	1093 (96%)	36 (3%)	4 (0%)	30	34

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	10	THR
1	A	5	GLN
1	B	6	GLY
1	B	5	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/226 (100%)	225 (100%)	1 (0%)	84	91
1	B	226/226 (100%)	225 (100%)	1 (0%)	84	91
1	C	226/226 (100%)	225 (100%)	1 (0%)	84	91
1	D	225/226 (100%)	224 (100%)	1 (0%)	84	91
All	All	903/904 (100%)	899 (100%)	4 (0%)	84	91

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	116	VAL
1	B	88	ARG
1	C	10	THR
1	D	116	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	172	GLN
1	D	172	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	HEM	A	303	1	50,50,50	1.40	8 (16%)	67,82,82	1.05	4 (5%)
3	HEM	D	301	1	50,50,50	1.42	8 (16%)	67,82,82	1.08	3 (4%)
2	GOL	B	301	-	5,5,5	0.92	0	5,5,5	1.11	0
2	GOL	C	301	-	5,5,5	0.87	0	5,5,5	1.12	0
2	GOL	A	302	-	5,5,5	0.93	0	5,5,5	1.02	0
2	GOL	C	305	-	5,5,5	0.92	0	5,5,5	1.08	0
3	HEM	C	302	1	50,50,50	1.40	9 (18%)	67,82,82	1.10	5 (7%)
2	GOL	D	303	-	5,5,5	0.95	0	5,5,5	1.10	0
2	GOL	A	304	-	5,5,5	0.91	0	5,5,5	1.10	0
3	HEM	B	302	1	50,50,50	1.39	8 (16%)	67,82,82	1.09	5 (7%)
2	GOL	A	301	-	5,5,5	0.93	0	5,5,5	1.08	0
2	GOL	B	304	-	5,5,5	0.93	0	5,5,5	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	303	1	-	5/14/54/54	-
3	HEM	D	301	1	-	4/14/54/54	-
2	GOL	B	301	-	-	0/4/4/4	-
2	GOL	C	301	-	-	2/4/4/4	-
2	GOL	A	302	-	-	2/4/4/4	-
2	GOL	C	305	-	-	2/4/4/4	-
3	HEM	C	302	1	-	4/14/54/54	-
2	GOL	D	303	-	-	0/4/4/4	-
2	GOL	A	304	-	-	2/4/4/4	-
3	HEM	B	302	1	-	4/14/54/54	-
2	GOL	A	301	-	-	0/4/4/4	-
2	GOL	B	304	-	-	1/4/4/4	-

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	301	HEM	FE-NC	3.50	2.06	1.95
3	C	302	HEM	FE-NB	3.39	2.05	1.94
3	C	302	HEM	FE-NC	3.37	2.06	1.95
3	A	303	HEM	FE-NB	3.23	2.04	1.94
3	B	302	HEM	FE-NB	3.21	2.04	1.94

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	HEM	C4D-ND-C1D	2.48	108.14	105.21
3	D	301	HEM	C4D-ND-C1D	2.41	108.06	105.21
3	A	303	HEM	C1B-NB-C4B	2.40	108.06	105.21
3	C	302	HEM	C4D-ND-C1D	2.38	108.02	105.21
3	D	301	HEM	C1B-NB-C4B	2.38	108.02	105.21

There are no chirality outliers.

5 of 26 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	304	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	301	GOL	O1-C1-C2-C3
2	A	302	GOL	C1-C2-C3-O3
2	C	305	GOL	O1-C1-C2-C3
2	A	302	GOL	O2-C2-C3-O3


There are no ring outliers.


6 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	303	HEM	1	0
3	D	301	HEM	3	0
3	C	302	HEM	2	0
2	D	303	GOL	1	0
3	B	302	HEM	4	0
2	A	301	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/286 (100%)	0.20	12 (4%) 40 46	34, 44, 64, 92	0
1	B	285/286 (99%)	0.25	11 (3%) 43 49	34, 47, 68, 89	0
1	C	286/286 (100%)	0.55	23 (8%) 18 21	36, 50, 76, 105	0
1	D	284/286 (99%)	0.62	24 (8%) 16 18	36, 50, 77, 105	0
All	All	1141/1144 (99%)	0.40	70 (6%) 27 31	34, 48, 72, 105	0

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	287	ALA	7.8
1	D	3	PHE	6.4
1	A	287	ALA	6.0
1	C	9	LEU	5.5
1	C	10	THR	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	B	304	6/6	0.68	0.26	69,77,81,96	0
4	CL	D	302	1/1	0.71	0.32	75,75,75,75	0
2	GOL	A	304	6/6	0.79	0.20	59,69,70,71	0
2	GOL	A	301	6/6	0.82	0.18	55,59,60,62	0
4	CL	C	304	1/1	0.84	0.15	81,81,81,81	0
2	GOL	D	303	6/6	0.89	0.14	53,55,56,62	0
2	GOL	C	305	6/6	0.90	0.13	52,56,61,61	0
2	GOL	C	301	6/6	0.92	0.13	49,54,55,58	0
4	CL	B	303	1/1	0.92	0.21	86,86,86,86	0
2	GOL	B	301	6/6	0.93	0.12	39,46,49,54	0
2	GOL	A	302	6/6	0.93	0.12	45,52,56,57	0
4	CL	C	303	1/1	0.95	0.11	62,62,62,62	0
3	HEM	D	301	43/43	0.97	0.08	37,47,54,60	0
3	HEM	C	302	43/43	0.98	0.07	36,46,52,56	0
3	HEM	A	303	43/43	0.98	0.08	32,38,45,46	0
3	HEM	B	302	43/43	0.98	0.08	33,43,50,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around HEM D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM C 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM A 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.