



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:32 AM UTC

PDB ID : 1F25 / pdb_00001f25
Title : CRYSTAL STRUCTURE OF NO COMPLEX OF THR243ASN MUTANTS OF CYTOCHROME P450NOR
Authors : Shimizu, H.; Park, S.-Y.
Deposited on : 2000-05-23
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	402	 91% 7% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3728 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITRIC OXIDE REDUCTASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	399	3100	1971	530	586	13	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	243	ASN	THR	engineered mutation	UNP P23295

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	N	O	0	0
			2	1	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	577	Total 577	O 577	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: NITRIC OXIDE REDUCTASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.62Å 81.93Å 85.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.40	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-1.40)	Depositor
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.142 , 0.204	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3728	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, GOL, NO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/3167	1.32	24/4304 (0.6%)

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	393	ASP	CA-CB-CG	10.34	122.94	112.60
1	A	161	PHE	CA-CB-CG	8.16	121.96	113.80
1	A	319	ASN	OD1-CG-ND2	-8.04	114.56	122.60
1	A	182	ARG	CA-C-N	7.95	130.78	120.44
1	A	182	ARG	C-N-CA	7.95	130.78	120.44
1	A	366	PHE	CA-CB-CG	-7.19	106.61	113.80
1	A	182	ARG	O-C-N	-6.62	115.26	122.07
1	A	322	GLU	CB-CG-CD	6.54	123.71	112.60
1	A	174	ARG	CD-NE-CZ	6.34	133.28	124.40
1	A	130	GLN	CG-CD-NE2	6.11	125.56	116.40
1	A	26	ARG	NH1-CZ-NH2	-5.96	111.55	119.30
1	A	130	GLN	CB-CG-CD	5.91	122.65	112.60
1	A	177	GLY	O-C-N	5.81	128.40	122.24
1	A	58	GLU	CB-CG-CD	5.54	122.01	112.60
1	A	130	GLN	OE1-CD-NE2	-5.54	117.06	122.60
1	A	363	THR	CA-CB-OG1	5.45	117.78	109.60
1	A	259	PRO	CA-C-N	-5.45	111.61	121.14
1	A	259	PRO	C-N-CA	-5.45	111.61	121.14
1	A	26	ARG	CD-NE-CZ	5.42	131.99	124.40
1	A	126	GLU	CA-C-N	5.38	127.80	120.54
1	A	126	GLU	C-N-CA	5.38	127.80	120.54
1	A	86	PHE	CA-CB-CG	5.09	118.89	113.80
1	A	174	ARG	NE-CZ-NH1	-5.05	116.45	121.50
1	A	349	ASP	CA-CB-CG	-5.03	107.58	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3100	0	3114	13	0
2	A	43	0	30	1	0
3	A	2	0	0	0	0
4	A	6	0	5	0	0
5	A	577	0	0	7	0
All	All	3728	0	3149	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:102:GLU:HG2	5:A:1168:HOH:O	1.83	0.78
1:A:260:ASP:HB3	5:A:1176:HOH:O	1.86	0.75
1:A:9:PRO:HG2	5:A:1051:HOH:O	2.03	0.59
1:A:359:LYS:O	1:A:363:THR:HG23	2.05	0.55
1:A:293:THR:HG22	5:A:632:HOH:O	2.06	0.54
1:A:240:GLY:HA3	2:A:501:HEM:C1C	2.45	0.51
1:A:203:LEU:CD1	1:A:226:LYS:HD3	2.41	0.50
1:A:8:PHE:CG	1:A:9:PRO:HA	2.50	0.47
1:A:104:THR:HG23	5:A:997:HOH:O	2.15	0.46
1:A:202:ARG:CZ	1:A:207:LYS:HD2	2.48	0.44
1:A:253:ALA:O	1:A:257:GLN:HG2	2.17	0.44
1:A:363:THR:HG21	5:A:1144:HOH:O	2.19	0.42
1:A:111:LYS:HD3	5:A:1031:HOH:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/402 (99%)	391 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	340/341 (100%)	333 (98%)	7 (2%)	47	15

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	LYS
1	A	49	LYS
1	A	111	LYS
1	A	130	GLN
1	A	257	GLN
1	A	260	ASP
1	A	384	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	78	GLN
1	A	114	GLN
1	A	130	GLN
1	A	170	GLN
1	A	258	HIS
1	A	327	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NO	A	502	2	0,1,1	-	-	-		
2	HEM	A	501	1,3	50,50,50	1.80	13 (26%)	67,82,82	1.86	16 (23%)
4	GOL	A	503	-	5,5,5	3.17	3 (60%)	5,5,5	1.58	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	503	-	-	0/4/4/4	-
2	HEM	A	501	1,3	-	4/14/54/54	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	GOL	O2-C2	-5.11	1.28	1.43
2	A	501	HEM	CBC-CAC	-4.08	1.10	1.30
2	A	501	HEM	CBA-CGA	3.50	1.58	1.50
4	A	503	GOL	O1-C1	-3.39	1.28	1.42
2	A	501	HEM	C4A-C3A	3.37	1.51	1.43
4	A	503	GOL	O3-C3	-3.35	1.28	1.42
2	A	501	HEM	CAB-C3B	3.09	1.55	1.47
2	A	501	HEM	C4A-NA	-3.09	1.33	1.39
2	A	501	HEM	C3C-C4C	3.08	1.51	1.46
2	A	501	HEM	C1A-C2A	-3.02	1.38	1.44
2	A	501	HEM	C4D-ND	2.98	1.45	1.40
2	A	501	HEM	C1C-C2C	2.49	1.50	1.45
2	A	501	HEM	C3C-C2C	2.30	1.41	1.37
2	A	501	HEM	CMA-C3A	-2.28	1.46	1.50
2	A	501	HEM	CHB-C4A	-2.18	1.34	1.39
2	A	501	HEM	FE-NB	2.15	2.01	1.94

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	HEM	C4A-C3A-C2A	-4.88	101.22	106.82
2	A	501	HEM	C2A-C1A-NA	4.14	114.75	110.15
2	A	501	HEM	C3C-C2C-C1C	-3.91	103.34	107.05
2	A	501	HEM	CHC-C4B-NB	3.70	128.41	124.42
2	A	501	HEM	C4B-C3B-C2B	3.49	110.49	107.28
2	A	501	HEM	CMB-C2B-C1B	-3.44	119.66	125.03
2	A	501	HEM	C3A-C4A-NA	3.31	115.45	110.14
2	A	501	HEM	CMB-C2B-C3B	3.18	136.11	128.43
2	A	501	HEM	C3B-C2B-C1B	-3.00	104.16	106.41
2	A	501	HEM	C2C-C1C-NC	2.87	114.95	109.64
2	A	501	HEM	C4D-C3D-C2D	2.79	110.96	106.89
2	A	501	HEM	C4A-NA-C1A	-2.75	101.34	105.82
2	A	501	HEM	CMA-C3A-C4A	2.71	129.55	125.42
4	A	503	GOL	O1-C1-C2	2.51	121.67	110.38
2	A	501	HEM	C1D-C2D-C3D	-2.49	104.36	106.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	HEM	CHB-C4A-C3A	-2.29	120.77	127.43
2	A	501	HEM	CHD-C4C-C3C	-2.29	121.36	125.21
4	A	503	GOL	C3-C2-C1	2.18	119.78	111.80

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	HEM	CAA-CBA-CGA-O1A
2	A	501	HEM	CAA-CBA-CGA-O2A
2	A	501	HEM	CAD-CBD-CGD-O2D
2	A	501	HEM	CAD-CBD-CGD-O1D

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.