



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:08 AM UTC

PDB ID : 5F2I / pdb_00005f2i
Title : Tagatose-1,6-bisphosphate aldolase from Streptococcus pyogenes Glu163Gln mutant
Authors : LowKam, C.
Deposited on : 2015-12-01
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	2% 91% 7% ..
1	B	327	2% 90% 8% ..
1	C	327	4% 94%
1	D	327	3% 87% 11% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 21764 atoms, of which 10206 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tagatose 1,6-diphosphate aldolase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	324	Total	C	H	N	O	S	0	2	0
			5086	1624	2528	422	504	8			
1	B	324	Total	C	H	N	O	S	0	3	0
			5107	1626	2543	424	505	9			
1	C	324	Total	C	H	N	O	S	0	2	0
			5096	1623	2538	423	503	9			
1	D	326	Total	C	H	N	O	S	0	3	0
			5137	1638	2555	427	508	9			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	163	GLN	GLU	engineered mutation	UNP P63705
B	163	GLN	GLU	engineered mutation	UNP P63705
C	163	GLN	GLU	engineered mutation	UNP P63705
D	163	GLN	GLU	engineered mutation	UNP P63705

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Ca	0	0
			3	3		
2	B	4	Total	Ca	0	0
			4	4		
2	C	2	Total	Ca	0	0
			2	2		
2	D	2	Total	Ca	0	0
			2	2		

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	0	1
			27	6	15	6		
4	B	1	Total	C	H	O	0	0
			14	3	8	3		
4	C	1	Total	C	H	O	0	0
			14	3	8	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total 14	C 3	H 8	O 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	355	Total 355	O 355	0	0
5	B	358	Total 358	O 358	0	0
5	C	243	Total 243	O 243	0	0
5	D	295	Total 295	O 295	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tagatose 1,6-diphosphate aldolase 2




- Molecule 1: Tagatose 1,6-diphosphate aldolase 2



- Molecule 1: Tagatose 1,6-diphosphate aldolase 2



- Molecule 1: Tagatose 1,6-diphosphate aldolase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.08Å 98.54Å 237.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.37 – 1.95 49.37 – 1.95	Depositor EDS
% Data completeness (in resolution range)	84.4 (49.37-1.95) 84.8 (49.37-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.20 (at 1.94Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.165 , 0.202 0.174 , 0.209	Depositor DCC
R_{free} test set	2000 reflections (2.11%)	wwPDB-VP
Wilson B-factor (Å ²)	25.0	Xtriage
Anisotropy	0.786	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 55.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	21764	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, CA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	9/2612 (0.3%)	0.78	5/3528 (0.1%)
1	B	0.80	10/2619 (0.4%)	0.84	7/3534 (0.2%)
1	C	0.62	2/2609 (0.1%)	0.75	9/3522 (0.3%)
1	D	0.80	9/2633 (0.3%)	0.77	4/3555 (0.1%)
All	All	0.74	30/10473 (0.3%)	0.79	25/14139 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	6	GLU	C-N	-8.13	1.22	1.34
1	B	203	VAL	C-O	-6.50	1.17	1.24
1	B	90	ALA	C-O	-6.45	1.16	1.23
1	B	7[A]	ASN	C-N	-6.25	1.25	1.33
1	B	7[B]	ASN	C-N	-6.25	1.25	1.33
1	A	203	VAL	C-O	-6.24	1.17	1.24
1	A	315	LEU	C-O	-6.17	1.17	1.24
1	B	197	GLU	C-O	-6.08	1.17	1.24
1	A	11	SER	C-O	-6.00	1.17	1.24
1	B	315	LEU	C-O	-5.99	1.17	1.24
1	A	316	ASP	C-O	-5.88	1.17	1.24
1	D	203	VAL	C-O	-5.76	1.18	1.24
1	C	41	THR	C-O	-5.70	1.16	1.24
1	D	196	LYS	C-O	-5.65	1.17	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	8	LYS	C-O	-5.50	1.17	1.24
1	B	206	VAL	C-O	-5.48	1.17	1.23
1	A	250	ALA	C-O	-5.47	1.19	1.24
1	D	257	PHE	C-O	-5.47	1.17	1.24
1	D	206	VAL	C-O	-5.43	1.17	1.23
1	D	193	VAL	C-O	-5.27	1.18	1.24
1	A	249	SER	C-O	-5.21	1.17	1.24
1	B	89	LEU	C-O	-5.21	1.17	1.24
1	D	320	SER	C-N	5.21	1.40	1.33
1	B	11	SER	C-O	-5.17	1.18	1.24
1	A	7	ASN	C-O	-5.15	1.18	1.24
1	A	13	GLU	C-O	-5.15	1.18	1.24
1	C	34	ARG	CD-NE	-5.14	1.39	1.46
1	D	202	ASP	C-O	-5.08	1.17	1.23
1	D	321	PRO	N-CD	5.02	1.54	1.47
1	D	10	LYS	C-O	-5.01	1.18	1.24

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	248	LEU	CA-C-N	13.71	139.76	120.29
1	B	248	LEU	C-N-CA	13.71	139.76	120.29
1	A	252	VAL	N-CA-C	8.85	120.89	108.23
1	A	250	ALA	CA-C-N	8.74	137.44	121.70
1	A	250	ALA	C-N-CA	8.74	137.44	121.70
1	B	6	GLU	O-C-N	-7.75	113.79	122.08
1	C	39	HIS	N-CA-C	6.98	121.88	112.88
1	B	292	GLY	CA-C-N	6.73	125.97	118.97
1	B	292	GLY	C-N-CA	6.73	125.97	118.97
1	C	292	GLY	CA-C-N	6.04	125.25	118.97
1	C	292	GLY	C-N-CA	6.04	125.25	118.97
1	C	38	GLN	N-CA-C	6.03	117.85	111.28
1	B	61	THR	CA-C-N	-6.01	113.93	119.82
1	B	61	THR	C-N-CA	-6.01	113.93	119.82
1	C	325	LYS	N-CA-C	-5.89	106.73	114.04
1	D	320	SER	CA-C-N	-5.71	114.08	119.85
1	D	320	SER	C-N-CA	-5.71	114.08	119.85
1	C	61	THR	CA-C-N	-5.51	114.14	119.87
1	C	61	THR	C-N-CA	-5.51	114.14	119.87
1	A	292	GLY	CA-C-N	5.50	124.68	118.97
1	A	292	GLY	C-N-CA	5.50	124.68	118.97
1	C	37	ALA	CA-C-N	5.40	127.52	120.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	37	ALA	C-N-CA	5.40	127.52	120.28
1	D	292	GLY	CA-C-N	5.37	124.55	118.97
1	D	292	GLY	C-N-CA	5.37	124.55	118.97

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	325	LYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2558	2528	2526	28	0
1	B	2564	2543	2528	18	0
1	C	2558	2538	2532	24	0
1	D	2582	2555	2560	31	0
2	A	3	0	0	0	0
2	B	4	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	4	3	3	1	0
4	B	18	23	24	0	0
4	C	6	8	8	3	0
4	D	6	8	8	0	0
5	A	355	0	0	12	1
5	B	358	0	0	7	1
5	C	243	0	0	3	0
5	D	295	0	0	5	0
All	All	11558	10206	10189	101	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:39:HIS:NE2	1:C:286:LYS:HG2	1.16	1.48
1:C:39:HIS:NE2	1:C:286:LYS:CG	2.01	1.23
1:A:275:LEU:CD2	5:A:669:HOH:O	1.92	1.14
1:C:39:HIS:CD2	1:C:286:LYS:HG2	1.84	1.12
1:D:326:MET:HE2	1:D:326:MET:HA	1.53	0.90
1:D:6:GLU:N	1:D:6:GLU:OE1	2.17	0.77
1:A:275:LEU:HD21	5:A:669:HOH:O	1.70	0.77
1:D:326:MET:HA	1:D:326:MET:CE	2.17	0.72
1:C:39:HIS:NE2	1:C:286:LYS:CD	2.53	0.72
1:D:325:LYS:HE3	5:D:549:HOH:O	1.89	0.71
1:C:31:ALA:HA	1:C:34:ARG:NH2	2.07	0.70
1:A:275:LEU:HD23	5:A:669:HOH:O	1.75	0.69
1:D:1:THR:HA	1:D:197:GLU:HB2	1.74	0.69
1:C:38:GLN:C	1:C:39:HIS:HD1	2.01	0.68
1:C:39:HIS:CD2	1:C:286:LYS:CG	2.65	0.68
1:C:31:ALA:HA	1:C:34:ARG:CZ	2.23	0.67
1:D:173:ASP:OD2	5:D:501:HOH:O	2.14	0.66
1:A:228:GLU:OE2	1:A:235:ARG:NH2	2.30	0.64
1:C:174[B]:ASN:ND2	5:C:505:HOH:O	2.29	0.64
1:D:2:ILE:HG22	1:D:3:THR:N	2.12	0.64
5:B:787:HOH:O	1:D:140:GLU:HG2	1.96	0.64
3:A:404:ACT:OXT	5:A:501:HOH:O	2.15	0.63
1:D:325:LYS:HG3	5:D:549:HOH:O	1.98	0.63
1:B:9:ARG:HD3	5:B:527:HOH:O	1.99	0.63
1:C:316:ASP:OD1	4:C:403:GOL:H31	2.00	0.62
1:D:1:THR:HG22	1:D:1:THR:O	1.99	0.62
1:A:250:ALA:CB	1:A:251:GLY:HA3	2.30	0.61
1:D:326:MET:HE2	1:D:326:MET:CA	2.29	0.61
1:B:163:GLN:HB2	1:B:205:LYS:HZ3	1.68	0.59
1:C:205:LYS:CE	1:C:275:LEU:HD22	2.33	0.59
1:D:3:THR:HG23	1:D:202:ASP:OD1	2.03	0.59
1:A:2:ILE:HG21	5:A:799:HOH:O	2.03	0.59
1:A:205:LYS:HE3	5:A:727:HOH:O	2.03	0.59
1:A:192:LYS:NZ	1:B:17:VAL:HG22	2.19	0.58
1:A:275:LEU:CD2	5:A:727:HOH:O	2.51	0.58
1:A:25:ALA:HB2	1:A:275:LEU:HG	1.85	0.56
1:B:205:LYS:HG2	1:B:246:ILE:HD11	1.87	0.56
4:C:403:GOL:H11	5:C:596:HOH:O	2.08	0.54
1:A:250:ALA:HB3	1:A:251:GLY:HA3	1.91	0.53
1:D:325:LYS:CE	5:D:549:HOH:O	2.52	0.53
1:A:2:ILE:HG22	1:A:3:THR:N	2.23	0.52
1:A:192:LYS:HZ2	1:B:17:VAL:HG22	1.74	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:39:HIS:HD1	1:C:39:HIS:N	2.07	0.52
1:D:2:ILE:CG2	1:D:3:THR:N	2.73	0.52
1:B:7[B]:ASN:ND2	5:B:512:HOH:O	2.43	0.52
1:B:36:MET:HE2	1:B:289:ILE:HD11	1.92	0.51
1:D:216:GLU:HG3	1:D:223:VAL:HG11	1.92	0.51
1:A:2:ILE:CG2	1:A:3:THR:N	2.73	0.51
1:C:30:GLY:C	1:C:34:ARG:NH1	2.69	0.51
1:C:39:HIS:CE1	1:C:286:LYS:HE2	2.46	0.50
1:D:325:LYS:O	1:D:326:MET:HB2	2.11	0.50
1:B:27:ASP:OD1	5:B:501:HOH:O	2.19	0.50
1:A:275:LEU:HD22	5:A:727:HOH:O	2.11	0.49
1:C:39:HIS:CE1	1:C:286:LYS:HG2	2.23	0.49
1:C:39:HIS:CD2	1:C:286:LYS:CB	2.96	0.49
1:D:2:ILE:HG12	1:D:243:LEU:HD23	1.94	0.49
1:D:205:LYS:HG2	1:D:246:ILE:HD11	1.95	0.49
1:A:251:GLY:C	1:A:252:VAL:CG1	2.85	0.48
1:A:251:GLY:C	1:A:252:VAL:HG13	2.37	0.48
1:C:61:THR:N	1:C:62:PRO:HD2	2.28	0.48
1:A:275:LEU:HD23	5:A:727:HOH:O	2.11	0.47
1:C:205:LYS:NZ	1:C:275:LEU:HD13	2.30	0.47
1:A:2:ILE:HD12	1:A:2:ILE:N	2.30	0.46
1:A:61:THR:N	1:A:62:PRO:CD	2.79	0.46
1:D:1:THR:C	1:D:2:ILE:HG13	2.40	0.46
1:B:25:ALA:HB2	1:B:275:LEU:HG	1.97	0.46
1:D:61:THR:N	1:D:62:PRO:CD	2.79	0.46
1:D:272:ASN:HA	1:D:320:SER:O	2.15	0.45
1:D:286:LYS:NZ	1:D:290:GLU:OE2	2.45	0.45
1:A:205:LYS:HG2	1:A:246:ILE:HD11	1.98	0.45
1:B:205:LYS:HE2	1:B:275:LEU:HD22	1.98	0.45
1:B:213:LYS:NZ	5:B:517:HOH:O	2.49	0.45
1:A:226:THR:HB	5:A:700:HOH:O	2.16	0.45
1:B:61:THR:N	1:B:62:PRO:CD	2.80	0.45
1:B:161:TYR:HD2	1:B:205:LYS:HD2	1.82	0.45
1:D:4:LEU:HA	1:D:8:LYS:HD2	2.00	0.44
1:A:197:GLU:OE1	5:A:502:HOH:O	2.21	0.44
1:B:274:VAL:HG22	1:B:315:LEU:HD11	2.00	0.44
1:B:222:GLU:O	5:B:502:HOH:O	2.20	0.44
1:C:39:HIS:N	1:C:39:HIS:ND1	2.64	0.44
1:A:161:TYR:HB3	1:A:205:LYS:HD2	2.00	0.44
1:D:35:MET:O	1:D:38:GLN:HG2	2.18	0.44
1:C:316:ASP:OD1	4:C:403:GOL:C3	2.65	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:174[B]:ASN:OD1	5:D:502:HOH:O	2.21	0.43
1:B:36:MET:HE1	1:B:48:GLN:HB3	1.99	0.43
1:D:205:LYS:NZ	1:D:275:LEU:HD22	2.34	0.43
1:D:243:LEU:O	1:D:325:LYS:HE2	2.18	0.43
1:B:141:GLN:OE1	5:B:503:HOH:O	2.22	0.43
1:D:2:ILE:HG12	1:D:243:LEU:CD2	2.50	0.42
1:C:205:LYS:HE3	1:C:275:LEU:HD22	2.01	0.42
1:D:249:SER:HA	1:D:257:PHE:CE1	2.55	0.42
1:C:25:ALA:HB2	1:C:275:LEU:HG	2.02	0.42
1:A:250:ALA:HB1	1:A:251:GLY:HA3	2.02	0.41
1:A:250:ALA:CB	1:A:251:GLY:CA	2.99	0.41
1:C:205:LYS:NZ	1:C:275:LEU:HD22	2.35	0.41
1:D:131:ASP:OD1	1:D:186:LYS:NZ	2.51	0.41
1:D:3:THR:O	1:D:4:LEU:HG	2.21	0.40
1:B:315:LEU:HD23	1:B:315:LEU:HA	1.93	0.40
1:A:2:ILE:CG2	5:A:799:HOH:O	2.67	0.40
1:C:286:LYS:HG3	5:C:546:HOH:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:690:HOH:O	5:B:796:HOH:O[3_655]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	324/327 (99%)	315 (97%)	9 (3%)	0	100	100
1	B	325/327 (99%)	317 (98%)	8 (2%)	0	100	100
1	C	324/327 (99%)	315 (97%)	9 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	327/327 (100%)	317 (97%)	10 (3%)	0	100	100
All	All	1300/1308 (99%)	1264 (97%)	36 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	273/274 (100%)	272 (100%)	1 (0%)	84	85
1	B	274/274 (100%)	273 (100%)	1 (0%)	84	85
1	C	272/274 (99%)	269 (99%)	3 (1%)	65	64
1	D	275/274 (100%)	272 (99%)	3 (1%)	65	64
All	All	1094/1096 (100%)	1086 (99%)	8 (1%)	76	76

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	192	LYS
1	B	205	LYS
1	C	34	ARG
1	C	39	HIS
1	C	196	LYS
1	D	4	LEU
1	D	5	THR
1	D	14	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	HIS
1	B	39	HIS
1	C	232	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 17 ligands modelled in this entry, 11 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	B	405[B]	-	5,5,5	0.36	0	5,5,5	0.23	0
3	ACT	A	404	-	3,3,3	0.66	0	3,3,3	1.44	0
4	GOL	C	403	-	5,5,5	1.30	0	5,5,5	2.17	1 (20%)
4	GOL	B	406	-	5,5,5	0.39	0	5,5,5	0.38	0
4	GOL	D	403	-	5,5,5	0.37	0	5,5,5	0.29	0
4	GOL	B	405[A]	-	5,5,5	0.37	0	5,5,5	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	405[B]	-	-	2/4/4/4	-
4	GOL	C	403	-	-	4/4/4/4	-
4	GOL	B	406	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	403	-	-	2/4/4/4	-
4	GOL	B	405[A]	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	403	GOL	O3-C3-C2	-4.29	91.08	110.38

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	405[B]	GOL	O1-C1-C2-C3
4	B	406	GOL	O1-C1-C2-C3
4	C	403	GOL	O1-C1-C2-C3
4	C	403	GOL	C1-C2-C3-O3
4	B	405[B]	GOL	O1-C1-C2-O2
4	D	403	GOL	O1-C1-C2-C3
4	C	403	GOL	O2-C2-C3-O3
4	D	403	GOL	O1-C1-C2-O2
4	B	406	GOL	O1-C1-C2-O2
4	C	403	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	404	ACT	1	0
4	C	403	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	324/327 (99%)	-0.07	7 (2%) 62 69	15, 33, 67, 94	1 (0%)
1	B	324/327 (99%)	-0.16	6 (1%) 66 74	12, 30, 58, 98	3 (0%)
1	C	324/327 (99%)	0.21	12 (3%) 45 52	21, 43, 81, 124	1 (0%)
1	D	326/327 (99%)	0.16	10 (3%) 51 58	12, 38, 74, 107	4 (1%)
All	All	1298/1308 (99%)	0.04	35 (2%) 56 62	12, 36, 74, 124	9 (0%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	2	ILE	7.4
1	D	4	LEU	7.1
1	D	326	MET	6.4
1	D	1	THR	6.2
1	D	5	THR	5.9
1	A	2	ILE	5.8
1	D	3	THR	5.5
1	D	250	ALA	3.9
1	C	3	THR	3.9
1	B	249	SER	3.8
1	C	41	THR	3.0
1	A	41	THR	2.9
1	B	205	LYS	2.8
1	B	3	THR	2.8
1	C	326	MET	2.7
1	D	251	GLY	2.6
1	C	251	GLY	2.6
1	D	325	LYS	2.6
1	A	252	VAL	2.4
1	D	242	ASP	2.3
1	C	286	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	98	ALA	2.3
1	C	4	LEU	2.3
1	B	326	MET	2.2
1	A	282	ALA	2.2
1	C	35	MET	2.2
1	C	84	GLU	2.2
1	A	279	ALA	2.2
1	C	289	ILE	2.1
1	B	224	LEU	2.1
1	B	279	ALA	2.1
1	A	3	THR	2.1
1	C	38	GLN	2.1
1	C	250	ALA	2.0
1	A	220	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	C	403	6/6	0.68	0.20	67,81,82,82	0
4	GOL	D	403	6/6	0.68	0.20	77,93,94,94	0
3	ACT	A	404	4/4	0.76	0.13	30,32,36,36	0
4	GOL	B	405[A]	6/6	0.86	0.23	52,62,65,66	14
4	GOL	B	405[B]	6/6	0.86	0.23	51,55,65,66	13
4	GOL	B	406	6/6	0.88	0.11	50,60,61,62	0
2	CA	A	403	1/1	0.90	0.10	83,83,83,83	0
2	CA	B	401	1/1	0.92	0.13	70,70,70,70	0
2	CA	B	402	1/1	0.92	0.09	61,61,61,61	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CA	D	402	1/1	0.92	0.14	52,52,52,52	0
2	CA	D	401	1/1	0.94	0.06	35,35,35,35	0
2	CA	B	404	1/1	0.95	0.07	67,67,67,67	0
2	CA	C	402	1/1	0.95	0.07	50,50,50,50	0
2	CA	A	401	1/1	0.96	0.07	53,53,53,53	0
2	CA	C	401	1/1	0.97	0.04	27,27,27,27	0
2	CA	B	403	1/1	0.99	0.03	25,25,25,25	0
2	CA	A	402	1/1	0.99	0.03	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.