



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:56 PM UTC

PDB ID : 7F2F / pdb\_00007f2f  
Title : The complex of DNA with the C-terminal domain of TYE7 from *Saccharomyces cerevisiae*.  
Authors : Gui, W.  
Deposited on : 2021-06-10  
Resolution : 2.55 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	136	
1	B	136	
2	E	15	
3	H	15	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2178 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine-rich protein TYE7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	90	764	483	136	143	2	0	1	0
1	B	94	802	508	145	147	2	0	1	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	164	MET	-	expression tag	UNP P33122
A	292	LEU	-	expression tag	UNP P33122
A	293	GLU	-	expression tag	UNP P33122
A	294	HIS	-	expression tag	UNP P33122
A	295	HIS	-	expression tag	UNP P33122
A	296	HIS	-	expression tag	UNP P33122
A	297	HIS	-	expression tag	UNP P33122
A	298	HIS	-	expression tag	UNP P33122
A	299	HIS	-	expression tag	UNP P33122
B	164	MET	-	expression tag	UNP P33122
B	292	LEU	-	expression tag	UNP P33122
B	293	GLU	-	expression tag	UNP P33122
B	294	HIS	-	expression tag	UNP P33122
B	295	HIS	-	expression tag	UNP P33122
B	296	HIS	-	expression tag	UNP P33122
B	297	HIS	-	expression tag	UNP P33122
B	298	HIS	-	expression tag	UNP P33122
B	299	HIS	-	expression tag	UNP P33122

- Molecule 2 is a DNA chain called DNA (5'-D(\*CP\*AP\*GP\*AP\*TP\*CP\*AP\*TP\*GP\*TP\*GP\*TP\*GP\*CP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	E	14	288	137	52	85	14	0	0	0


- Molecule 3 is a DNA chain called DNA (5'-D(\*GP\*GP\*GP\*CP\*AP\*CP\*AP\*CP\*AP\*TP\*GP\*AP\*TP\*CP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	H	15	324	156	63	91	14	0	1	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine-rich protein TYE7




- Molecule 1: Serine-rich protein TYE7



- Molecule 2: DNA (5'-D(\*CP\*AP\*GP\*AP\*TP\*CP\*AP\*TP\*GP\*TP\*GP\*TP\*GP\*CP\*C)-3')



- Molecule 3: DNA (5'-D(\*GP\*GP\*GP\*CP\*AP\*CP\*AP\*CP\*AP\*TP\*GP\*AP\*TP\*CP\*T)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.03Å 73.03Å 181.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	67.76 – 2.55 67.76 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.7 (67.76-2.55) 99.7 (67.76-2.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.58 (at 2.54Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, $R_{free}$	0.246 , 0.284 0.251 , 0.291	Depositor DCC
$R_{free}$ test set	838 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	64.1	Xtrriage
Anisotropy	0.152	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 64.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2178	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	72.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.09	0/773	1.19	1/1037 (0.1%)
1	B	1.15	0/813	1.16	2/1090 (0.2%)
2	E	0.49	0/322	1.13	3/495 (0.6%)
3	H	0.45	0/364	1.22	6/561 (1.1%)
All	All	0.97	0/2272	1.18	12/3183 (0.4%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	2	DG	C2'-C3'-O3'	7.61	122.92	111.50
3	H	4	DC	C4'-C3'-O3'	7.33	121.00	110.00
3	H	5	DA	C4'-C3'-O3'	6.81	120.21	110.00
2	E	9	DG	C2'-C3'-O3'	5.96	120.44	111.50
1	A	287	LYS	N-CA-C	5.92	118.52	111.71
3	H	5	DA	C5'-C4'-C3'	-5.85	106.12	114.90
3	H	14	DC	C2'-C3'-O3'	5.84	120.27	111.50
2	E	3	DG	C2'-C3'-O3'	5.72	120.08	111.50
3	H	2	DG	C4'-C3'-O3'	-5.53	101.71	110.00
2	E	2	DA	C2'-C3'-O3'	5.47	119.70	111.50
1	B	252	SER	N-CA-C	5.38	117.84	111.33
1	B	185	HIS	N-CA-C	-5.14	105.12	111.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	764	0	784	6	0
1	B	802	0	824	8	0
2	E	288	0	159	5	0
3	H	324	0	182	5	0
All	All	2178	0	1949	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:192:TYR:OH	2:E:6:DC:OP2	1.92	0.87
3:H:4:DC:H2'	3:H:5:DA:C8	2.30	0.66
1:A:206:ILE:HD11	1:B:263:LEU:HD21	1.88	0.55
1:A:206:ILE:HD11	1:B:263:LEU:CD2	2.37	0.55
2:E:10:DT:H2'	2:E:11:DG:C8	2.43	0.54
2:E:15:DC:N3	3:H:2:DG:O6	2.42	0.53
1:B:253:MET:HE2	1:B:257:LYS:HG2	1.94	0.48
1:A:252:SER:HB2	1:B:199:LYS:HE3	1.94	0.48
3:H:4:DC:H2''	3:H:5:DA:O5'	2.15	0.46
1:A:283:ILE:O	1:A:284:ASP:C	2.59	0.46
1:A:262:ILE:CG2	1:B:206:ILE:HD12	2.47	0.45
1:B:213:GLU:HB2	1:B:249:LEU:HD12	1.99	0.45
3:H:4:DC:H2''	3:H:5:DA:C5'	2.47	0.45
1:A:193[A]:ARG:NH2	2:E:8:DT:OP2	2.49	0.42
3:H:1[B]:DG:H2''	3:H:2:DG:C8	2.55	0.41
1:B:189:GLU:CD	2:E:6:DC:H41	2.26	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	87/136 (64%)	79 (91%)	7 (8%)	1 (1%)	11	15
1	B	91/136 (67%)	88 (97%)	3 (3%)	0	100	100
All	All	178/272 (65%)	167 (94%)	10 (6%)	1 (1%)	21	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	182	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/121 (70%)	83 (98%)	2 (2%)	43	61
1	B	89/121 (74%)	83 (93%)	6 (7%)	15	21
All	All	174/242 (72%)	166 (95%)	8 (5%)	23	36

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	206	ILE
1	A	257	LYS
1	B	177	LEU
1	B	178	THR
1	B	194	ILE
1	B	212	SER
1	B	252	SER
1	B	285	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	183	GLN
1	A	250	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	290	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	90/136 (66%)	1.59	29 (32%) <b>1</b> <b>0</b>	26, 60, 124, 141	1 (1%)
1	B	94/136 (69%)	1.80	28 (29%) <b>1</b> <b>1</b>	29, 64, 117, 145	1 (1%)
2	E	14/15 (93%)	1.08	1 (7%) 22 21	49, 64, 113, 123	0
3	H	15/15 (100%)	1.75	4 (26%) <b>1</b> <b>1</b>	39, 72, 92, 125	1 (6%)
All	All	213/302 (70%)	1.66	62 (29%) <b>1</b> <b>1</b>	26, 64, 123, 145	3 (1%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	1[A]	DG	9.3
1	A	177	LEU	9.0
1	B	186	ASN	7.0
1	B	184	ALA	6.8
1	B	182	LYS	6.6
1	B	189	GLU	6.6
1	B	177	LEU	6.2
1	A	180	PHE	6.1
1	B	180	PHE	6.1
1	B	185	HIS	5.6
1	B	247	THR	5.5
1	B	219	VAL	5.1
1	A	185	HIS	4.9
1	B	253	MET	4.8
1	A	186	ASN	4.7
1	A	189	GLU	4.7
1	B	187	LYS	4.3
1	A	188	ILE	4.1
1	B	294	HIS	4.1
1	A	179	PRO	4.1
1	A	247	THR	3.9

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
3	H	2	DG	3.8
1	A	289	ASP	3.6
1	A	184	ALA	3.5
1	B	188	ILE	3.5
1	B	220	GLY	3.3
1	A	182	LYS	3.3
1	A	284	ASP	3.2
1	A	246	SER	3.1
1	A	248	LYS	3.0
1	A	250	ASN	3.0
1	B	246	SER	3.0
1	B	181	GLN	2.8
1	B	295	HIS	2.7
1	A	181	GLN	2.6
1	B	274	MET	2.6
3	H	10	DT	2.6
1	B	178	THR	2.6
1	B	216	ALA	2.6
1	A	253	MET	2.5
1	B	191	ARG	2.5
1	B	179	PRO	2.5
3	H	9	DA	2.5
1	A	282	GLU	2.5
1	A	187	LYS	2.5
1	A	288	GLN	2.4
1	B	273	GLU	2.4
1	A	214	GLN	2.4
1	A	275	GLU	2.4
1	A	220	GLY	2.4
2	E	15	DC	2.3
1	A	213	GLU	2.2
1	A	290	GLN	2.2
1	A	285	THR	2.2
1	B	192	TYR	2.2
1	B	218	GLU	2.1
1	A	221	ASP	2.1
1	B	289	ASP	2.1
1	A	257	LYS	2.1
1	A	178	THR	2.1
1	B	270	ARG	2.0
1	B	282	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.