



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:58 PM UTC

PDB ID : 1F3D / pdb_00001f3d
Title : CATALYTIC ANTIBODY 4B2 IN COMPLEX WITH ITS AMIDINIUM HAPTEN.
Authors : Golinelli-Pimpaneau, B.; Goncalves, O.; Dintinger, T.; Blanchard, D.; Knos-sow, M.; Tellier, C.
Deposited on : 2000-06-02
Resolution : 1.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1220 (1.88-1.88)
Clashscore	190562	1234 (1.88-1.88)
Ramachandran outliers	187476	1222 (1.88-1.88)
Sidechain outliers	187428	1222 (1.88-1.88)
RSRZ outliers	180081	1220 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	J	219	4% 93% 6%
1	L	219	10% 89% 11%
2	H	217	7% 83% 16% .
2	K	217	9% 85% 13% .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7035 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CATALYTIC ANTIBODY 4B2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	218	Total	C	N	O	S	0	1	0
			1691	1061	284	340	6			
1	J	218	Total	C	N	O	S	0	3	0
			1703	1067	286	344	6			

- Molecule 2 is a protein called CATALYTIC ANTIBODY 4B2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	217	Total	C	N	O	S	0	3	0
			1648	1051	269	322	6			
2	K	215	Total	C	N	O	S	0	6	0
			1650	1054	266	324	6			

- Molecule 3 is 2-(4-AMINO BENZYLAMINO)-3,4,5,6-TETRAHYDROPYRIDINIUM (CCD ID: TPM) (formula: C₁₂H₁₈N₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	L	1	Total	C	N	0	0
			15	12	3		
3	J	1	Total	C	N	0	0
			15	12	3		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	K	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	52	Total	O	0	0
			52	52		
5	H	111	Total	O	0	0
			111	111		
5	J	65	Total	O	0	0
			65	65		
5	K	80	Total	O	0	0
			80	80		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CATALYTIC ANTIBODY 4B2




- Molecule 1: CATALYTIC ANTIBODY 4B2



- Molecule 2: CATALYTIC ANTIBODY 4B2



- Molecule 2: CATALYTIC ANTIBODY 4B2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.00Å 87.20Å 128.80Å 90.00° 97.60° 90.00°	Depositor
Resolution (Å)	20.00 – 1.87 20.00 – 1.87	Depositor EDS
% Data completeness (in resolution range)	97.5 (20.00-1.87) 97.9 (20.00-1.87)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 1.87Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.198 , 0.218 0.203 , 0.224	Depositor DCC
R_{free} test set	7999 reflections (10.05%)	wwPDB-VP
Wilson B-factor (Å ²)	25.5	Xtriage
Anisotropy	0.337	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.021 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7035	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, TPM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	J	0.36	0/1743	0.84	1/2364 (0.0%)
1	L	0.35	0/1731	0.85	2/2348 (0.1%)
2	H	0.38	0/1695	0.92	8/2318 (0.3%)
2	K	0.37	0/1696	0.93	9/2323 (0.4%)
All	All	0.37	0/6865	0.89	20/9353 (0.2%)

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	60	ASN	N-CA-C	-8.06	96.91	109.25
2	K	60	ASN	N-CA-C	-7.74	96.61	109.46
1	J	137	ASN	N-CA-C	6.49	120.62	110.17
1	L	137	ASN	N-CA-C	6.48	120.61	110.17
2	K	151	LYS	N-CA-C	6.46	119.93	109.40
2	H	151	LYS	N-CA-C	6.40	119.83	109.40
2	K	52	VAL	N-CA-C	-6.18	101.43	107.55
2	H	128	SER	N-CA-C	-5.87	99.62	109.07
2	K	207	HIS	CA-C-N	5.78	125.91	119.32
2	K	207	HIS	C-N-CA	5.78	125.91	119.32
2	K	219	VAL	CA-C-N	5.73	125.74	119.89
2	K	219	VAL	C-N-CA	5.73	125.74	119.89
2	H	207	HIS	CA-C-N	5.57	125.03	119.24
2	H	207	HIS	C-N-CA	5.57	125.03	119.24
2	K	194	SER	N-CA-C	-5.47	106.61	113.28
2	H	92	CYS	N-CA-C	-5.41	101.34	109.95
2	K	82(C)	LEU	N-CA-C	5.20	117.76	110.23
2	H	141	ASN	N-CA-C	5.13	121.73	110.80
1	L	206	VAL	N-CA-C	5.10	115.82	108.48
2	H	52	VAL	N-CA-C	-5.07	102.53	107.55

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	J	1703	0	1634	9	0
1	L	1691	0	1626	15	0
2	H	1648	0	1585	17	0
2	K	1650	0	1590	15	0
3	J	15	0	18	0	0
3	L	15	0	18	0	0
4	K	5	0	0	0	0
5	H	111	0	0	0	0
5	J	65	0	0	0	0
5	K	80	0	0	0	0
5	L	52	0	0	0	0
All	All	7035	0	6471	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:53:LYS:NZ	2:K:139:GLN:H	1.73	0.85
1:L:53:LYS:HZ1	2:K:139:GLN:H	1.24	0.84
1:L:142:LYS:H	1:L:142:LYS:CE	2.13	0.61
2:K:127:PRO:HB3	2:K:153:TYR:HB3	1.81	0.61
2:K:59:PHE:HB2	2:K:64:LYS:HE2	1.83	0.60
2:H:6:GLN:HE21	2:H:104:GLY:HA3	1.67	0.60
1:J:142:LYS:HE3	1:J:142:LYS:H	1.67	0.59
1:J:181:LEU:HD22	1:J:185:GLU:HG2	1.86	0.57
2:H:40:SER:O	2:H:41[B]:HIS:HD2	1.89	0.56
1:J:142:LYS:H	1:J:142:LYS:CE	2.19	0.56
1:L:142:LYS:H	1:L:142:LYS:HE3	1.71	0.55
2:H:141:ASN:HA	2:H:193:SER:CB	2.37	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:K:6:GLN:HE22	2:K:91:TYR:HA	1.74	0.53
2:K:6:GLN:HE21	2:K:104:GLY:HA3	1.74	0.53
2:K:12:VAL:HG21	2:K:82(C):LEU:HD13	1.94	0.50
2:H:6:GLN:HE22	2:H:91:TYR:HA	1.77	0.49
1:J:30:LYS:HD3	1:J:50:LYS:HE3	1.94	0.49
2:H:220:PRO:O	2:H:221:ARG:HB2	2.12	0.49
2:H:127:PRO:HB3	2:H:153:TYR:HB3	1.94	0.49
1:L:184:ASP:O	1:L:188:ARG:HG3	2.12	0.49
2:H:40:SER:O	2:H:41[B]:HIS:CD2	2.67	0.48
2:K:163:ASN:HD22	2:K:167:LEU:HD13	1.79	0.47
2:H:216[A]:LYS:HA	2:H:216[A]:LYS:HD2	1.70	0.47
1:J:160:LEU:HD11	2:K:179:GLN:HB3	1.96	0.47
2:K:65:GLY:O	2:K:82(A)[A]:ASN:ND2	2.47	0.47
1:L:27(E):SER:C	1:L:29:GLY:H	2.23	0.46
2:H:29:PHE:CE2	2:H:71:VAL:CG2	2.99	0.46
2:H:135:GLY:HA2	2:H:222:ASP:CB	2.44	0.46
1:J:83:LEU:HD11	1:J:106:ILE:HD11	1.96	0.46
2:K:12:VAL:HG21	2:K:82(C):LEU:CD1	2.46	0.46
2:H:59:PHE:HB2	2:H:64:LYS:HE2	1.97	0.45
2:H:47:TRP:CZ2	2:H:49:GLY:HA2	2.52	0.45
1:L:144:ILE:HG22	1:L:163:TRP:HH2	1.82	0.44
1:L:190:ASN:HD21	1:L:212:ASN:H	1.65	0.44
2:K:68:THR:HB	2:K:81:HIS:HB2	1.98	0.44
1:L:125:LEU:O	1:L:183:LYS:HD2	2.18	0.44
1:J:184:ASP:O	1:J:188:ARG:HG3	2.18	0.43
2:H:12:VAL:HG21	2:H:82(C):LEU:HD13	2.00	0.43
1:J:207:LYS:HD3	1:J:207:LYS:HA	1.96	0.42
2:K:47:TRP:CZ2	2:K:49:GLY:HA2	2.54	0.42
2:H:68:THR:HB	2:H:81:HIS:HB2	2.00	0.42
2:H:159:THR:OG1	2:H:206:ALA:HB3	2.19	0.42
1:L:30:LYS:NZ	2:K:139:GLN:NE2	2.68	0.41
1:L:18:GLN:HG3	1:L:76[A]:SER:HA	2.02	0.41
1:J:9:LEU:HD23	1:J:9:LEU:HA	1.83	0.41
1:L:105:GLU:OE2	1:L:142:LYS:HE2	2.20	0.41
2:H:29:PHE:HE2	2:H:71:VAL:CG2	2.34	0.41
2:H:174:PHE:HA	2:H:175:PRO:HD3	1.92	0.40
1:L:144:ILE:HG22	1:L:163:TRP:CH2	2.56	0.40
1:L:207:LYS:HD3	1:L:207:LYS:HA	1.94	0.40
1:L:161:ASN:HD22	1:L:177:SER:HA	1.86	0.40
2:K:159:THR:OG1	2:K:206:ALA:HB3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	J	219/219 (100%)	215 (98%)	4 (2%)	0	100	100
1	L	217/219 (99%)	212 (98%)	4 (2%)	1 (0%)	24	13
2	H	218/217 (100%)	202 (93%)	13 (6%)	3 (1%)	9	1
2	K	219/217 (101%)	210 (96%)	7 (3%)	2 (1%)	14	4
All	All	873/872 (100%)	839 (96%)	28 (3%)	6 (1%)	18	7

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	142	SER
2	K	142	SER
2	H	135	GLY
1	L	68	GLY
2	H	221	ARG
2	K	137	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	J	196/196 (100%)	193 (98%)	3 (2%)	57	46
1	L	194/196 (99%)	191 (98%)	3 (2%)	57	46
2	H	184/188 (98%)	181 (98%)	3 (2%)	55	43
2	K	186/188 (99%)	185 (100%)	1 (0%)	81	76

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	760/768 (99%)	750 (99%)	10 (1%)	61	50

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	33	LEU
1	L	142	LYS
1	L	185	GLU
2	H	157	PRO
2	H	185	LEU
2	H	221	ARG
1	J	142	LYS
1	J	179	LEU
1	J	187	GLU
2	K	185	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	137	ASN
1	L	161	ASN
1	L	190	ASN
2	H	6	GLN
2	H	82(A)	ASN
1	J	137	ASN
1	J	161	ASN
2	K	6	GLN
2	K	139	GLN
2	K	141	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TPM	J	401	-	14,16,16	1.81	1 (7%)	15,20,20	0.55	0
4	SO4	K	402	-	4,4,4	0.27	0	6,6,6	0.13	0
3	TPM	L	400	-	14,16,16	1.80	1 (7%)	15,20,20	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TPM	J	401	-	-	2/5/13/13	0/2/2/2
3	TPM	L	400	-	-	2/5/13/13	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	J	401	TPM	C8-N3	6.46	1.37	1.27
3	L	400	TPM	C8-N3	6.44	1.37	1.27

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	L	400	TPM	N3-C8-N2-C7
3	J	401	TPM	N3-C8-N2-C7
3	L	400	TPM	C4-C7-N2-C8
3	J	401	TPM	C4-C7-N2-C8

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	J	218/219 (99%)	0.32	8 (3%)	45	49	8, 28, 49, 69	4 (1%)
1	L	218/219 (99%)	0.59	21 (9%)	13	15	17, 32, 52, 63	1 (0%)
2	H	217/217 (100%)	0.33	16 (7%)	20	23	13, 24, 44, 83	3 (1%)
2	K	215/217 (99%)	0.40	19 (8%)	15	17	11, 25, 51, 84	6 (2%)
All	All	868/872 (99%)	0.41	64 (7%)	20	23	8, 27, 50, 84	14 (1%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	K	137	ALA	11.1
2	H	138	ALA	9.3
2	K	136	SER	8.9
2	H	223	CYS	7.9
2	H	136	SER	7.1
2	K	138	ALA	6.2
1	J	208	SER	5.9
2	K	140	THR	5.7
2	H	137	ALA	5.5
1	J	213	ALA	5.5
2	H	139	GLN	5.4
2	H	140	THR	5.1
2	K	139	GLN	4.9
2	H	135	GLY	4.9
2	K	141	ASN	4.7
2	H	141	ASN	4.6
1	L	29	GLY	4.5
2	H	222	ASP	4.3
1	J	212	ASN	4.2
2	K	135	GLY	4.2
2	K	221	ARG	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	L	213	ALA	4.0
2	K	42	GLY	3.9
1	L	27(C)	PHE	3.5
1	L	28	ASP	3.4
1	L	56	SER	3.3
1	L	76[A]	SER	3.3
2	H	42	GLY	3.1
1	L	27	GLN	3.0
2	H	84	PHE	3.0
1	L	27(E)	SER	2.9
2	K	203	CYS	2.8
1	L	69	THR	2.8
2	K	43	LYS	2.7
2	H	41[A]	HIS	2.7
1	L	27(B)	ILE	2.7
2	H	221	ARG	2.6
1	L	24	ARG	2.6
1	L	190	ASN	2.6
1	J	184	ASP	2.5
2	K	41	HIS	2.5
2	K	142	SER	2.4
1	L	157	ASN	2.4
2	K	198	SER	2.3
1	L	163	TRP	2.3
1	L	156	GLN	2.3
1	L	20	SER	2.2
2	H	166	SER	2.2
1	J	211	ARG	2.2
2	K	199	GLU	2.2
2	K	65	GLY	2.2
1	L	188	ARG	2.2
1	L	14	SER	2.2
1	J	126	THR	2.1
1	L	3	LEU	2.1
2	H	40	SER	2.1
2	K	195	THR	2.1
2	K	64	LYS	2.1
1	L	40	PRO	2.1
1	J	41	GLY	2.0
1	J	157	ASN	2.0
1	L	58	VAL	2.0
2	H	142	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	K	5	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	TPM	L	400	15/15	0.83	0.13	33,34,36,38	0
3	TPM	J	401	15/15	0.86	0.12	25,27,28,28	0
4	SO4	K	402	5/5	0.97	0.09	35,39,39,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.