



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:17 PM UTC

PDB ID : 6F3G / pdb_00006f3g
Title : IRAK4 IN COMPLEX WITH inhibitor
Authors : Xue, Y.; Degorce, S.L.; Robb, G.R.; Ferguson, A.D.
Deposited on : 2017-11-28
Resolution : 2.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7164 (2.40-2.36)
Clashscore	190562	7722 (2.40-2.36)
Ramachandran outliers	187476	7626 (2.40-2.36)
Sidechain outliers	187428	7627 (2.40-2.36)
RSRZ outliers	180081	7170 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	34% 82% 11% • 6%
2	B	295	30% 81% 11% • 5%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4544 atoms, of which 4 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Interleukin-1 receptor-associated kinase 4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	P	S	0	0	0
			2209	1385	371	436	3	14			

- Molecule 2 is a protein called Interleukin-1 receptor-associated kinase 4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	279	Total	C	N	O	P	S	0	0	0
			2211	1390	373	432	2	14			

- Molecule 3 is {N}4, {N}4-dimethyl- {N}1-(5-propan-2-ylpyrrolo[3,2-d]pyrimidin-4-yl)cyclohexane-1,4-diamine (CCD ID: CJN) (formula: C₁₇H₂₇N₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	N	0	0
			24	17	2	5		
3	B	1	Total	C	H	N	0	0
			24	17	2	5		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	37	Total	O	0	0
			37	37		
5	B	34	Total	O	0	0
			34	34		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Interleukin-1 receptor-associated kinase 4



- Molecule 2: Interleukin-1 receptor-associated kinase 4



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	87.51Å 110.28Å 142.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.13 – 2.37 29.13 – 2.37	Depositor EDS
% Data completeness (in resolution range)	99.9 (29.13-2.37) 99.9 (29.13-2.37)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 2.36Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.290 , 0.314 0.286 , 0.308	Depositor DCC
R_{free} test set	1384 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	49.9	Xtriage
Anisotropy	0.535	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 71.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4544	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.98 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.7309e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, SEP, TPO, CJN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/2209	1.25	5/2972 (0.2%)
2	B	0.81	1/2224 (0.0%)	1.26	7/2991 (0.2%)
All	All	0.79	1/4433 (0.0%)	1.25	12/5963 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	237	MET	SD-CE	-5.14	1.66	1.79

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	175	ASN	CA-CB-CG	-6.53	106.07	112.60
1	A	179	ASN	CA-CB-CG	6.14	118.74	112.60
2	B	181	ASP	CA-CB-CG	6.05	118.66	112.60
2	B	185	ILE	CA-C-N	5.78	128.30	120.38
2	B	185	ILE	C-N-CA	5.78	128.30	120.38
1	A	412	ASP	CA-CB-CG	5.50	118.10	112.60
2	B	185	ILE	N-CA-CB	5.43	116.53	110.51
1	A	206	ASN	CA-CB-CG	5.40	118.00	112.60
1	A	194	GLU	CB-CG-CD	5.39	121.77	112.60
2	B	412	ASP	CA-CB-CG	5.30	117.90	112.60
2	B	446	ASP	CA-CB-CG	5.05	117.66	112.60
1	A	446	ASP	CA-CB-CG	5.03	117.63	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2209	0	2180	13	0
2	B	2211	0	2182	20	4
3	A	22	2	0	0	0
3	B	22	2	0	0	0
4	B	5	0	0	0	0
5	A	37	0	0	0	0
5	B	34	0	0	1	0
All	All	4540	4	4362	32	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:178:ASN:C	2:B:179:ASN:N	2.17	1.03
1:A:169:SER:HB3	1:A:172:GLU:HG3	1.59	0.83
1:A:185:ILE:HD11	1:A:192:MET:HG2	1.65	0.78
2:B:210:VAL:CG1	2:B:261:VAL:HG13	2.20	0.71
2:B:192:MET:SD	2:B:264:TYR:HE1	2.19	0.66
2:B:210:VAL:HG11	2:B:261:VAL:HG13	1.80	0.62
1:A:237:MET:SD	1:A:330:PHE:CD2	2.98	0.57
1:A:237:MET:SD	1:A:330:PHE:HD2	2.28	0.56
1:A:390:HIS:O	2:B:390:HIS:O	2.24	0.56
2:B:210:VAL:HG13	2:B:261:VAL:HG13	1.86	0.56
1:A:409:THR:HG23	1:A:412:ASP:H	1.70	0.56
2:B:210:VAL:HG11	2:B:261:VAL:CG1	2.36	0.56
2:B:266:PRO:CG	2:B:321:GLU:HG3	2.37	0.54
1:A:222:THR:HG22	1:A:225:GLU:HG3	1.89	0.53
1:A:246:VAL:HG11	1:A:318:LEU:HD12	1.91	0.53
1:A:233:GLU:O	1:A:237:MET:HG2	2.10	0.52
2:B:174:LYS:HE2	2:B:179:ASN:OD1	2.10	0.51
2:B:266:PRO:HG3	2:B:321:GLU:HG3	1.93	0.50
2:B:210:VAL:CG1	2:B:261:VAL:CG1	2.92	0.47
1:A:345:TPO:HB	1:A:364:ILE:HD11	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:210:VAL:HG22	2:B:249:LEU:HD12	1.98	0.44
2:B:265:MET:HE1	5:B:624:HOH:O	2.17	0.44
2:B:237:MET:SD	2:B:248:LEU:HB2	2.59	0.43
2:B:246:VAL:HG21	2:B:318:LEU:HD12	2.00	0.42
2:B:192:MET:SD	2:B:264:TYR:CE1	3.08	0.42
1:A:222:THR:HG23	1:A:225:GLU:H	1.85	0.41
2:B:345:TPO:HB	2:B:364:ILE:HD11	2.00	0.41
2:B:414:ILE:HG12	2:B:426:VAL:HG11	2.01	0.41
2:B:311:ASP:HB2	2:B:332:LEU:HD12	2.02	0.41
1:A:168:PHE:HE2	1:A:250:GLY:HA3	1.86	0.41
2:B:169:SER:HB2	2:B:172:GLU:CD	2.46	0.41
1:A:414:ILE:HG12	1:A:426:VAL:HG11	2.02	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:172:GLU:OE1	2:B:172:GLU:OE1[4_545]	0.41	1.79
2:B:172:GLU:CD	2:B:172:GLU:OE1[4_545]	1.55	0.65
2:B:172:GLU:CG	2:B:172:GLU:CG[4_545]	1.81	0.39
2:B:172:GLU:CD	2:B:172:GLU:CD[4_545]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/295 (89%)	256 (97%)	7 (3%)	0	100	100
2	B	267/295 (90%)	258 (97%)	8 (3%)	1 (0%)	30	40
All	All	530/590 (90%)	514 (97%)	15 (3%)	1 (0%)	43	56

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	181	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/254 (96%)	232 (96%)	11 (4%)	24	39
2	B	243/255 (95%)	230 (95%)	13 (5%)	20	33
All	All	486/509 (96%)	462 (95%)	24 (5%)	22	36

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	164	ARG
1	A	187	VAL
1	A	194	GLU
1	A	206	ASN
1	A	328	SER
1	A	344	MET
1	A	352	THR
1	A	389	GLU
1	A	401	GLU
1	A	408	LYS
1	A	455	GLN
2	B	169	SER
2	B	185	ILE
2	B	187	VAL
2	B	206	ASN
2	B	210	VAL
2	B	213	LYS
2	B	214	LYS
2	B	290	LYS
2	B	321	GLU
2	B	389	GLU
2	B	401	GLU
2	B	408	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	455	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	206	ASN
1	A	390	HIS
1	A	435	GLN
1	A	452	GLN
2	B	390	HIS
2	B	435	GLN
2	B	452	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	A	342	1	8,10,11	1.11	0	10,14,16	1.11	2 (20%)
1	SEP	A	346	1	8,9,10	0.87	0	7,12,14	4.91	2 (28%)
1	TPO	A	345	1	8,10,11	0.98	0	10,14,16	2.22	3 (30%)
2	SEP	B	346	2	8,9,10	0.80	0	7,12,14	2.36	3 (42%)
2	TPO	B	345	2	8,10,11	1.04	0	10,14,16	1.36	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	342	1	-	1/9/11/13	-
1	SEP	A	346	1	-	3/6/8/10	-
1	TPO	A	345	1	-	2/9/11/13	-
2	SEP	B	346	2	-	0/6/8/10	-
2	TPO	B	345	2	-	3/9/11/13	-

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	346	SEP	OG-CB-CA	12.31	120.12	108.14
2	B	346	SEP	OG-CB-CA	4.86	112.88	108.14
1	A	345	TPO	O3P-P-OG1	4.85	124.74	105.85
1	A	346	SEP	OG-P-O1P	3.38	115.57	106.44
2	B	346	SEP	O3P-P-OG	2.95	114.36	106.67
2	B	345	TPO	O3P-P-OG1	2.91	117.20	105.85
1	A	345	TPO	O3P-P-O2P	-2.53	98.33	107.80
1	A	345	TPO	O3P-P-O1P	-2.40	101.49	110.83
1	A	342	TPO	CG2-CB-CA	-2.28	108.83	113.26
2	B	346	SEP	OG-P-O1P	2.27	112.56	106.44
1	A	342	TPO	O2P-P-OG1	2.20	114.41	105.85

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	345	TPO	N-CA-CB-OG1
1	A	345	TPO	O-C-CA-CB
1	A	346	SEP	N-CA-CB-OG
1	A	346	SEP	C-CA-CB-OG
2	B	345	TPO	N-CA-CB-OG1
2	B	345	TPO	O-C-CA-CB
1	A	346	SEP	CB-OG-P-O2P
1	A	342	TPO	CA-CB-OG1-P
2	B	345	TPO	CB-OG1-P-O1P

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	345	TPO	1	0
2	B	345	TPO	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	B	501	-	4,4,4	0.25	0	6,6,6	0.16	0
3	CJN	B	502	-	23,24,24	1.16	2 (8%)	25,34,34	2.34	11 (44%)
3	CJN	A	501	-	23,24,24	1.24	3 (13%)	25,34,34	2.64	8 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CJN	B	502	-	-	2/12/22/22	0/3/3/3
3	CJN	A	501	-	-	0/12/22/22	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	CJN	C11-N12	2.77	1.39	1.35
3	A	501	CJN	C11-N12	2.57	1.39	1.35
3	A	501	CJN	C7-C6	2.51	1.46	1.42
3	A	501	CJN	C24-N4	2.04	1.42	1.38
3	B	502	CJN	C24-C7	2.03	1.43	1.41

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	CJN	C9-N8-C7	7.88	122.63	115.72
3	A	501	CJN	N10-C9-N8	-6.37	118.93	128.58
3	B	502	CJN	N10-C9-N8	-5.46	120.31	128.58
3	B	502	CJN	C9-N8-C7	5.40	120.45	115.72
3	B	502	CJN	C22-N21-C17	4.50	120.13	112.39
3	A	501	CJN	C7-C24-C11	-4.27	114.60	119.59
3	A	501	CJN	C22-N21-C17	3.62	118.61	112.39
3	B	502	CJN	C7-C24-C11	-3.19	115.86	119.59
3	B	502	CJN	C16-C15-C13	-3.12	108.07	111.49
3	A	501	CJN	C16-C15-C13	-3.01	108.19	111.49
3	B	502	CJN	C9-N10-C11	2.66	124.05	115.24
3	A	501	CJN	C9-N10-C11	2.60	123.87	115.24
3	B	502	CJN	C6-C5-N4	2.42	112.19	110.31
3	B	502	CJN	C11-N12-C13	-2.37	120.18	124.18
3	A	501	CJN	C3-C2-C1	-2.09	106.78	112.46
3	A	501	CJN	C2-N4-C5	-2.04	121.63	124.63
3	B	502	CJN	C23-N21-C17	2.03	115.88	112.39
3	B	502	CJN	C1-C2-N4	2.01	113.59	110.27
3	B	502	CJN	C3-C2-C1	-2.01	107.00	112.46

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	502	CJN	C16-C17-N21-C23
3	B	502	CJN	C19-C17-N21-C23

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand CJN B 502



Ligand CJN A 501



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	178:ASN	C	179:ASN	N	2.17

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/295 (92%)	1.76	100 (36%) 1 0	32, 64, 161, 194	0
2	B	277/295 (93%)	1.67	89 (32%) 1 1	29, 62, 109, 145	0
All	All	551/590 (93%)	1.71	189 (34%) 1 0	29, 63, 139, 194	0

All (189) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	170	PHE	6.9
1	A	205	VAL	6.8
1	A	165	PHE	6.5
2	B	258	LEU	6.3
1	A	168	PHE	5.7
2	B	199	VAL	5.7
2	B	187	VAL	5.6
2	B	180	PHE	5.5
2	B	170	PHE	5.2
1	A	198	GLY	5.1
2	B	176	VAL	4.9
2	B	185	ILE	4.8
2	B	230	PHE	4.8
2	B	214	LYS	4.7
1	A	343	VAL	4.7
2	B	259	CYS	4.7
1	A	260	LEU	4.7
1	A	237	MET	4.6
1	A	248	LEU	4.6
1	A	439	GLU	4.6
2	B	226	LEU	4.6
1	A	251	PHE	4.5
1	A	216	ALA	4.4
1	A	175	ASN	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	189	GLY	4.4
1	A	171	TYR	4.4
1	A	176	VAL	4.4
2	B	343	VAL	4.4
1	A	185	ILE	4.3
2	B	190	ASN	4.3
1	A	206	ASN	4.3
2	B	201	TYR	4.2
1	A	330	PHE	4.2
1	A	245	LEU	4.2
1	A	199	VAL	4.2
1	A	263	VAL	4.2
1	A	173	LEU	4.2
1	A	212	VAL	4.2
2	B	179	ASN	4.2
1	A	204	TYR	4.2
1	A	230	PHE	4.2
1	A	187	VAL	4.1
2	B	253	SER	4.1
2	B	252	SER	4.1
2	B	336	SER	4.1
2	B	196	GLY	4.0
2	B	257	ASP	4.0
2	B	348	ILE	4.0
2	B	410	ILE	4.0
1	A	262	TYR	3.9
2	B	186	SER	3.9
2	B	172	GLU	3.9
1	A	254	ASP	3.8
1	A	453	LEU	3.8
2	B	173	LEU	3.8
1	A	229	GLN	3.7
2	B	349	VAL	3.7
2	B	168	PHE	3.7
2	B	206	ASN	3.6
1	A	349	VAL	3.6
1	A	335	ALA	3.6
1	A	166	HIS	3.6
1	A	200	VAL	3.6
1	A	227	LYS	3.5
1	A	167	SER	3.5
2	B	198	GLY	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	375	VAL	3.4
1	A	186	SER	3.4
2	B	344	MET	3.4
1	A	331	GLY	3.4
2	B	204	TYR	3.4
1	A	450	VAL	3.4
1	A	180	PHE	3.4
1	A	192	MET	3.3
1	A	226	LEU	3.3
1	A	258	LEU	3.3
2	B	177	THR	3.3
2	B	197	PHE	3.3
2	B	229	GLN	3.3
1	A	208	THR	3.2
1	A	344	MET	3.2
2	B	335	ALA	3.2
2	B	188	GLY	3.2
1	A	179	ASN	3.2
1	A	249	LEU	3.2
1	A	223	THR	3.2
1	A	259	CYS	3.1
1	A	174	LYS	3.1
1	A	234	ILE	3.1
1	A	190	ASN	3.1
1	A	164	ARG	3.1
1	A	348	ILE	3.0
1	A	236	VAL	3.0
2	B	397	LEU	3.0
1	A	390	HIS	3.0
2	B	251	PHE	3.0
2	B	212	VAL	3.0
2	B	192	MET	3.0
2	B	373	PHE	2.9
2	B	169	SER	2.9
2	B	178	ASN	2.9
1	A	177	THR	2.9
1	A	261	VAL	2.9
2	B	171	TYR	2.9
1	A	357	PRO	2.9
2	B	223	THR	2.9
2	B	203	GLY	2.8
2	B	215	LEU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	252	SER	2.8
1	A	410	ILE	2.8
1	A	240	CYS	2.8
1	A	329	ASP	2.8
2	B	195	GLY	2.8
1	A	172	GLU	2.8
2	B	227	LYS	2.8
2	B	205	VAL	2.7
2	B	234	ILE	2.7
1	A	215	LEU	2.7
2	B	332	LEU	2.7
1	A	209	THR	2.7
1	A	246	VAL	2.7
1	A	211	ALA	2.6
1	A	241	GLN	2.6
1	A	353	ALA	2.6
1	A	210	VAL	2.6
2	B	184	PRO	2.6
1	A	178	ASN	2.6
2	B	279	GLY	2.6
1	A	169	SER	2.6
2	B	455	GLN	2.6
1	A	257	ASP	2.6
1	A	279	GLY	2.6
2	B	240	CYS	2.5
1	A	362	GLY	2.5
1	A	239	LYS	2.5
1	A	332	LEU	2.5
1	A	183	ARG	2.5
2	B	167	SER	2.4
2	B	328	SER	2.4
1	A	201	TYR	2.4
2	B	181	ASP	2.4
1	A	356	ALA	2.4
2	B	238	ALA	2.4
2	B	200	VAL	2.4
2	B	231	ASP	2.4
2	B	364	ILE	2.4
2	B	327	ILE	2.3
2	B	228	GLN	2.3
2	B	395	LEU	2.3
1	A	253	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	361	ARG	2.3
2	B	419	ASN	2.3
2	B	193	GLY	2.3
1	A	233	GLU	2.3
1	A	312	ILE	2.3
1	A	207	ASN	2.3
2	B	374	GLY	2.3
2	B	390	HIS	2.3
1	A	326	LYS	2.3
2	B	207	ASN	2.2
2	B	166	HIS	2.2
2	B	352	THR	2.2
2	B	175	ASN	2.2
1	A	434	SER	2.2
2	B	370	ILE	2.2
1	A	184	PRO	2.2
2	B	232	GLN	2.2
1	A	328	SER	2.2
2	B	183	ARG	2.1
2	B	236	VAL	2.1
1	A	222	THR	2.1
1	A	333	ALA	2.1
2	B	237	MET	2.1
1	A	242	HIS	2.1
2	B	375	VAL	2.1
2	B	241	GLN	2.1
2	B	191	LYS	2.1
1	A	327	ILE	2.1
2	B	249	LEU	2.1
2	B	330	PHE	2.1
1	A	247	GLU	2.1
1	A	334	ARG	2.1
2	B	347	ARG	2.1
1	A	194	GLU	2.0
2	B	359	ALA	2.0
1	A	403	ILE	2.0
2	B	440	LYS	2.0
1	A	423	SER	2.0
1	A	360	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	A	346	10/11	0.30	0.16	128,133,140,141	4
2	SEP	B	346	10/11	0.30	0.14	141,150,162,162	0
1	TPO	A	342	11/12	0.58	0.18	116,120,125,128	4
2	TPO	B	345	11/12	0.60	0.16	129,138,143,144	0
1	TPO	A	345	11/12	0.76	0.15	121,127,133,135	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CJN	A	501	22/22	0.75	0.17	34,45,47,51	0
4	SO4	B	501	5/5	0.78	0.12	92,94,95,96	0
3	CJN	B	502	22/22	0.85	0.13	28,35,44,45	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around CJN A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around CJN B 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.