



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 02:46 AM UTC

PDB ID : 1F4Y / pdb_00001f4y
Title : CRYSTAL STRUCTURE OF AN ANTI-CARBOHYDRATE ANTIBODY
DIRECTED AGAINST VIBRIO CHOLERAЕ O1 IN COMPLEX WITH
ANTIGEN
Authors : Alzari, P.M.; Souchon, H.
Deposited on : 2000-06-10
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	210	17% 42% 33% 8%
2	H	216	18% 37% 32% 13%
3	A	2	100%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3231 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ANTIBODY S-20-4, FAB FRAGMENT, LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	210	Total	C	N	O	S	0	0	0
			1578	989	262	321	6			

- Molecule 2 is a protein called ANTIBODY S-20-4, FAB FRAGMENT, HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	216	Total	C	N	O	S	0	0	0
			1616	1022	262	324	8			

- Molecule 3 is an oligosaccharide called 4,6-dideoxy-4-{[(2R)-2,4-dihydroxybutanoyl]amino}-2-O-methyl- α -D-mannopyranose-(1-2)-methyl 4,6-dideoxy-4-{[(2R)-2,4-dihydroxybutanoyl]amino}- α -D-mannopyranoside.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	2	Total	C	N	O		0	0	0
			37	22	2	13				

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: ANTIBODY S-20-4, FAB FRAGMENT, LIGHT CHAIN



- Molecule 2: ANTIBODY S-20-4, FAB FRAGMENT, HEAVY CHAIN



- Molecule 3: 4,6-dideoxy-4-[(2R)-2,4-dihydroxybutanoyl]amino}-2-O-methyl-alpha-D-mannopyranose-(1-2)-methyl 4,6-dideoxy-4-[(2R)-2,4-dihydroxybutanoyl]amino}-alpha-D-mannopyranoside



ZD01
ZC22

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.69Å 113.36Å 45.58Å 90.00° 93.93° 90.00°	Depositor
Resolution (Å)	10.00 – 2.80	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-2.80)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.226 , 0.352	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3231	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZD0, ZCZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	1.18	0/1615	3.06	203/2208 (9.2%)
2	H	1.25	3/1659 (0.2%)	3.12	206/2272 (9.1%)
All	All	1.21	3/3274 (0.1%)	3.09	409/4480 (9.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	2
2	H	0	1
All	All	0	3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	215	VAL	C-O	-5.60	1.17	1.24
2	H	98	ARG	CD-NE	-5.44	1.38	1.46
2	H	216	PRO	N-CD	5.13	1.54	1.47

The worst 5 of 409 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	98	ARG	CD-NE-CZ	33.66	171.52	124.40
1	L	56	ARG	CD-NE-CZ	17.67	149.14	124.40
1	L	55	ASN	CA-CB-CG	15.43	128.03	112.60
1	L	190	ARG	N-CA-C	14.21	126.46	110.97
2	H	2	VAL	N-CA-C	13.72	127.87	109.21

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	209	LYS	Mainchain
1	L	117	SER	Mainchain
1	L	174	LYS	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1578	0	1523	92	0
2	H	1616	0	1554	108	0
3	A	37	0	0	7	0
All	All	3231	0	3077	203	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 203 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1:ZD0:C5	3:A:1:ZD0:C1M	2.02	1.35
3:A:1:ZD0:C1M	3:A:1:ZD0:C3	2.21	1.17
1:L:41:LYS:HD2	1:L:42:PRO:HD2	1.42	1.02
3:A:1:ZD0:C1M	3:A:1:ZD0:C4	2.40	0.99
2:H:102:ALA:HA	3:A:2:ZCZ:C2M	1.96	0.95

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	208/210 (99%)	178 (86%)	20 (10%)	10 (5%)	2	6
2	H	214/216 (99%)	179 (84%)	22 (10%)	13 (6%)	1	3
All	All	422/426 (99%)	357 (85%)	42 (10%)	23 (6%)	1	4

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	43	ASP
1	L	53	THR
1	L	103	GLY
2	H	29	PHE
2	H	62	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	176/177 (99%)	134 (76%)	42 (24%)	1	3
2	H	182/186 (98%)	124 (68%)	58 (32%)	0	1
All	All	358/363 (99%)	258 (72%)	100 (28%)	0	1

5 of 100 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	40	THR
2	H	93	MET
2	H	214	ILE
2	H	47	TRP
2	H	74	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	39	GLN
2	H	82	GLN
2	H	203	HIS
2	H	175	GLN
2	H	3	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	ZD0	A	1	3	19,19,19	0.91	1 (5%)	22,26,26	3.15	11 (50%)
3	ZCZ	A	2	3	18,18,19	1.22	1 (5%)	17,24,26	4.07	9 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ZD0	A	1	3	-	6/13/33/33	0/1/1/1
3	ZCZ	A	2	3	-	9/13/30/33	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2	ZCZ	C4-N4	3.78	1.51	1.45
3	A	1	ZD0	C4-N4	2.64	1.50	1.45

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2	ZCZ	O7-C7-N4	-9.56	105.85	122.96
3	A	2	ZCZ	C4-N4-C7	-9.48	108.78	123.20
3	A	1	ZD0	C4-N4-C7	-8.26	110.63	123.20
3	A	1	ZD0	C1-O5-C5	-5.18	104.80	113.63
3	A	2	ZCZ	O5-C1-C2	5.02	119.68	109.51

There are no chirality outliers.

5 of 15 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1	ZD0	N4-C7-C8-C9
3	A	1	ZD0	N4-C7-C8-O8
3	A	1	ZD0	O7-C7-C8-C9
3	A	1	ZD0	O8-C8-C9-C10
3	A	2	ZCZ	C3-C2-O2-C2M

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2	ZCZ	4	0
3	A	1	ZD0	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.