



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:03 PM UTC

PDB ID : 4F4K / pdb_00004f4k
Title : DNA Polymerase I Large Fragment Complex 6
Authors : Wang, W.; Beese, L.S.
Deposited on : 2012-05-10
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	592	3% 89% 9% .
1	D	592	2% 91% 7% .
2	B	9	78% 22%
2	E	9	78% 11% 11%
3	C	13	69% 15% 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	13	 A horizontal bar chart showing the quality of chain F. The bar is divided into three segments: a green segment representing 62%, a yellow segment representing 31%, and a grey segment representing 8%. The percentages are labeled below the corresponding segments.

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 21725 atoms, of which 10028 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	579	Total	C	H	N	O	S	0	6	0
			9450	2979	4768	814	872	17			
1	D	579	Total	C	H	N	O	S	0	10	0
			9484	2992	4787	810	877	18			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	598	ALA	ASP	engineered mutation	UNP Q5KWC1
A	710	TYR	PHE	engineered mutation	UNP Q5KWC1
A	823	HIS	ARG	SEE REMARK 999	UNP Q5KWC1
D	598	ALA	ASP	engineered mutation	UNP Q5KWC1
D	710	TYR	PHE	engineered mutation	UNP Q5KWC1
D	823	HIS	ARG	SEE REMARK 999	UNP Q5KWC1

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(DDG))-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	9	Total	C	H	N	O	P	0	0	0
			277	86	100	31	52	8			
2	E	9	Total	C	H	N	O	P	0	0	0
			277	86	100	31	52	8			

- Molecule 3 is a DNA chain called DNA (5'-D(*CP*AP*TP*TP*CP*GP*AP*GP*TP*CP*AP*GP*G)-3').


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	11	Total	C	H	N	O	P	0	0	0
			352	108	124	42	67	11			
3	F	12	Total	C	H	N	O	P	0	0	0
			381	118	135	47	70	11			

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is 2'-3'-DIDEOXYGUANOSINE-5'-TRIPHOSPHATE (CCD ID: DG3) (formula: C₁₀H₁₆N₅O₁₂P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	D	1	Total	C	N	O	P	0	0
			30	10	5	12	3		

- Molecule 6 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	D	1	Total	Mn	0	0
			1	1		

- Molecule 7 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	D	1	Total	C	H	O	0	0
			22	6	14	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	493	Total	O	0	0
			493	493		
8	D	735	Total	O	0	0
			735	735		
8	B	37	Total	O	0	0
			37	37		
8	C	60	Total	O	0	0
			60	60		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	E	34	Total 34	O 34	0	0
8	F	77	Total 77	O 77	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase




- Molecule 1: DNA polymerase



- Molecule 2: DNA (5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(DDG))-3')



- Molecule 2: DNA (5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(DDG))-3')




- Molecule 3: DNA (5'-D(*CP*AP*TP*TP*CP*GP*AP*GP*TP*CP*AP*GP*G)-3')

Chain C:  69% 15% 15%



- Molecule 3: DNA (5'-D(*CP*AP*TP*TP*CP*GP*AP*GP*TP*CP*AP*GP*G)-3')

Chain F:  62% 31% 8%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	93.81Å 109.22Å 150.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.06 – 1.60 30.06 – 1.60	Depositor EDS
% Data completeness (in resolution range)	88.2 (30.06-1.60) 88.2 (30.06-1.60)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.06 (at 1.60Å)	Xtriage
Refinement program	PHENIX dev_1026	Depositor
R, R_{free}	0.171 , 0.194 0.172 , 0.195	Depositor DCC
R_{free} test set	8168 reflections (4.55%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 48.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.75$, $\langle L^2 \rangle = 0.66$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	21725	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, DDG, MN, MPD, DG3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/4784	0.73	0/6464
1	D	0.72	0/4812	0.80	0/6503
2	B	0.56	0/173	1.22	1/264 (0.4%)
2	E	0.56	0/173	1.23	1/264 (0.4%)
3	C	0.52	0/255	1.26	0/392
3	F	0.58	0/276	1.16	1/425 (0.2%)
All	All	0.63	0/10473	0.82	3/14312 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	26	DC	N1-C1'-C2'	-7.28	102.58	113.50
2	B	26	DC	N1-C1'-C2'	-5.64	105.03	113.50
3	F	8	DT	O4'-C1'-N1	-5.31	100.44	108.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4682	4768	4757	31	0
1	D	4697	4787	4775	32	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	177	100	103	2	0
2	E	177	100	103	2	0
3	C	228	124	125	2	0
3	F	246	135	137	2	0
4	A	5	0	0	0	0
4	D	10	0	0	0	0
5	D	30	0	12	3	0
6	D	1	0	0	0	0
7	D	8	14	14	0	0
8	A	493	0	0	11	0
8	B	37	0	0	3	0
8	C	60	0	0	2	0
8	D	735	0	0	14	0
8	E	34	0	0	0	0
8	F	77	0	0	0	0
All	All	11697	10028	10026	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:416:MET:HE3	8:D:1689:HOH:O	1.63	0.97
5:D:901:DG3:HN1	3:F:3:DT:H3	1.12	0.96
2:B:27:DT:OP2	8:B:230:HOH:O	1.90	0.88
1:D:812[A]:ASN:OD1	8:D:1178:HOH:O	1.91	0.88
2:B:27:DT:OP1	8:B:237:HOH:O	1.91	0.87
3:C:3:DT:O4	8:C:151:HOH:O	1.94	0.83
1:D:340:GLU:OE2	8:D:1702:HOH:O	2.06	0.72
1:D:305:ASP:OD1	8:D:1704:HOH:O	2.11	0.68
1:A:860:VAL:O	8:A:1402:HOH:O	2.13	0.66
1:D:764[B]:THR:HG21	8:D:1596:HOH:O	1.97	0.64
1:A:655:SER:O	1:A:660:ARG:NH1	2.30	0.63
1:D:658:GLU:CD	5:D:901:DG3:H2'1	2.25	0.60
1:D:814:ARG:NH2	1:D:847[A]:LEU:HD13	2.18	0.59
1:A:323:VAL:HG12	1:A:435:ARG:NH1	2.18	0.58
1:D:831:GLU:OE2	8:D:1102:HOH:O	2.17	0.58
1:D:456:GLU:HG3	8:D:1722:HOH:O	2.04	0.57
1:A:812[B]:ASN:ND2	8:A:1433:HOH:O	2.33	0.55
1:D:754:VAL:HG12	1:D:758:LYS:HE2	1.88	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:754:VAL:CG1	1:D:758:LYS:HE2	2.38	0.54
1:D:774:PRO:HA	8:D:1654:HOH:O	2.08	0.53
1:A:677:ARG:HB2	1:A:679:LEU:HG	1.89	0.52
1:A:463:ASP:O	1:A:467[B]:ARG:HG3	2.09	0.52
1:D:459:ARG:NH2	8:D:1503:HOH:O	2.43	0.51
1:D:713:VAL:HG12	1:D:739[B]:TYR:OH	2.10	0.51
1:D:435:ARG:NH2	8:D:1347:HOH:O	2.43	0.51
1:D:581:GLY:HA3	8:D:1323:HOH:O	2.11	0.51
1:A:381:LYS:HE3	8:A:1474:HOH:O	2.10	0.50
1:D:713:VAL:CG1	1:D:739[B]:TYR:OH	2.60	0.50
1:A:665:ILE:HG22	1:A:749:TYR:CE1	2.47	0.50
1:D:726:ASN:HB2	8:D:1466:HOH:O	2.12	0.49
1:A:315:LYS:NZ	8:A:1430:HOH:O	2.39	0.49
1:A:559:ASP:OD2	1:A:563:LYS:HE2	2.11	0.49
1:A:517:ARG:NH1	1:A:520:GLU:OE1	2.46	0.49
1:D:764[B]:THR:HG23	1:D:769:ARG:O	2.13	0.48
1:D:515:GLU:HG2	1:D:519:TYR:CE2	2.49	0.47
1:A:468:ASN:HA	8:A:1363:HOH:O	2.12	0.47
1:A:325:GLU:OE1	8:A:1399:HOH:O	2.20	0.47
1:A:672:MET:HE3	1:A:859:ARG:HB3	1.95	0.47
1:D:503[B]:MET:HE1	1:D:635:LYS:O	2.15	0.47
1:D:578:ARG:HD3	2:E:26:DC:OP1	2.14	0.47
3:C:11:DG:OP1	8:C:158:HOH:O	2.21	0.46
1:A:631:GLU:OE2	1:A:635:LYS:NZ	2.43	0.45
1:A:848:VAL:HB	1:A:849:PRO:HD3	1.98	0.45
1:A:607[A]:ASN:OD1	8:A:1483:HOH:O	2.21	0.44
1:D:814:ARG:CZ	1:D:847[A]:LEU:CD1	2.95	0.44
1:A:782:ASN:O	1:A:786:PHE:HD2	2.00	0.44
1:A:682:HIS:CE1	1:A:706:LYS:HD3	2.54	0.43
1:D:324:GLU:HG2	8:D:1582:HOH:O	2.18	0.43
1:D:764[B]:THR:HG22	1:D:765:THR:N	2.33	0.43
1:A:503:MET:HE1	1:A:635:LYS:O	2.19	0.43
1:A:579:GLN:HG2	8:B:233:HOH:O	2.18	0.43
1:A:812[A]:ASN:CG	8:A:1433:HOH:O	2.62	0.43
1:D:690:PHE:O	1:D:691:GLN:C	2.62	0.43
1:A:692:VAL:HG21	1:A:701:MET:HE1	2.00	0.42
3:F:9:DC:H2''	3:F:10:DA:C8	2.53	0.42
1:D:764[B]:THR:HG22	1:D:765:THR:H	1.84	0.42
5:D:901:DG3:O4'	2:E:29:DDG:H2''	2.19	0.42
1:D:836:ALA:HB3	1:D:841:MET:CE	2.50	0.42
1:A:840:GLU:O	1:A:844:LEU:HD13	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:645:SER:O	1:D:646:ASP:HB2	2.20	0.42
1:A:748:ARG:HG3	8:A:1426:HOH:O	2.20	0.42
1:A:439:ASP:HB3	8:A:1485:HOH:O	2.20	0.41
1:A:803:ILE:HG12	1:A:856:VAL:HG21	2.01	0.41
1:D:383:LYS:HE3	8:D:1442:HOH:O	2.19	0.41
1:D:712:ILE:HA	1:D:716:ILE:HG22	2.02	0.41
1:D:690:PHE:O	1:D:692:VAL:HG13	2.21	0.41
1:A:459:ARG:HB3	1:A:460:PRO:HD3	2.03	0.41
1:A:719:TYR:O	1:A:723:GLN:HG2	2.21	0.41
1:A:739:TYR:CD2	1:A:739:TYR:C	2.99	0.41
1:A:828:VAL:HB	1:A:831:GLU:HG3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	583/592 (98%)	570 (98%)	12 (2%)	1 (0%)	43	24
1	D	587/592 (99%)	576 (98%)	11 (2%)	0	100	100
All	All	1170/1184 (99%)	1146 (98%)	23 (2%)	1 (0%)	48	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	628	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	501/507 (99%)	497 (99%)	4 (1%)	73	59
1	D	505/507 (100%)	500 (99%)	5 (1%)	68	50
All	All	1006/1014 (99%)	997 (99%)	9 (1%)	73	55

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	349	GLU
1	A	477	LEU
1	A	511	LEU
1	A	850	GLU
1	D	303	LEU
1	D	477	LEU
1	D	728	SER
1	D	812[A]	ASN
1	D	812[B]	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	418	GLN
1	A	524	GLN
1	A	724	ASN
1	D	625	ASN
1	D	867	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DDG	B	29	2,3	20,23,24	0.69	0	27,33,36	0.74	0
2	DDG	E	29	2,3	20,23,24	0.64	0	27,33,36	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DDG	B	29	2,3	-	0/7/18/19	0/3/3/3
2	DDG	E	29	2,3	-	0/7/18/19	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	29	DDG	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	MPD	D	905	-	7,7,7	0.32	0	9,10,10	0.41	0
4	SO4	D	904	-	4,4,4	0.28	0	6,6,6	0.05	0
5	DG3	D	901	6	31,32,32	1.98	3 (9%)	44,50,50	1.79	15 (34%)
4	SO4	D	903	-	4,4,4	0.28	0	6,6,6	0.46	0
4	SO4	A	901	-	4,4,4	0.30	0	6,6,6	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	MPD	D	905	-	-	3/5/5/5	-
5	DG3	D	901	6	-	6/22/31/31	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	901	DG3	O6-C6	8.33	1.39	1.23
5	D	901	DG3	C2-N2	3.96	1.43	1.34
5	D	901	DG3	C6-N1	-3.25	1.32	1.38

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	901	DG3	C5-C4-N3	-4.00	122.03	128.39
5	D	901	DG3	O4'-C1'-N9	3.22	113.58	107.86
5	D	901	DG3	C2'-C1'-N9	-2.94	106.83	112.40
5	D	901	DG3	C2-N3-C4	2.90	117.30	112.30
5	D	901	DG3	C2-N1-C6	-2.75	120.12	125.11
5	D	901	DG3	O2B-PB-O3B	2.74	114.68	107.27
5	D	901	DG3	O2G-PG-O3B	2.66	113.55	104.64
5	D	901	DG3	O4'-C1'-C2'	-2.54	103.41	106.41
5	D	901	DG3	O3B-PB-O1B	-2.51	103.17	110.70
5	D	901	DG3	N9-C8-N7	-2.41	108.93	113.40
5	D	901	DG3	C5-C6-N1	2.40	119.36	113.25
5	D	901	DG3	O3A-PB-O1B	-2.34	103.67	110.70
5	D	901	DG3	N9-C4-N3	2.27	130.50	125.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	901	DG3	C6-C5-N7	2.14	134.18	130.29
5	D	901	DG3	O6-C6-C5	-2.02	121.19	126.53

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	D	901	DG3	PB-O3B-PG-O2G
7	D	905	MPD	C2-C3-C4-C5
5	D	901	DG3	PG-O3B-PB-O1B
5	D	901	DG3	PB-O3A-PA-O2A
5	D	901	DG3	PB-O3B-PG-O1G
5	D	901	DG3	PB-O3A-PA-O1A
7	D	905	MPD	O2-C2-C3-C4
7	D	905	MPD	C2-C3-C4-O4
5	D	901	DG3	PG-O3B-PB-O2B

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	901	DG3	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	579/592 (97%)	0.20	17 (2%) 53 56	10, 35, 65, 95	6 (1%)
1	D	579/592 (97%)	-0.29	9 (1%) 70 74	6, 22, 42, 54	10 (1%)
2	B	8/9 (88%)	-0.20	0 100 100	20, 25, 43, 52	0
2	E	8/9 (88%)	-0.31	0 100 100	15, 23, 43, 53	0
3	C	11/13 (84%)	-0.13	0 100 100	17, 25, 42, 61	0
3	F	12/13 (92%)	-0.25	0 100 100	12, 24, 47, 61	0
All	All	1197/1228 (97%)	-0.05	26 (2%) 62 66	6, 28, 55, 95	16 (1%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	679	LEU	5.2
1	A	692	VAL	3.6
1	D	735	PHE	3.2
1	D	699	PRO	3.1
1	A	693	SER	3.1
1	A	670	ASN	3.0
1	A	714	TYR	2.9
1	A	710	TYR	2.9
1	D	704	GLN	2.8
1	A	816	LYS	2.8
1	A	697	VAL	2.8
1	D	703	ARG	2.8
1	D	689	ILE	2.8
1	A	669	ASP	2.6
1	A	298	LYS	2.5
1	A	695	ASP	2.4
1	A	846	ARG	2.4
1	A	819	ARG	2.4
1	A	842	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	695	ASP	2.3
1	A	690	PHE	2.3
1	D	707	ALA	2.2
1	A	703	ARG	2.2
1	D	782	ASN	2.2
1	A	860	VAL	2.1
1	D	687	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DDG	B	29	21/22	0.96	0.06	18,20,29,29	0
2	DDG	E	29	21/22	0.98	0.04	9,12,16,17	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	D	904	5/5	0.83	0.11	49,50,55,59	0
7	MPD	D	905	8/8	0.84	0.13	37,52,63,63	0
4	SO4	A	901	5/5	0.89	0.11	33,34,40,46	0
5	DG3	D	901	30/30	0.95	0.09	13,18,22,23	0
4	SO4	D	903	5/5	0.97	0.08	29,30,39,42	0
6	MN	D	902	1/1	0.99	0.16	17,17,17,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.