



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 04:38 AM UTC

PDB ID : 6F4Q / pdb_00006f4q
Title : Human JMJD5 (Q275C) in complex with Mn(II), NOG and RPS6-A138C (129-144) (complex-2)
Authors : Chowdhury, R.; Islam, M.S.; Schofield, C.J.
Deposited on : 2017-11-29
Resolution : 1.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1985 (1.14-1.10)
Clashscore	190562	2021 (1.14-1.10)
Ramachandran outliers	187476	1978 (1.14-1.10)
Sidechain outliers	187428	1974 (1.14-1.10)
RSRZ outliers	180081	1984 (1.14-1.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	255	7% 83% 9% 8%
2	B	16	12% 12% 88%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4774 atoms, of which 2165 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called JmjC domain-containing protein 5.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	235	Total	C	H	N	O	S	0	78	0
			4306	1404	2118	370	407	7			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	162	HIS	-	expression tag	UNP Q8N371
A	163	HIS	-	expression tag	UNP Q8N371
A	164	HIS	-	expression tag	UNP Q8N371
A	165	HIS	-	expression tag	UNP Q8N371
A	166	HIS	-	expression tag	UNP Q8N371
A	167	HIS	-	expression tag	UNP Q8N371
A	168	SER	-	expression tag	UNP Q8N371
A	169	SER	-	expression tag	UNP Q8N371
A	170	GLY	-	expression tag	UNP Q8N371
A	171	VAL	-	expression tag	UNP Q8N371
A	172	ASP	-	expression tag	UNP Q8N371
A	173	LEU	-	expression tag	UNP Q8N371
A	174	GLY	-	expression tag	UNP Q8N371
A	175	THR	-	expression tag	UNP Q8N371
A	176	GLU	-	expression tag	UNP Q8N371
A	177	ASN	-	expression tag	UNP Q8N371
A	178	LEU	-	expression tag	UNP Q8N371
A	179	TYR	-	expression tag	UNP Q8N371
A	180	PHE	-	expression tag	UNP Q8N371
A	181	GLN	-	expression tag	UNP Q8N371
A	182	SER	-	expression tag	UNP Q8N371
A	217	ALA	CYS	engineered mutation	UNP Q8N371
A	232	ALA	CYS	engineered mutation	UNP Q8N371
A	275	CYS	GLN	engineered mutation	UNP Q8N371
A	295	ALA	CYS	engineered mutation	UNP Q8N371
A	384	ALA	CYS	engineered mutation	UNP Q8N371

- Molecule 2 is a protein called 40S ribosomal protein S6.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	2	Total	C	H	N	O	S	0	2	0
			33	9	16	5	2	1			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	138	CYS	ALA	engineered mutation	UNP P62753

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		

- Molecule 4 is N-OXALYLGLYCINE (CCD ID: OGA) (formula: C₄H₅NO₅).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	H	N	O		0	0
			13	4	3	1	5			

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
5	A	1	Total	C	H	N	O	0	0
			20	4	12	1	3		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			14	3	8	3		
6	A	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	360	Total 372	O 372	0	25
7	B	1	Total 1	O 1	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: JmjC domain-containing protein 5



- Molecule 2: 40S ribosomal protein S6



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.41Å 65.07Å 77.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.91 – 1.12 49.91 – 1.12	Depositor EDS
% Data completeness (in resolution range)	98.7 (49.91-1.12) 98.7 (49.91-1.12)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 1.12Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.139 , 0.153 (Not available) , 0.164	Depositor DCC
R_{free} test set	4770 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	12.8	Xtriage
Anisotropy	0.254	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4774	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, MN, OGA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/2443	0.65	0/3325
2	B	0.39	0/16	0.57	0/19
All	All	0.47	0/2459	0.65	0/3344

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2188	2118	1901	23	0
2	B	17	16	16	1	0
3	A	1	0	0	0	0
4	A	10	3	3	0	0
5	A	8	12	12	3	0
6	A	12	16	16	1	0
7	A	372	0	0	12	0
7	B	1	0	0	0	0
All	All	2609	2165	1948	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:503:TRS:H32	7:A:636:HOH:O	1.53	1.06
1:A:359[A]:ASN:HB3	7:A:794:HOH:O	1.76	0.85
1:A:364[A]:ASP:OD1	7:A:601[A]:HOH:O	1.97	0.80
1:A:253[B]:MET:HE2	1:A:257:GLU:HB3	1.71	0.72
1:A:182[B]:SER:N	7:A:608:HOH:O	2.34	0.60
1:A:265[A]:ASN:O	1:A:267[A]:PRO:HD3	2.05	0.56
1:A:275[B]:CYS:HB3	1:A:310:TRP:CH2	2.42	0.55
1:A:265[A]:ASN:O	1:A:267[A]:PRO:CD	2.54	0.55
1:A:263[A]:ILE:HG23	7:A:882:HOH:O	2.07	0.54
1:A:229[B]:ILE:HG21	1:A:285:LEU:HD21	1.91	0.53
2:B:137[A]:ARG:O	2:B:138[A]:CYS:HB2	2.10	0.50
1:A:201:VAL:O	5:A:503:TRS:O2	2.26	0.49
1:A:196[B]:ARG:CZ	7:A:701:HOH:O	2.62	0.47
1:A:358[A]:HIS:CE1	7:A:794:HOH:O	2.67	0.47
1:A:299[A]:ASP:OD2	7:A:602:HOH:O	2.20	0.46
5:A:503:TRS:C3	7:A:636:HOH:O	2.34	0.45
1:A:253[B]:MET:HE2	1:A:257:GLU:CB	2.45	0.44
1:A:197[B]:GLU:HG3	1:A:198[B]:GLN:OE1	2.17	0.44
1:A:196[A]:ARG:NH2	7:A:612:HOH:O	2.47	0.44
1:A:197[B]:GLU:HG3	1:A:198[B]:GLN:CD	2.43	0.43
1:A:189[B]:ARG:HD2	7:A:869:HOH:O	2.17	0.43
1:A:188:HIS:NE2	1:A:189[B]:ARG:NH1	2.65	0.42
1:A:189[B]:ARG:HG3	6:A:505:GOL:H12	2.01	0.42
1:A:372[B]:LYS:NZ	7:A:610:HOH:O	2.40	0.42
1:A:333[B]:MET:O	1:A:408:SER:HA	2.20	0.41
1:A:194:HIS:CE1	1:A:198[A]:GLN:HG3	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/255 (113%)	281 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/229 (112%)	257 (100%)	0	100	100
2	B	2/15 (13%)	2 (100%)	0	100	100
All	All	259/244 (106%)	259 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	265[A]	ASN
1	A	325	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	A	504	-	5,5,5	1.31	1 (20%)	5,5,5	1.00	0
4	OGA	A	502	3	9,9,9	2.17	4 (44%)	8,11,11	1.42	0
6	GOL	A	505	-	5,5,5	1.02	0	5,5,5	1.28	1 (20%)
5	TRS	A	503	-	7,7,7	1.44	1 (14%)	9,9,9	2.08	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	504	-	-	2/4/4/4	-
4	OGA	A	502	3	-	0/8/9/9	-
6	GOL	A	505	-	-	2/4/4/4	-
5	TRS	A	503	-	-	6/9/9/9	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	502	OGA	C2-C1	-3.86	1.49	1.54
5	A	503	TRS	C2-C	3.26	1.61	1.53
4	A	502	OGA	O2'-C2	2.88	1.29	1.23
6	A	504	GOL	C1-C2	2.31	1.60	1.51
4	A	502	OGA	O4-C5	2.29	1.29	1.22
4	A	502	OGA	O3-C5	-2.14	1.23	1.30

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	503	TRS	C2-C-C1	4.29	122.07	110.66
5	A	503	TRS	C3-C-C1	-3.98	100.07	110.66
6	A	505	GOL	C3-C2-C1	-2.60	102.27	111.80

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	503	TRS	N-C-C1-O1
5	A	503	TRS	N-C-C3-O3
6	A	504	GOL	C1-C2-C3-O3
6	A	505	GOL	O1-C1-C2-C3
5	A	503	TRS	C3-C-C1-O1
6	A	504	GOL	O2-C2-C3-O3
6	A	505	GOL	O1-C1-C2-O2
5	A	503	TRS	C2-C-C1-O1
5	A	503	TRS	C1-C-C3-O3
5	A	503	TRS	C2-C-C3-O3

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	505	GOL	1	0
5	A	503	TRS	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	235/255 (92%)	0.12	18 (7%)	19 26	6, 15, 32, 59	52 (22%)
2	B	2/16 (12%)	4.04	2 (100%)	0 0	24, 24, 24, 39	2 (100%)
All	All	237/271 (87%)	0.16	20 (8%)	17 23	6, 16, 33, 59	54 (22%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	138[A]	CYS	5.4
1	A	355	HIS	4.2
1	A	182[A]	SER	3.5
1	A	356	LEU	3.4
1	A	357	LEU	3.1
1	A	246[A]	GLU	3.0
1	A	353	ASP	3.0
1	A	183[A]	THR	2.9
1	A	214[A]	HIS	2.8
1	A	247[A]	GLU	2.7
2	B	137[A]	ARG	2.7
1	A	248[A]	TRP	2.6
1	A	265[A]	ASN	2.6
1	A	354	THR	2.5
1	A	358[A]	HIS	2.5
1	A	264[A]	VAL	2.4
1	A	219[A]	GLN	2.4
1	A	250[A]	GLN	2.4
1	A	267[A]	PRO	2.4
1	A	240	GLY	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	GOL	A	504	6/6	0.63	0.14	40,63,77,78	0
6	GOL	A	505	6/6	0.84	0.14	41,50,61,73	0
5	TRS	A	503	8/8	0.88	0.14	19,45,54,54	0
4	OGA	A	502	10/10	0.99	0.04	9,10,12,12	0
3	MN	A	501	1/1	1.00	0.01	9,9,9,9	0

6.5 Other polymers [i](#)

There are no such residues in this entry.