



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 01:50 PM UTC

PDB ID : 1F5F / pdb_00001f5f
Title : CRYSTAL STRUCTURE OF THE N-TERMINAL G-DOMAIN OF SHBG
IN COMPLEX WITH ZINC
Authors : Avvakumov, V.A.; Muller, Y.A.; Hammond, G.L.
Deposited on : 2000-06-14
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	205	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	IPA	A	254	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 1524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SEX HORMONE-BINDING GLOBULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	172	Total	C	N	O	S	0	5	0
			1366	864	243	251	8			


- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		

- Molecule 4 is 5-ALPHA-DIHYDROTESTOSTERONE (CCD ID: DHT) (formula: C₁₉H₃₀O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			21	19	2		

- Molecule 5 is ISOPROPYL ALCOHOL (CCD ID: IPA) (formula: C_3H_8O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	3	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	130	Total 130	O 130	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SEX HORMONE-BINDING GLOBULIN



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	104.02Å 104.02Å 84.71Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.70 20.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	96.3 (20.00-1.70) 96.3 (20.00-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 1.70Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.197 , 0.259 0.202 , 0.260	Depositor DCC
R_{free} test set	1868 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1524	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IPA, CA, ZN, DHT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	0/1420	1.93	38/1928 (2.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	11

There are no bond length outliers.

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	154	ARG	NE-CZ-NH1	-11.14	110.36	121.50
1	A	60	THR	N-CA-C	-9.42	101.03	114.12
1	A	71	LEU	CA-C-O	9.08	130.33	120.71
1	A	154	ARG	NH1-CZ-NH2	8.48	130.32	119.30
1	A	125[A]	ARG	CA-C-O	8.45	131.50	121.36
1	A	125[B]	ARG	CA-C-O	8.45	131.50	121.36
1	A	125[A]	ARG	O-C-N	-8.03	114.07	123.06
1	A	125[B]	ARG	O-C-N	-8.03	114.07	123.06
1	A	127	VAL	CA-C-N	-7.59	111.06	122.74
1	A	127	VAL	C-N-CA	-7.59	111.06	122.74
1	A	121	VAL	N-CA-C	-7.26	106.14	113.47
1	A	120	GLU	CA-C-N	-7.21	114.14	122.14
1	A	120	GLU	C-N-CA	-7.21	114.14	122.14
1	A	110	ASP	CA-C-O	-7.17	111.25	119.78
1	A	83	HIS	CA-CB-CG	-6.97	106.83	113.80
1	A	113	LEU	N-CA-CB	6.57	121.36	110.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	128	SER	CA-C-O	-6.49	113.12	120.32
1	A	72	ARG	CD-NE-CZ	6.49	133.48	124.40
1	A	67	PHE	CA-CB-CG	6.27	120.07	113.80
1	A	54	VAL	CA-C-O	-6.14	113.23	121.02
1	A	60	THR	N-CA-CB	6.06	119.25	110.88
1	A	118	GLY	CA-C-N	-6.04	114.66	123.00
1	A	118	GLY	C-N-CA	-6.04	114.66	123.00
1	A	160	ALA	CA-C-O	6.04	127.65	120.70
1	A	42	SER	CA-CB-OG	5.98	123.06	111.10
1	A	110	ASP	O-C-N	5.87	129.57	122.35
1	A	114	LEU	O-C-N	-5.70	116.66	123.27
1	A	114	LEU	CA-C-O	5.67	126.36	120.30
1	A	75	ARG	O-C-N	5.62	128.02	121.39
1	A	158	VAL	CA-C-O	5.58	122.44	119.15
1	A	138	ILE	O-C-N	5.45	128.94	123.10
1	A	110	ASP	N-CA-C	-5.42	106.54	112.72
1	A	65	ASP	CA-CB-CG	5.36	117.96	112.60
1	A	165	LEU	N-CA-CB	-5.32	101.60	111.13
1	A	140	ARG	CA-C-O	-5.20	115.20	120.71
1	A	128	SER	O-C-N	5.05	129.22	123.31
1	A	157	LEU	N-CA-CB	5.03	119.77	111.62
1	A	121	VAL	CA-C-O	5.01	123.49	118.98

There are no chirality outliers.

All (11) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	111	SER	Mainchain
1	A	164	CYS	Mainchain
1	A	169	SER	Mainchain
1	A	26	PRO	Mainchain
1	A	28	ALA	Mainchain
1	A	39	LYS	Mainchain
1	A	53	GLY	Mainchain
1	A	54	VAL	Mainchain
1	A	64	ASP	Mainchain
1	A	70	GLY	Mainchain
1	A	85	ALA	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1366	0	1347	37	0
2	A	1	0	0	0	0
3	A	2	0	0	0	0
4	A	21	0	30	0	0
5	A	4	0	8	1	0
6	A	130	0	0	7	0
All	All	1524	0	1385	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17[A]:HIS:CD2	1:A:187:SER:O	2.22	0.92
1:A:102:GLN:HE21	1:A:102:GLN:H	1.14	0.91
1:A:107[B]:MET:HE1	1:A:131:LEU:HG	1.55	0.87
1:A:17[B]:HIS:HD2	1:A:19:SER:H	1.23	0.85
1:A:17[A]:HIS:HD2	1:A:187:SER:O	1.60	0.85
1:A:17[B]:HIS:CD2	1:A:19:SER:H	2.02	0.77
1:A:107[A]:MET:HE1	1:A:131:LEU:HG	1.71	0.72
1:A:25:GLU:HB2	1:A:26:PRO:HD2	1.74	0.69
1:A:40:THR:CG2	1:A:131:LEU:HD12	2.30	0.61
1:A:40:THR:HG22	1:A:131:LEU:HD12	1.84	0.60
1:A:19:SER:HB2	6:A:367:HOH:O	2.00	0.59
1:A:34:LEU:HD23	1:A:136:HIS:O	2.02	0.59
1:A:43:SER:HB3	1:A:104:GLU:HG2	1.83	0.58
1:A:186:ARG:HD3	6:A:380:HOH:O	2.02	0.58
1:A:102:GLN:NE2	1:A:117:ASP:OD1	2.37	0.58
1:A:20:ASN:HB2	6:A:347:HOH:O	2.06	0.55
1:A:102:GLN:HE21	1:A:102:GLN:N	1.94	0.54
1:A:32:PHE:HB3	1:A:37:ILE:HD11	1.90	0.53
1:A:34:LEU:HA	1:A:37:ILE:HD12	1.91	0.52
1:A:22:PRO:HG2	6:A:360:HOH:O	2.10	0.51
1:A:20:ASN:HD21	1:A:186:ARG:HE	1.59	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:GLU:OE1	5:A:254:IPA:H32	2.12	0.49
1:A:45:GLU:OE2	1:A:167:ARG:NH1	2.45	0.48
1:A:119:GLU:HG2	6:A:316:HOH:O	2.15	0.47
1:A:107[B]:MET:CE	1:A:131:LEU:HG	2.39	0.46
1:A:87:LEU:HD13	1:A:89:VAL:HG23	1.98	0.45
1:A:21:GLY:HA3	1:A:22:PRO:HD3	1.80	0.45
1:A:123:ARG:HH11	1:A:123:ARG:HD3	1.67	0.44
1:A:40:THR:HG21	1:A:131:LEU:HD12	2.00	0.43
1:A:24:GLN:HB2	6:A:360:HOH:O	2.18	0.42
1:A:64:ASP:HB3	1:A:65:ASP:OD1	2.19	0.42
1:A:34:LEU:HD13	1:A:139:MET:HB2	2.00	0.42
1:A:23:GLY:HA2	6:A:366:HOH:O	2.19	0.42
1:A:107[A]:MET:HB3	1:A:107[A]:MET:HE3	1.71	0.42
1:A:157:LEU:C	1:A:157:LEU:HD12	2.46	0.41
1:A:37:ILE:HG13	1:A:175:ALA:CB	2.51	0.41
1:A:131:LEU:HD23	1:A:131:LEU:HA	1.87	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	173/205 (84%)	165 (95%)	8 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	152/176 (86%)	137 (90%)	15 (10%)	7 1

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	SER
1	A	34	LEU
1	A	36	LYS
1	A	37	ILE
1	A	39	LYS
1	A	63	LYS
1	A	102	GLN
1	A	107[A]	MET
1	A	107[B]	MET
1	A	131	LEU
1	A	157	LEU
1	A	173	LYS
1	A	183	THR
1	A	187	SER
1	A	188	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	102	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DHT	A	253	-	24,24,24	0.90	0	39,39,39	1.84	11 (28%)
5	IPA	A	254	-	3,3,3	0.53	0	3,3,3	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DHT	A	253	-	-	-	0/4/4/4

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	253	DHT	C10-C9-C8	4.22	116.72	112.43
4	A	253	DHT	C2-C1-C10	4.05	118.50	113.44
4	A	253	DHT	C2-C3-C4	3.61	121.36	115.89
4	A	253	DHT	C9-C10-C5	-3.44	103.74	108.51
4	A	253	DHT	C12-C13-C17	2.94	119.70	115.23
4	A	253	DHT	C11-C9-C10	-2.93	110.61	113.91
4	A	253	DHT	O3-C3-C4	-2.77	118.16	121.95
4	A	253	DHT	C16-C17-C13	2.76	106.70	104.52
4	A	253	DHT	C6-C7-C8	-2.54	108.10	112.16
4	A	253	DHT	C19-C10-C9	2.47	114.49	111.18
4	A	253	DHT	C6-C5-C4	2.37	114.48	111.17

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	254	IPA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	172/205 (83%)	0.70	14 (8%) 18 19	12, 32, 76, 90	5 (2%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	21	GLY	4.2
1	A	182	PRO	4.0
1	A	188	CYS	3.9
1	A	185	LEU	3.6
1	A	183	THR	3.5
1	A	22	PRO	3.4
1	A	181	ALA	3.3
1	A	131	LEU	2.8
1	A	20	ASN	2.8
1	A	130	PRO	2.7
1	A	186	ARG	2.1
1	A	164	CYS	2.1
1	A	23	GLY	2.1
1	A	63	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	IPA	A	254	4/4	0.67	0.46	39,41,42,43	4
4	DHT	A	253	21/21	0.94	0.07	18,20,26,30	0
3	ZN	A	251	1/1	0.96	0.07	40,40,40,40	0
2	CA	A	250	1/1	0.97	0.10	31,31,31,31	0
3	ZN	A	252	1/1	0.99	0.13	47,47,47,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.