



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 12:44 PM UTC

PDB ID : 9F5M / pdb_00009f5m
Title : Structure of the C-terminal domain of bacteriophage K gp155, native
Authors : Pichel, A.; van Raaij, M.J.
Deposited on : 2024-04-29
Resolution : 1.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5890 (1.54-1.50)
Clashscore	190562	6116 (1.54-1.50)
Ramachandran outliers	187476	6002 (1.54-1.50)
Sidechain outliers	187428	5999 (1.54-1.50)
RSRZ outliers	180081	5891 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	235	4% 93% 7%
1	BBB	235	4% 94% 6%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4457 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GP-PDE domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	234	Total	C	N	O	S	0	4	0
			1877	1195	303	368	11			
1	BBB	234	Total	C	N	O	S	0	8	0
			1896	1210	303	371	12			

- Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	AAA	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	BBB	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	C	O	0	0
			6	3	3		
3	BBB	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	311	Total	O	0	0
			311	311		
4	BBB	337	Total	O	0	0
			337	337		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GP-PDE domain-containing protein



- Molecule 1: GP-PDE domain-containing protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.13Å 43.33Å 98.98Å 90.00° 94.08° 90.00°	Depositor
Resolution (Å)	62.38 – 1.52 62.38 – 1.52	Depositor EDS
% Data completeness (in resolution range)	95.3 (62.38-1.52) 95.3 (62.38-1.52)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 1.52Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.188 , 0.213 0.197 , 0.222	Depositor DCC
R_{free} test set	4666 reflections (4.64%)	wwPDB-VP
Wilson B-factor (Å ²)	22.2	Xtriage
Anisotropy	0.076	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 32.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4457	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	0.97	0/1925	1.19	0/2598
1	BBB	0.95	0/1956	1.20	3/2639 (0.1%)
All	All	0.96	0/3881	1.19	3/5237 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	790	PHE	CA-C-N	5.96	128.75	120.29
1	BBB	790	PHE	C-N-CA	5.96	128.75	120.29
1	BBB	664	ASP	CA-CB-CG	5.05	117.65	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1877	0	1860	10	0
1	BBB	1896	0	1891	6	0
2	AAA	12	0	13	0	0
2	BBB	12	0	13	0	0
3	AAA	6	0	8	0	0
3	BBB	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	AAA	311	0	0	1	0
4	BBB	337	0	0	0	0
All	All	4457	0	3793	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:826:MET:HB2	1:AAA:848:ILE:HD11	1.87	0.56
1:AAA:653:LEU:HG	1:AAA:725[A]:MET:HG2	1.90	0.53
1:AAA:777:PHE:CE1	1:AAA:782:MET:HE2	2.48	0.49
1:BBB:848:ILE:HG22	1:BBB:848:ILE:O	2.13	0.48
1:AAA:750:ILE:HG21	1:AAA:761:ILE:HD13	1.96	0.48
1:AAA:725[A]:MET:HE2	1:AAA:725[A]:MET:HB3	1.72	0.47
1:AAA:683:SER:O	1:AAA:687[A]:THR:HG23	2.16	0.46
1:AAA:826:MET:CB	1:AAA:848:ILE:HD11	2.46	0.45
1:BBB:721:PHE:CD2	1:BBB:756:GLU:HG2	2.52	0.44
1:BBB:745:ARG:NH1	1:BBB:767:ASN:O	2.49	0.44
1:BBB:717:THR:HA	1:BBB:725[A]:MET:HE1	1.99	0.43
1:AAA:791:TYR:CZ	1:AAA:814:LYS:HG3	2.55	0.42
1:AAA:751:GLN:HA	1:AAA:771:ALA:O	2.21	0.41
1:AAA:736:LYS:NZ	4:AAA:1007:HOH:O	2.54	0.40
1:BBB:646:TYR:CE2	1:BBB:712:LYS:HB3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	236/235 (100%)	232 (98%)	4 (2%)	0	100	100
1	BBB	240/235 (102%)	234 (98%)	6 (2%)	0	100	100
All	All	476/470 (101%)	466 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	210/207 (101%)	209 (100%)	1 (0%)	81	66
1	BBB	214/207 (103%)	213 (100%)	1 (0%)	81	66
All	All	424/414 (102%)	422 (100%)	2 (0%)	81	66

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	837	PHE
1	BBB	837	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MES	BBB	901	-	12,12,12	0.70	0	15,16,16	0.38	0
3	GOL	AAA	902	-	5,5,5	0.11	0	5,5,5	0.25	0
2	MES	AAA	901	-	12,12,12	0.74	0	15,16,16	0.39	0
3	GOL	BBB	902	-	5,5,5	0.10	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MES	BBB	901	-	-	5/6/14/14	0/1/1/1
3	GOL	AAA	902	-	-	1/4/4/4	-
2	MES	AAA	901	-	-	3/6/14/14	0/1/1/1
3	GOL	BBB	902	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	AAA	901	MES	C7-C8-S-O1S
2	AAA	901	MES	C7-C8-S-O2S

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	BBB	901	MES	C7-C8-S-O2S
2	AAA	901	MES	C7-C8-S-O3S
2	BBB	901	MES	C7-C8-S-O3S
2	BBB	901	MES	C8-C7-N4-C5
3	AAA	902	GOL	O2-C2-C3-O3
2	BBB	901	MES	C7-C8-S-O1S
2	BBB	901	MES	C8-C7-N4-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	AAA	234/235 (99%)	0.43	9 (3%)	44	51	12, 23, 34, 50	4 (1%)
1	BBB	234/235 (99%)	0.39	9 (3%)	44	51	12, 22, 34, 52	8 (3%)
All	All	468/470 (99%)	0.41	18 (3%)	44	51	12, 22, 34, 52	12 (2%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	848	ILE	6.5
1	AAA	848	ILE	5.6
1	AAA	615	ASP	3.9
1	BBB	709	THR	3.8
1	AAA	709	THR	2.9
1	AAA	735	ALA	2.8
1	AAA	741	ILE	2.8
1	BBB	710	LYS	2.7
1	BBB	791	TYR	2.6
1	AAA	710	LYS	2.6
1	AAA	776	THR	2.5
1	BBB	615	ASP	2.4
1	BBB	797	TYR	2.3
1	BBB	756	GLU	2.3
1	AAA	756	GLU	2.2
1	AAA	742	GLY	2.1
1	BBB	708	GLY	2.1
1	BBB	725[A]	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	AAA	902	6/6	0.71	0.19	38,46,47,48	0
3	GOL	BBB	902	6/6	0.81	0.16	36,41,43,46	0
2	MES	BBB	901	12/12	0.92	0.13	34,40,42,42	0
2	MES	AAA	901	12/12	0.95	0.11	33,40,41,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.