



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 03:44 PM UTC

PDB ID : 4F6E / pdb_00004f6e
Title : Crystal Structure of the K182R, A183P mutant manganese superoxide dismutase from *Sacchomyces cerevisiae*
Authors : Sheng, Y.; Cascio, D.; Valentine, J.S.
Deposited on : 2012-05-14
Resolution : 1.60 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	 94% 6%
1	B	207	 97% ..
1	C	207	 91% 8%
1	D	207	 92% 8%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7238 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Superoxide dismutase [Mn], mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	207	1670	1084	279	305	2	0	0	0
1	B	205	1656	1073	276	305	2	0	1	0
1	C	207	1677	1089	279	307	2	0	2	0
1	D	206	1662	1078	278	304	2	0	0	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	ARG	LYS	engineered mutation	UNP P00447
A	183	PRO	ALA	engineered mutation	UNP P00447
B	182	ARG	LYS	engineered mutation	UNP P00447
B	183	PRO	ALA	engineered mutation	UNP P00447
C	182	ARG	LYS	engineered mutation	UNP P00447
C	183	PRO	ALA	engineered mutation	UNP P00447
D	182	ARG	LYS	engineered mutation	UNP P00447
D	183	PRO	ALA	engineered mutation	UNP P00447

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Mn 1	0	0
2	B	1	Total 1	Mn 1	0	0
2	C	1	Total 1	Mn 1	0	0
2	D	1	Total 1	Mn 1	0	0

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	C	1	8	4	1	3	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	D	1	6	3	3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	132	Total 132	O 132	0	0
5	B	142	Total 142	O 142	0	0
5	C	142	Total 142	O 142	0	0
5	D	139	Total 139	O 139	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Superoxide dismutase [Mn], mitochondrial

Chain A:  94% 6%



- Molecule 1: Superoxide dismutase [Mn], mitochondrial

Chain B:  97%



- Molecule 1: Superoxide dismutase [Mn], mitochondrial

Chain C:  91% 8%



- Molecule 1: Superoxide dismutase [Mn], mitochondrial

Chain D:  92% 8%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	65.54Å 66.25Å 66.62Å 112.58° 103.63° 110.27°	Depositor
Resolution (Å)	37.65 – 1.60 37.65 – 1.60	Depositor EDS
% Data completeness (in resolution range)	90.5 (37.65-1.60) 90.6 (37.65-1.60)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.93 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.7.3_928	Depositor
R, R_{free}	0.164 , 0.195 0.159 , 0.189	Depositor DCC
R_{free} test set	5252 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	13.9	Xtriage
Anisotropy	0.341	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for h+k+l,-h,-k 0.000 for -k,-l,h+k+l 0.009 for k,h,-h-k-l 0.010 for l,-h-k-l,h 0.008 for -l,-k,-h 0.013 for -h-k-l,l,k 0.008 for -h,h+k+l,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7238	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, TRS, MN, MLY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/1551	0.68	0/2129
1	B	0.40	0/1551	0.66	0/2129
1	C	0.39	0/1564	0.67	0/2147
1	D	0.38	0/1543	0.68	0/2118
All	All	0.39	0/6209	0.67	0/8523

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1670	0	1635	8	0
1	B	1656	0	1613	2	0
1	C	1677	0	1647	12	0
1	D	1662	0	1624	15	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	C	8	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	6	0	8	0	0
5	A	132	0	0	0	0
5	B	142	0	0	0	0
5	C	142	0	0	0	0
5	D	139	0	0	0	0
All	All	7238	0	6539	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 34 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:45:PHE:CE1	1:C:64:MET:HE1	2.35	0.61
1:B:65:ILE:O	1:B:68:GLN:HG3	2.06	0.56
1:C:44:GLN:HB3	1:C:67:ILE:HD13	1.89	0.55
1:C:149:VAL:HG21	1:C:157:VAL:HG11	1.88	0.54
1:A:182:ARG:HG2	1:A:183:PRO:HD3	1.89	0.53

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	191/207 (92%)	187 (98%)	4 (2%)	0	100	100
1	B	191/207 (92%)	187 (98%)	4 (2%)	0	100	100
1	C	193/207 (93%)	188 (97%)	5 (3%)	0	100	100
1	D	191/207 (92%)	186 (97%)	5 (3%)	0	100	100
All	All	766/828 (92%)	748 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/156 (100%)	155 (99%)	1 (1%)	78	66
1	B	156/156 (100%)	155 (99%)	1 (1%)	78	66
1	C	158/156 (101%)	155 (98%)	3 (2%)	50	27
1	D	155/156 (99%)	154 (99%)	1 (1%)	78	66
All	All	625/624 (100%)	619 (99%)	6 (1%)	68	50

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	69	GLN
1	C	87	ASN
1	D	87	ASN
1	B	87	ASN
1	A	87	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	32	GLN
1	D	32	GLN
1	D	68	GLN
1	D	46	GLN
1	B	69	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

59 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MLY	B	63	1	9,10,11	0.55	0	6,11,13	0.85	0
1	MLY	A	63	1	9,10,11	0.55	0	6,11,13	0.82	0
1	MLY	A	145	1	9,10,11	0.62	0	6,11,13	0.51	0
1	MLY	D	1	1	9,10,11	0.56	0	6,11,13	0.79	0
1	MLY	C	187	1	9,10,11	0.58	0	6,11,13	0.72	0
1	MLY	C	105	1	9,10,11	0.56	0	6,11,13	0.70	0
1	MLY	D	72	1	9,10,11	0.61	0	6,11,13	0.77	0
1	MLY	A	1	1	9,10,11	0.59	0	6,11,13	0.87	0
1	MLY	C	119	1	9,10,11	0.48	0	6,11,13	0.82	0
1	MLY	C	138	1	9,10,11	0.53	0	6,11,13	0.87	0
1	MLY	B	72	1	9,10,11	0.57	0	6,11,13	0.60	0
1	MLY	D	105	1	9,10,11	0.56	0	6,11,13	0.85	0
1	MLY	C	196	1	9,10,11	0.56	0	6,11,13	0.78	0
1	MLY	A	29	1	9,10,11	0.55	0	6,11,13	0.87	0
1	MLY	D	196	1	9,10,11	0.61	0	6,11,13	0.73	0
1	MLY	B	1	1	9,10,11	0.56	0	6,11,13	0.85	0
1	MLY	B	124	1	9,10,11	0.56	0	6,11,13	0.98	0
1	MLY	C	63	1	9,10,11	0.57	0	6,11,13	0.88	0
1	MLY	D	29	1	9,10,11	0.59	0	6,11,13	0.65	0
1	MLY	D	54	1	9,10,11	0.64	0	6,11,13	0.81	0
1	MLY	D	8	1	9,10,11	0.52	0	6,11,13	0.75	0
1	MLY	D	187	1	9,10,11	0.58	0	6,11,13	0.86	0
1	MLY	B	145	1	9,10,11	0.61	0	6,11,13	0.92	0
1	MLY	C	54	1	9,10,11	0.57	0	6,11,13	0.73	0
1	MLY	B	181	1	9,10,11	0.65	0	6,11,13	0.37	0
1	MLY	A	105	1	9,10,11	0.56	0	6,11,13	0.93	0
1	MLY	B	29	1	9,10,11	0.58	0	6,11,13	0.70	0
1	MLY	C	8	1	9,10,11	0.51	0	6,11,13	0.80	0
1	MLY	D	119	1	9,10,11	0.54	0	6,11,13	0.81	0
1	MLY	A	196	1	9,10,11	0.61	0	6,11,13	0.80	0
1	MLY	C	1	1	9,10,11	0.60	0	6,11,13	0.82	0
1	MLY	D	138	1	9,10,11	0.68	0	6,11,13	0.72	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MLY	C	29	1	9,10,11	0.60	0	6,11,13	0.70	0
1	MLY	A	54	1	9,10,11	0.55	0	6,11,13	0.82	0
1	MLY	B	8	1	9,10,11	0.45	0	6,11,13	0.75	0
1	MLY	A	187	1	9,10,11	0.60	0	6,11,13	0.81	0
1	MLY	A	206	1	9,10,11	0.55	0	6,11,13	0.78	0
1	MLY	B	138	1	9,10,11	0.63	0	6,11,13	1.06	1 (16%)
1	MLY	C	72	1	9,10,11	0.61	0	6,11,13	0.78	0
1	MLY	A	181	1	9,10,11	0.51	0	6,11,13	0.82	0
1	MLY	D	145	1	9,10,11	0.60	0	6,11,13	0.87	0
1	MLY	B	187	1	9,10,11	0.73	0	6,11,13	0.62	0
1	MLY	B	105	1	9,10,11	0.53	0	6,11,13	0.74	0
1	MLY	C	124	1	9,10,11	0.59	0	6,11,13	0.73	0
1	MLY	D	124	1	9,10,11	0.55	0	6,11,13	0.83	0
1	MLY	B	196	1	9,10,11	0.63	0	6,11,13	0.80	0
1	MLY	A	8	1	9,10,11	0.54	0	6,11,13	0.60	0
1	MLY	D	206	1	9,10,11	0.66	0	6,11,13	0.66	0
1	MLY	C	181	1	9,10,11	0.54	0	6,11,13	0.77	0
1	MLY	D	63	1	9,10,11	0.47	0	6,11,13	0.97	0
1	MLY	A	119	1	9,10,11	0.65	0	6,11,13	0.74	0
1	MLY	C	145	1	9,10,11	0.63	0	6,11,13	0.87	0
1	MLY	C	206	1	9,10,11	0.57	0	6,11,13	0.77	0
1	MLY	A	72	1	9,10,11	0.68	0	6,11,13	0.66	0
1	MLY	D	181	1	9,10,11	0.51	0	6,11,13	0.70	0
1	MLY	A	138	1	9,10,11	0.49	0	6,11,13	0.76	0
1	MLY	B	119	1	9,10,11	0.63	0	6,11,13	0.72	0
1	MLY	B	54	1	9,10,11	0.56	0	6,11,13	0.76	0
1	MLY	A	124	1	9,10,11	0.55	0	6,11,13	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	B	63	1	-	0/8/9/11	-
1	MLY	A	63	1	-	0/8/9/11	-
1	MLY	A	145	1	-	1/8/9/11	-
1	MLY	D	1	1	-	0/8/9/11	-
1	MLY	C	187	1	-	0/8/9/11	-
1	MLY	C	105	1	-	0/8/9/11	-
1	MLY	D	72	1	-	0/8/9/11	-
1	MLY	A	1	1	-	2/8/9/11	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	C	119	1	-	1/8/9/11	-
1	MLY	C	138	1	-	0/8/9/11	-
1	MLY	B	72	1	-	1/8/9/11	-
1	MLY	D	105	1	-	0/8/9/11	-
1	MLY	C	196	1	-	0/8/9/11	-
1	MLY	A	29	1	-	0/8/9/11	-
1	MLY	D	196	1	-	0/8/9/11	-
1	MLY	B	1	1	-	1/8/9/11	-
1	MLY	B	124	1	-	0/8/9/11	-
1	MLY	C	63	1	-	0/8/9/11	-
1	MLY	D	29	1	-	1/8/9/11	-
1	MLY	D	54	1	-	2/8/9/11	-
1	MLY	D	8	1	-	1/8/9/11	-
1	MLY	D	187	1	-	0/8/9/11	-
1	MLY	B	145	1	-	0/8/9/11	-
1	MLY	C	54	1	-	1/8/9/11	-
1	MLY	B	181	1	-	3/8/9/11	-
1	MLY	A	105	1	-	0/8/9/11	-
1	MLY	B	29	1	-	0/8/9/11	-
1	MLY	C	8	1	-	2/8/9/11	-
1	MLY	D	119	1	-	0/8/9/11	-
1	MLY	A	196	1	-	0/8/9/11	-
1	MLY	C	1	1	-	0/8/9/11	-
1	MLY	D	138	1	-	0/8/9/11	-
1	MLY	C	29	1	-	1/8/9/11	-
1	MLY	A	54	1	-	1/8/9/11	-
1	MLY	B	8	1	-	1/8/9/11	-
1	MLY	A	187	1	-	0/8/9/11	-
1	MLY	A	206	1	-	0/8/9/11	-
1	MLY	B	138	1	-	2/8/9/11	-
1	MLY	C	72	1	-	0/8/9/11	-
1	MLY	A	181	1	-	0/8/9/11	-
1	MLY	D	145	1	-	0/8/9/11	-
1	MLY	B	187	1	-	2/8/9/11	-
1	MLY	B	105	1	-	2/8/9/11	-
1	MLY	C	124	1	-	1/8/9/11	-
1	MLY	D	124	1	-	0/8/9/11	-
1	MLY	B	196	1	-	2/8/9/11	-
1	MLY	A	8	1	-	0/8/9/11	-
1	MLY	D	206	1	-	1/8/9/11	-
1	MLY	C	181	1	-	0/8/9/11	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	D	63	1	-	0/8/9/11	-
1	MLY	A	119	1	-	3/8/9/11	-
1	MLY	C	145	1	-	1/8/9/11	-
1	MLY	C	206	1	-	0/8/9/11	-
1	MLY	A	72	1	-	0/8/9/11	-
1	MLY	D	181	1	-	0/8/9/11	-
1	MLY	A	138	1	-	0/8/9/11	-
1	MLY	B	119	1	-	1/8/9/11	-
1	MLY	B	54	1	-	3/8/9/11	-
1	MLY	A	124	1	-	0/8/9/11	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	138	MLY	CD-CE-NZ	2.28	119.61	113.71

There are no chirality outliers.

5 of 37 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	54	MLY	O-C-CA-CB
1	B	54	MLY	O-C-CA-CB
1	C	119	MLY	C-CA-CB-CG
1	A	119	MLY	CD-CE-NZ-CH2
1	D	54	MLY	CD-CE-NZ-CH2

There are no ring outliers.

11 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	63	MLY	1	0
1	C	105	MLY	2	0
1	A	1	MLY	1	0
1	C	119	MLY	1	0
1	D	105	MLY	1	0
1	D	119	MLY	2	0
1	B	105	MLY	1	0
1	D	206	MLY	1	0
1	C	181	MLY	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	119	MLY	1	0
1	A	72	MLY	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TRS	C	302	-	7,7,7	0.30	0	9,9,9	0.51	0
4	GOL	D	303	-	5,5,5	0.34	0	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	C	302	-	-	0/9/9/9	-
4	GOL	D	303	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	192/207 (92%)	-0.24	1 (0%) 87 90	8, 14, 27, 37	0
1	B	191/207 (92%)	-0.32	2 (1%) 79 83	8, 14, 25, 35	1 (0%)
1	C	192/207 (92%)	-0.24	3 (1%) 70 74	6, 14, 28, 36	2 (1%)
1	D	191/207 (92%)	-0.23	4 (2%) 63 67	8, 15, 26, 29	0
All	All	766/828 (92%)	-0.26	10 (1%) 75 79	6, 14, 27, 37	3 (0%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	45	PHE	4.2
1	A	45	PHE	3.7
1	B	205	GLY	3.6
1	D	205	GLY	3.4
1	D	64	MET	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	MLY	D	206	11/12	0.75	0.22	31,45,53,53	0
1	MLY	B	54	11/12	0.76	0.19	31,39,51,51	0
1	MLY	C	54	11/12	0.81	0.17	28,36,51,51	0
1	MLY	A	54	11/12	0.82	0.18	31,35,51,52	0
1	MLY	C	105	11/12	0.86	0.20	22,28,54,54	0
1	MLY	C	119	11/12	0.88	0.19	17,24,50,53	0
1	MLY	A	1	11/12	0.88	0.15	16,20,44,46	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	MLY	D	1	11/12	0.89	0.14	23,29,45,45	0
1	MLY	B	1	11/12	0.89	0.14	20,30,44,47	0
1	MLY	A	196	11/12	0.90	0.20	13,18,50,51	0
1	MLY	D	54	11/12	0.91	0.14	14,19,44,45	0
1	MLY	A	105	11/12	0.91	0.15	17,28,46,49	0
1	MLY	B	63	11/12	0.92	0.12	17,18,37,39	0
1	MLY	B	72	11/12	0.92	0.12	10,12,35,36	0
1	MLY	B	105	11/12	0.92	0.12	20,28,46,47	0
1	MLY	C	206	11/12	0.92	0.10	11,20,31,35	0
1	MLY	B	119	11/12	0.92	0.15	15,23,47,50	0
1	MLY	B	181	11/12	0.92	0.14	10,13,37,42	0
1	MLY	D	196	11/12	0.92	0.12	13,15,36,41	0
1	MLY	C	8	11/12	0.92	0.12	13,19,39,39	0
1	MLY	A	181	11/12	0.93	0.12	11,15,37,38	0
1	MLY	A	72	11/12	0.93	0.12	10,12,33,33	0
1	MLY	D	8	11/12	0.93	0.13	14,18,39,40	0
1	MLY	C	63	11/12	0.93	0.10	17,20,36,40	0
1	MLY	D	105	11/12	0.93	0.12	20,22,42,45	0
1	MLY	D	119	11/12	0.93	0.15	14,21,47,48	0
1	MLY	A	63	11/12	0.93	0.11	18,23,39,40	0
1	MLY	B	196	11/12	0.93	0.13	13,17,41,45	0
1	MLY	C	1	11/12	0.94	0.10	15,19,29,34	0
1	MLY	D	29	11/12	0.94	0.11	12,15,31,33	0
1	MLY	A	8	11/12	0.94	0.10	15,19,29,33	0
1	MLY	D	72	11/12	0.94	0.11	10,17,32,34	0
1	MLY	C	124	11/12	0.94	0.10	14,17,35,36	0
1	MLY	C	181	11/12	0.94	0.13	9,12,43,45	0
1	MLY	B	187	11/12	0.94	0.15	12,15,44,45	0
1	MLY	A	119	11/12	0.94	0.14	12,19,39,39	0
1	MLY	B	138	11/12	0.95	0.09	10,13,25,35	0
1	MLY	B	8	11/12	0.95	0.11	11,18,35,38	0
1	MLY	D	63	11/12	0.95	0.10	13,17,37,39	0
1	MLY	C	187	11/12	0.95	0.11	11,15,40,43	0
1	MLY	C	196	11/12	0.95	0.12	13,15,45,46	0
1	MLY	A	138	11/12	0.95	0.08	9,11,19,25	0
1	MLY	D	138	11/12	0.95	0.08	10,11,19,25	0
1	MLY	D	187	11/12	0.95	0.12	11,15,39,43	0
1	MLY	A	124	11/12	0.95	0.10	11,15,32,35	0
1	MLY	B	124	11/12	0.95	0.10	11,16,36,39	0
1	MLY	A	187	11/12	0.96	0.11	10,15,39,39	0
1	MLY	C	138	11/12	0.96	0.07	9,11,19,24	0
1	MLY	A	206	11/12	0.96	0.07	11,16,23,23	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	MLY	D	181	11/12	0.96	0.07	8,12,19,22	0
1	MLY	C	72	11/12	0.96	0.09	9,13,28,31	0
1	MLY	B	29	11/12	0.96	0.09	10,12,30,35	0
1	MLY	C	29	11/12	0.96	0.07	9,12,27,35	0
1	MLY	A	145	11/12	0.97	0.06	11,14,19,31	0
1	MLY	D	145	11/12	0.97	0.07	11,16,21,28	0
1	MLY	B	145	11/12	0.97	0.07	11,16,25,26	0
1	MLY	A	29	11/12	0.97	0.08	10,14,31,40	0
1	MLY	C	145	11/12	0.97	0.06	11,14,21,25	0
1	MLY	D	124	11/12	0.97	0.07	11,15,29,30	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	TRS	C	302	8/8	0.80	0.15	27,38,41,43	0
4	GOL	D	303	6/6	0.95	0.10	16,18,20,21	0
2	MN	C	301	1/1	1.00	0.01	8,8,8,8	0
2	MN	D	302	1/1	1.00	0.01	10,10,10,10	0
2	MN	A	301	1/1	1.00	0.01	9,9,9,9	0
2	MN	B	301	1/1	1.00	0.01	9,9,9,9	0

6.5 Other polymers [i](#)

There are no such residues in this entry.