



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 08:56 PM UTC

PDB ID : 8F6B / pdb_00008f6b
Title : Crystal structure of murine PolG2 hexamer bound to DNA
Authors : Wojtaszek, J.L.; Hoff, K.E.; Williams, R.S.
Deposited on : 2022-11-16
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	455	 82% 8% 10%
1	B	455	 83% 7% 10%
1	C	455	 83% 6% 11%
1	D	455	 84% 7% 9%
1	E	455	 82% 6% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	455	 82% 10% 9%
2	G	18	 44% 56%
2	H	18	 78% 22%
2	I	18	 89% 11%
2	J	18	 61% 39%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 20672 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PolG2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	408	Total	C	N	O	S	0	0	0
			3118	1995	538	571	14			
1	B	411	Total	C	N	O	S	0	1	0
			3169	2023	550	581	15			
1	C	404	Total	C	N	O	S	0	1	0
			3120	1991	545	571	13			
1	D	414	Total	C	N	O	S	0	3	0
			3223	2054	565	589	15			
1	E	401	Total	C	N	O	S	0	1	0
			3091	1971	539	567	14			
1	F	416	Total	C	N	O	S	0	2	0
			3248	2071	569	593	15			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	16	MET	-	initiating methionine	UNP Q0VES3
A	460	ALA	-	expression tag	UNP Q0VES3
A	461	ALA	-	expression tag	UNP Q0VES3
A	462	ALA	-	expression tag	UNP Q0VES3
A	463	LEU	-	expression tag	UNP Q0VES3
A	464	GLU	-	expression tag	UNP Q0VES3
A	465	HIS	-	expression tag	UNP Q0VES3
A	466	HIS	-	expression tag	UNP Q0VES3
A	467	HIS	-	expression tag	UNP Q0VES3
A	468	HIS	-	expression tag	UNP Q0VES3
A	469	HIS	-	expression tag	UNP Q0VES3
A	470	HIS	-	expression tag	UNP Q0VES3
B	16	MET	-	initiating methionine	UNP Q0VES3
B	460	ALA	-	expression tag	UNP Q0VES3
B	461	ALA	-	expression tag	UNP Q0VES3
B	462	ALA	-	expression tag	UNP Q0VES3
B	463	LEU	-	expression tag	UNP Q0VES3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	464	GLU	-	expression tag	UNP Q0VES3
B	465	HIS	-	expression tag	UNP Q0VES3
B	466	HIS	-	expression tag	UNP Q0VES3
B	467	HIS	-	expression tag	UNP Q0VES3
B	468	HIS	-	expression tag	UNP Q0VES3
B	469	HIS	-	expression tag	UNP Q0VES3
B	470	HIS	-	expression tag	UNP Q0VES3
C	16	MET	-	initiating methionine	UNP Q0VES3
C	460	ALA	-	expression tag	UNP Q0VES3
C	461	ALA	-	expression tag	UNP Q0VES3
C	462	ALA	-	expression tag	UNP Q0VES3
C	463	LEU	-	expression tag	UNP Q0VES3
C	464	GLU	-	expression tag	UNP Q0VES3
C	465	HIS	-	expression tag	UNP Q0VES3
C	466	HIS	-	expression tag	UNP Q0VES3
C	467	HIS	-	expression tag	UNP Q0VES3
C	468	HIS	-	expression tag	UNP Q0VES3
C	469	HIS	-	expression tag	UNP Q0VES3
C	470	HIS	-	expression tag	UNP Q0VES3
D	16	MET	-	initiating methionine	UNP Q0VES3
D	460	ALA	-	expression tag	UNP Q0VES3
D	461	ALA	-	expression tag	UNP Q0VES3
D	462	ALA	-	expression tag	UNP Q0VES3
D	463	LEU	-	expression tag	UNP Q0VES3
D	464	GLU	-	expression tag	UNP Q0VES3
D	465	HIS	-	expression tag	UNP Q0VES3
D	466	HIS	-	expression tag	UNP Q0VES3
D	467	HIS	-	expression tag	UNP Q0VES3
D	468	HIS	-	expression tag	UNP Q0VES3
D	469	HIS	-	expression tag	UNP Q0VES3
D	470	HIS	-	expression tag	UNP Q0VES3
E	16	MET	-	initiating methionine	UNP Q0VES3
E	460	ALA	-	expression tag	UNP Q0VES3
E	461	ALA	-	expression tag	UNP Q0VES3
E	462	ALA	-	expression tag	UNP Q0VES3
E	463	LEU	-	expression tag	UNP Q0VES3
E	464	GLU	-	expression tag	UNP Q0VES3
E	465	HIS	-	expression tag	UNP Q0VES3
E	466	HIS	-	expression tag	UNP Q0VES3
E	467	HIS	-	expression tag	UNP Q0VES3
E	468	HIS	-	expression tag	UNP Q0VES3
E	469	HIS	-	expression tag	UNP Q0VES3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	470	HIS	-	expression tag	UNP Q0VES3
F	16	MET	-	initiating methionine	UNP Q0VES3
F	460	ALA	-	expression tag	UNP Q0VES3
F	461	ALA	-	expression tag	UNP Q0VES3
F	462	ALA	-	expression tag	UNP Q0VES3
F	463	LEU	-	expression tag	UNP Q0VES3
F	464	GLU	-	expression tag	UNP Q0VES3
F	465	HIS	-	expression tag	UNP Q0VES3
F	466	HIS	-	expression tag	UNP Q0VES3
F	467	HIS	-	expression tag	UNP Q0VES3
F	468	HIS	-	expression tag	UNP Q0VES3
F	469	HIS	-	expression tag	UNP Q0VES3
F	470	HIS	-	expression tag	UNP Q0VES3

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*TP*GP*GP*TP*AP*GP*GP*CP*GP*CP*CP*TP*AP*CP*CP*AP*G)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	18	Total	C	N	O	P	0	0	0
			366	174	69	106	17			
2	H	18	Total	C	N	O	P	0	0	0
			366	174	69	106	17			
2	I	18	Total	C	N	O	P	0	0	0
			366	174	69	106	17			
2	J	18	Total	C	N	O	P	0	0	0
			366	174	69	106	17			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			8	6	2		
5	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	1	Total 1	Na 1	0	0
6	I	1	Total 1	Na 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	15	Total 15	O 15	0	0
7	B	21	Total 21	O 21	0	0
7	C	13	Total 13	O 13	0	0
7	D	6	Total 6	O 6	0	0
7	E	4	Total 4	O 4	0	0
7	F	11	Total 11	O 11	0	0
7	G	1	Total 1	O 1	0	0
7	H	4	Total 4	O 4	0	0
7	I	6	Total 6	O 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PolG2




• Molecule 1: PolG2



• Molecule 1: PolG2




• Molecule 1: PolG2






- Molecule 1: PolG2

Chain E: 82% 6% 12%



- Molecule 1: PolG2

Chain F: 82% 10% 9%



- Molecule 2: DNA (5'-D(*CP*TP*GP*GP*TP*AP*GP*GP*CP*GP*CP*CP*TP*AP*CP*CP*AP*G)-3')

Chain G: 44% 56%



- Molecule 2: DNA (5'-D(*CP*TP*GP*GP*TP*AP*GP*GP*CP*GP*CP*CP*TP*AP*CP*CP*AP*G)-3')

Chain H: 78% 22%




- Molecule 2: DNA (5'-D(*CP*TP*GP*GP*TP*AP*GP*GP*CP*GP*CP*CP*TP*AP*CP*CP*AP*G)-3')

Chain I: 89% 11%



- Molecule 2: DNA (5'-D(*CP*TP*GP*GP*TP*AP*GP*GP*CP*GP*CP*CP*TP*AP*CP*CP*AP*G)-3')

Chain J:  61% 39%



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	116.22Å 116.22Å 250.94Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.04 – 2.75 38.04 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.04-2.75) 99.9 (38.04-2.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.77Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.196 , 0.222 0.200 , 0.228	Depositor DCC
R_{free} test set	1978 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å ²)	82.2	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 53.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.011 for -h,-k,l 0.468 for h,-h-k,-l 0.015 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	20672	wwPDB-VP
Average B, all atoms (Å ²)	96.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, EDO, NA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.11	0/3190	0.24	0/4342
1	B	0.06	0/3244	0.20	0/4409
1	C	0.09	0/3195	0.22	0/4346
1	D	0.06	0/3306	0.20	0/4492
1	E	0.06	0/3166	0.20	0/4308
1	F	0.07	0/3328	0.22	0/4519
2	G	0.19	0/410	0.41	0/631
2	H	0.17	0/410	0.34	0/631
2	I	0.18	0/410	0.33	0/631
2	J	0.19	0/410	0.37	0/631
All	All	0.09	0/21069	0.23	0/28940

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3118	0	3038	18	0
1	B	3169	0	3104	20	0
1	C	3120	0	3058	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	3223	0	3174	17	0
1	E	3091	0	3012	15	0
1	F	3248	0	3218	28	0
2	G	366	0	203	7	0
2	H	366	0	203	4	0
2	I	366	0	203	1	0
2	J	366	0	203	4	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	20	0	30	0	0
4	B	16	0	24	0	0
4	C	12	0	18	0	0
4	D	36	0	54	0	0
4	E	16	0	24	0	0
4	F	28	0	42	0	0
5	A	8	0	14	1	0
5	B	8	0	14	1	0
6	H	1	0	0	0	0
6	I	1	0	0	0	0
7	A	15	0	0	0	0
7	B	21	0	0	0	0
7	C	13	0	0	0	0
7	D	6	0	0	0	0
7	E	4	0	0	0	0
7	F	11	0	0	0	0
7	G	1	0	0	0	0
7	H	4	0	0	0	0
7	I	6	0	0	0	0
All	All	20672	0	19652	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 114 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:239:LYS:HB2	1:A:460:ALA:HB1	1.80	0.62
2:J:5:DT:H2"	2:J:6:DA:C8	2.37	0.59
1:E:175:ASN:ND2	1:F:393:GLU:OE1	2.35	0.59
1:F:337:ARG:NH2	2:H:18:DG:N7	2.47	0.58
1:F:337:ARG:NH1	2:H:18:DG:O6	2.38	0.57

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/455 (88%)	394 (98%)	8 (2%)	0	100	100
1	B	406/455 (89%)	399 (98%)	7 (2%)	0	100	100
1	C	399/455 (88%)	393 (98%)	6 (2%)	0	100	100
1	D	413/455 (91%)	407 (98%)	6 (2%)	0	100	100
1	E	396/455 (87%)	389 (98%)	7 (2%)	0	100	100
1	F	414/455 (91%)	405 (98%)	9 (2%)	0	100	100
All	All	2430/2730 (89%)	2387 (98%)	43 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	331/398 (83%)	327 (99%)	4 (1%)	63	78
1	B	340/398 (85%)	336 (99%)	4 (1%)	63	78
1	C	335/398 (84%)	333 (99%)	2 (1%)	78	88
1	D	349/398 (88%)	347 (99%)	2 (1%)	78	88
1	E	331/398 (83%)	328 (99%)	3 (1%)	70	82
1	F	354/398 (89%)	352 (99%)	2 (1%)	78	88

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	2040/2388 (85%)	2023 (99%)	17 (1%)	73 84

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	343	ARG
1	F	435	THR
1	B	459	VAL
1	C	116	ASP
1	C	205	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	378	ASN
1	F	184	GLN
1	D	147	ASN
1	D	113	GLN
1	E	147	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 38 ligands modelled in this entry, 2 are monoatomic - leaving 36 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	D	509	-	3,3,3	0.43	0	2,2,2	0.38	0
4	EDO	E	501	-	3,3,3	0.44	0	2,2,2	0.38	0
3	GOL	B	506	-	5,5,5	0.89	0	5,5,5	1.37	1 (20%)
4	EDO	A	504	-	3,3,3	0.43	0	2,2,2	0.39	0
4	EDO	D	502	-	3,3,3	0.42	0	2,2,2	0.38	0
4	EDO	B	502	-	3,3,3	0.43	0	2,2,2	0.39	0
5	MPD	B	505	-	7,7,7	0.32	0	9,10,10	0.36	0
3	GOL	A	501	-	5,5,5	0.93	0	5,5,5	1.08	0
4	EDO	D	506	-	3,3,3	0.43	0	2,2,2	0.36	0
5	MPD	A	507	-	7,7,7	0.31	0	9,10,10	0.37	0
4	EDO	D	504	-	3,3,3	0.43	0	2,2,2	0.38	0
4	EDO	A	506	-	3,3,3	0.44	0	2,2,2	0.35	0
4	EDO	F	504	-	3,3,3	0.42	0	2,2,2	0.39	0
4	EDO	F	506	-	3,3,3	0.43	0	2,2,2	0.38	0
4	EDO	A	503	-	3,3,3	0.43	0	2,2,2	0.39	0
4	EDO	D	505	-	3,3,3	0.42	0	2,2,2	0.38	0
4	EDO	F	501	-	3,3,3	0.42	0	2,2,2	0.41	0
4	EDO	C	503	-	3,3,3	0.42	0	2,2,2	0.38	0
4	EDO	B	504	-	3,3,3	0.42	0	2,2,2	0.38	0
4	EDO	B	501	-	3,3,3	0.43	0	2,2,2	0.38	0
4	EDO	C	501	-	3,3,3	0.42	0	2,2,2	0.37	0
4	EDO	B	503	-	3,3,3	0.42	0	2,2,2	0.39	0
4	EDO	A	505	-	3,3,3	0.43	0	2,2,2	0.39	0
4	EDO	D	503	-	3,3,3	0.43	0	2,2,2	0.38	0
4	EDO	A	502	-	3,3,3	0.43	0	2,2,2	0.38	0
4	EDO	F	507	-	3,3,3	0.43	0	2,2,2	0.39	0
4	EDO	F	505	-	3,3,3	0.43	0	2,2,2	0.37	0
4	EDO	D	501	-	3,3,3	0.43	0	2,2,2	0.38	0
4	EDO	D	508	-	3,3,3	0.42	0	2,2,2	0.39	0
4	EDO	E	502	-	3,3,3	0.42	0	2,2,2	0.38	0
4	EDO	C	502	-	3,3,3	0.43	0	2,2,2	0.38	0
4	EDO	E	504	-	3,3,3	0.43	0	2,2,2	0.37	0
4	EDO	F	503	-	3,3,3	0.43	0	2,2,2	0.38	0
4	EDO	D	507	-	3,3,3	0.43	0	2,2,2	0.38	0
4	EDO	F	502	-	3,3,3	0.43	0	2,2,2	0.39	0
4	EDO	E	503	-	3,3,3	0.44	0	2,2,2	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	D	509	-	-	0/1/1/1	-
4	EDO	E	501	-	-	0/1/1/1	-
3	GOL	B	506	-	-	0/4/4/4	-
4	EDO	A	504	-	-	0/1/1/1	-
4	EDO	D	502	-	-	0/1/1/1	-
4	EDO	B	502	-	-	0/1/1/1	-
5	MPD	B	505	-	-	1/5/5/5	-
3	GOL	A	501	-	-	0/4/4/4	-
4	EDO	D	506	-	-	0/1/1/1	-
5	MPD	A	507	-	-	0/5/5/5	-
4	EDO	D	504	-	-	0/1/1/1	-
4	EDO	A	506	-	-	0/1/1/1	-
4	EDO	F	504	-	-	0/1/1/1	-
4	EDO	F	506	-	-	0/1/1/1	-
4	EDO	A	503	-	-	0/1/1/1	-
4	EDO	D	505	-	-	0/1/1/1	-
4	EDO	F	501	-	-	1/1/1/1	-
4	EDO	C	503	-	-	0/1/1/1	-
4	EDO	B	504	-	-	0/1/1/1	-
4	EDO	B	501	-	-	0/1/1/1	-
4	EDO	C	501	-	-	0/1/1/1	-
4	EDO	B	503	-	-	0/1/1/1	-
4	EDO	A	505	-	-	0/1/1/1	-
4	EDO	D	503	-	-	0/1/1/1	-
4	EDO	A	502	-	-	0/1/1/1	-
4	EDO	F	507	-	-	0/1/1/1	-
4	EDO	F	505	-	-	1/1/1/1	-
4	EDO	D	501	-	-	0/1/1/1	-
4	EDO	D	508	-	-	0/1/1/1	-
4	EDO	E	502	-	-	0/1/1/1	-
4	EDO	C	502	-	-	0/1/1/1	-
4	EDO	E	504	-	-	0/1/1/1	-
4	EDO	F	503	-	-	0/1/1/1	-
4	EDO	D	507	-	-	0/1/1/1	-
4	EDO	F	502	-	-	0/1/1/1	-
4	EDO	E	503	-	-	0/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	506	GOL	C3-C2-C1	-2.33	103.26	111.80

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	F	501	EDO	O1-C1-C2-O2
4	F	505	EDO	O1-C1-C2-O2
5	B	505	MPD	CM-C2-C3-C4

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	505	MPD	1	0
5	A	507	MPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å²)	Q<0.9	
1	A	408/455 (89%)	-1.00	0	100	100	54, 86, 169, 235	0
1	B	411/455 (90%)	-0.92	0	100	100	56, 86, 170, 233	1 (0%)
1	C	404/455 (88%)	-0.83	0	100	100	56, 95, 173, 216	1 (0%)
1	D	414/455 (90%)	-0.97	0	100	100	53, 87, 168, 223	3 (0%)
1	E	401/455 (88%)	-0.83	0	100	100	57, 94, 169, 203	1 (0%)
1	F	416/455 (91%)	-0.96	2 (0%)	87	87	54, 87, 168, 233	2 (0%)
2	G	18/18 (100%)	-1.01	0	100	100	85, 103, 124, 127	0
2	H	18/18 (100%)	-0.74	0	100	100	93, 103, 126, 126	0
2	I	18/18 (100%)	-0.93	0	100	100	90, 100, 116, 120	0
2	J	18/18 (100%)	-1.11	0	100	100	85, 101, 113, 117	0
All	All	2526/2802 (90%)	-0.92	2 (0%)	92	93	53, 90, 170, 235	8 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	130	ILE	2.3
1	F	462	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	D	504	4/4	0.89	0.15	91,94,101,118	0
4	EDO	F	501	4/4	0.92	0.13	123,123,124,146	0
4	EDO	A	502	4/4	0.94	0.11	82,98,99,113	0
4	EDO	D	503	4/4	0.94	0.16	120,121,124,132	0
5	MPD	A	507	8/8	0.94	0.11	80,122,130,138	0
4	EDO	B	502	4/4	0.95	0.14	94,95,104,107	0
4	EDO	D	502	4/4	0.95	0.10	101,101,103,116	0
4	EDO	A	503	4/4	0.95	0.18	94,103,103,109	0
4	EDO	D	506	4/4	0.96	0.13	69,87,103,105	0
4	EDO	B	503	4/4	0.96	0.07	82,84,84,107	0
4	EDO	A	505	4/4	0.96	0.10	81,88,102,108	0
5	MPD	B	505	8/8	0.96	0.08	79,114,123,128	0
4	EDO	D	505	4/4	0.97	0.18	91,96,105,126	0
4	EDO	C	501	4/4	0.97	0.16	85,90,102,110	0
4	EDO	E	503	4/4	0.97	0.16	80,87,89,101	0
4	EDO	A	506	4/4	0.97	0.15	69,81,96,116	0
4	EDO	F	503	4/4	0.97	0.17	73,84,86,100	0
4	EDO	F	506	4/4	0.97	0.19	81,92,102,105	0
3	GOL	B	506	6/6	0.97	0.10	121,124,167,190	0
3	GOL	A	501	6/6	0.97	0.07	80,102,106,112	0
4	EDO	E	502	4/4	0.98	0.12	86,92,102,103	0
4	EDO	B	504	4/4	0.98	0.06	75,77,86,91	0
4	EDO	E	504	4/4	0.98	0.16	77,93,99,106	0
4	EDO	A	504	4/4	0.98	0.08	75,84,85,90	0
4	EDO	F	502	4/4	0.98	0.10	78,82,86,86	0
4	EDO	D	501	4/4	0.98	0.08	77,80,83,83	0
4	EDO	F	504	4/4	0.98	0.13	79,83,85,90	0
4	EDO	F	505	4/4	0.98	0.10	73,82,105,118	0
4	EDO	B	501	4/4	0.98	0.11	75,79,94,100	0
4	EDO	F	507	4/4	0.98	0.15	72,79,80,91	0
4	EDO	D	508	4/4	0.98	0.12	80,82,87,92	0
4	EDO	D	509	4/4	0.98	0.15	66,82,82,85	0
6	NA	H	101	1/1	0.98	0.04	178,178,178,178	0
4	EDO	C	502	4/4	0.99	0.10	80,88,97,97	0
4	EDO	C	503	4/4	0.99	0.07	63,83,89,102	0
4	EDO	E	501	4/4	0.99	0.13	71,86,90,117	0
4	EDO	D	507	4/4	0.99	0.08	68,75,82,99	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	NA	I	101	1/1	0.99	0.04	215,215,215,215	0

6.5 Other polymers [i](#)

There are no such residues in this entry.