



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:48 AM UTC

PDB ID : 1F75 / pdb_00001f75
Title : CRYSTAL STRUCTURE OF UNDECAPRENYL DIPHOSPHATE SYNTHASE FROM MICROCOCCUS LUTEUS B-P 26
Authors : Fujihashi, M.; Zhang, Y.-W.; Higuchi, Y.; Li, X.-Y.; Koyama, T.; Miki, K.
Deposited on : 2000-06-26
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	249	
1	B	249	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3552 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called UNDECAPRENYL PYROPHOSPHATE SYNTHETASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	0	0
			1769	1140	294	328	7			
1	B	212	Total	C	N	O	S	0	0	0
			1735	1120	287	321	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	54	ARG	LYS	engineered mutation	UNP O82827
B	54	ARG	LYS	engineered mutation	UNP O82827

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	O S	0	0
			5	4 1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	19	Total	O	0	0
			19	19		
3	B	19	Total	O	0	0
			19	19		

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	127.20Å 60.16Å 75.70Å 90.00° 105.69° 90.00°	Depositor
Resolution (Å)	50.00 – 2.20	Depositor
% Data completeness (in resolution range)	(Not available) (50.00-2.20)	Depositor
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.843	Depositor
R, R_{free}	0.190 , 0.246	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3552	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/1808	0.84	1/2440 (0.0%)
1	B	0.38	0/1774	0.84	0/2393
All	All	0.39	0/3582	0.84	1/4833 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	221	ASP	N-CA-C	6.77	118.74	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1769	0	1761	30	0
1	B	1735	0	1728	19	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	19	0	0	0	0
3	B	19	0	0	1	0
All	All	3552	0	3489	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:LYS:HB3	1:A:40:MET:HE3	1.59	0.83
1:A:107:ASN:HD21	1:A:138:THR:H	1.31	0.78
1:A:204:LEU:HB2	1:B:217:PHE:CE1	2.28	0.69
1:A:107:ASN:ND2	1:A:138:THR:H	1.94	0.66
1:A:38:LYS:CB	1:A:40:MET:HE3	2.30	0.62
1:A:154:ILE:O	1:A:158:VAL:HG23	2.00	0.61
1:A:15:ILE:HD13	1:A:19:GLN:HG3	1.83	0.59
1:A:17:ALA:C	1:A:19:GLN:H	2.10	0.58
1:A:219:PHE:HE1	1:B:219:PHE:HE1	1.51	0.57
1:B:181:ASN:HB3	3:B:627:HOH:O	2.06	0.56
1:A:23:HIS:HD2	1:A:190:PRO:O	1.90	0.54
1:A:239:TYR:C	1:A:241:ASN:H	2.15	0.54
1:B:216:GLU:HG3	1:B:242:ARG:HH21	1.72	0.54
1:A:15:ILE:HD11	1:A:17:ALA:HB3	1.90	0.54
1:B:157:ALA:O	1:B:161:ILE:HG13	2.08	0.53
1:A:36:LYS:HE3	1:A:37:GLN:HE21	1.73	0.53
1:B:197:ARG:HB3	1:B:219:PHE:HD2	1.76	0.50
1:A:102:GLU:O	1:A:106:LYS:HB2	2.13	0.49
1:B:91:LEU:HB3	1:B:92:PRO:HD3	1.94	0.49
1:B:39:LYS:HB2	1:B:39:LYS:NZ	2.27	0.49
1:A:241:ASN:O	1:A:242:ARG:HB2	2.14	0.48
1:B:86:ASN:O	1:B:90:LYS:HE3	2.14	0.47
1:B:224:TRP:HB3	1:B:225:PRO:HD3	1.96	0.46
1:A:204:LEU:HB2	1:B:217:PHE:CZ	2.51	0.45
1:A:49:GLY:O	1:A:53:VAL:HG23	2.16	0.45
1:B:234:GLN:O	1:B:238:ILE:HG13	2.16	0.45
1:A:160:LEU:O	1:A:164:ARG:HG3	2.17	0.44
1:B:216:GLU:CG	1:B:242:ARG:HH21	2.30	0.44
1:A:166:LYS:HD2	1:B:165:TYR:CE2	2.53	0.44
1:A:34:TRP:CH2	1:A:48:GLU:HG3	2.53	0.44
1:A:52:THR:HA	1:A:55:LYS:HE2	2.00	0.44
1:A:17:ALA:O	1:A:19:GLN:N	2.49	0.43
1:A:204:LEU:HB2	1:B:217:PHE:HE1	1.77	0.43
1:A:15:ILE:HG21	1:A:19:GLN:HG3	1.99	0.43
1:A:204:LEU:HD23	1:A:205:SER:N	2.34	0.42
1:B:38:LYS:HB3	1:B:40:MET:HE3	2.02	0.42
1:A:189:MET:HA	1:A:190:PRO:HD3	1.84	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:165:TYR:CE2	1:B:166:LYS:HD2	2.56	0.41
1:B:122:HIS:CD2	1:B:122:HIS:H	2.37	0.41
1:A:197:ARG:HD3	1:A:201:GLU:O	2.20	0.41
1:B:23:HIS:CE1	1:B:192:PRO:HG3	2.55	0.41
1:A:201:GLU:HG2	1:A:203:ARG:HH21	1.86	0.41
1:A:224:TRP:HB3	1:A:225:PRO:HD3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/249 (86%)	207 (97%)	3 (1%)	3 (1%)	9	7
1	B	208/249 (84%)	200 (96%)	7 (3%)	1 (0%)	24	27
All	All	421/498 (84%)	407 (97%)	10 (2%)	4 (1%)	12	11

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	17	ALA
1	B	201	GLU
1	A	18	ALA
1	A	240	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/221 (87%)	190 (99%)	2 (1%)	68	81
1	B	189/221 (86%)	183 (97%)	6 (3%)	34	47
All	All	381/442 (86%)	373 (98%)	8 (2%)	47	63

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	106	LYS
1	A	221	ASP
1	B	54	ARG
1	B	87	TYR
1	B	106	LYS
1	B	140	LEU
1	B	160	LEU
1	B	208	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	HIS
1	A	37	GLN
1	A	107	ASN
1	A	147	ASN
1	B	122	HIS
1	B	147	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	901	-	4,4,4	1.76	1 (25%)	6,6,6	0.84	0
2	SO4	B	902	-	4,4,4	1.90	1 (25%)	6,6,6	0.81	0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	902	SO4	O1-S	3.33	1.64	1.44
2	A	901	SO4	O1-S	2.98	1.62	1.44

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.