



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:12 PM UTC

PDB ID : 4F7G / pdb_00004f7g
Title : Crystal structure of talin autoinhibition complex
Authors : Song, X.; Qin, J.; Ye, S.; Zhang, R.
Deposited on : 2012-05-16
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	222	4% 65% 18% • 14%
2	B	216	% 61% 15% • 21%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Talin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	192	1554	1000	265	282	7	0	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	184	MET	-	expression tag	UNP P26039
A	185	HIS	-	expression tag	UNP P26039
A	186	HIS	-	expression tag	UNP P26039
A	187	HIS	-	expression tag	UNP P26039
A	188	HIS	-	expression tag	UNP P26039
A	189	HIS	-	expression tag	UNP P26039
A	190	HIS	-	expression tag	UNP P26039
A	191	SER	-	expression tag	UNP P26039
A	192	SER	-	expression tag	UNP P26039
A	193	GLY	-	expression tag	UNP P26039
A	194	VAL	-	expression tag	UNP P26039
A	195	ASP	-	expression tag	UNP P26039
A	196	LEU	-	expression tag	UNP P26039
A	197	GLY	-	expression tag	UNP P26039
A	198	THR	-	expression tag	UNP P26039
A	199	GLU	-	expression tag	UNP P26039
A	200	ASN	-	expression tag	UNP P26039
A	201	LEU	-	expression tag	UNP P26039
A	202	TYR	-	expression tag	UNP P26039
A	203	PHE	-	expression tag	UNP P26039
A	204	GLN	-	expression tag	UNP P26039
A	205	SER	-	expression tag	UNP P26039

- Molecule 2 is a protein called Talin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	170	Total	C	N	O	S	0	0	0
			1243	765	214	257	7			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1632	MET	-	expression tag	UNP P26039
B	1633	HIS	-	expression tag	UNP P26039
B	1634	HIS	-	expression tag	UNP P26039
B	1635	HIS	-	expression tag	UNP P26039
B	1636	HIS	-	expression tag	UNP P26039
B	1637	HIS	-	expression tag	UNP P26039
B	1638	HIS	-	expression tag	UNP P26039
B	1639	SER	-	expression tag	UNP P26039
B	1640	SER	-	expression tag	UNP P26039
B	1641	GLY	-	expression tag	UNP P26039
B	1642	VAL	-	expression tag	UNP P26039
B	1643	ASP	-	expression tag	UNP P26039
B	1644	LEU	-	expression tag	UNP P26039
B	1645	GLY	-	expression tag	UNP P26039
B	1646	THR	-	expression tag	UNP P26039
B	1647	GLU	-	expression tag	UNP P26039
B	1648	ASN	-	expression tag	UNP P26039
B	1649	LEU	-	expression tag	UNP P26039
B	1650	TYR	-	expression tag	UNP P26039
B	1651	PHE	-	expression tag	UNP P26039
B	1652	GLN	-	expression tag	UNP P26039
B	1653	SER	-	expression tag	UNP P26039


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	60	Total	O	0	0
			60	60		
3	B	49	Total	O	0	0
			49	49		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Talin-1



• Molecule 2: Talin-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	36.66Å 86.88Å 113.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	69.02 – 2.05 69.02 – 2.05	Depositor EDS
% Data completeness (in resolution range)	87.8 (69.02-2.05) 87.7 (69.02-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.210 , 0.249 0.208 , 0.234	Depositor DCC
R_{free} test set	1207 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	48.8	Xtriage
Anisotropy	0.390	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 67.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2906	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/1586	1.02	5/2129 (0.2%)
2	B	0.88	0/1256	1.07	1/1708 (0.1%)
All	All	0.77	0/2842	1.04	6/3837 (0.2%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	307	THR	N-CA-C	5.89	119.94	112.87
2	B	1775	LEU	N-CA-C	-5.77	104.89	111.07
1	A	248	GLY	CA-C-N	5.23	125.24	119.90
1	A	248	GLY	C-N-CA	5.23	125.24	119.90
1	A	236	CYS	N-CA-C	5.04	116.47	111.07
1	A	373	TYR	N-CA-C	-5.04	107.08	113.43

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1554	0	1575	22	0
2	B	1243	0	1239	17	0
3	A	60	0	0	4	0
3	B	49	0	0	3	0
All	All	2906	0	2814	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:310:VAL:HG22	1:A:333:THR:CG2	2.16	0.75
2:B:1712:LEU:HD11	2:B:1733:GLN:HB3	1.69	0.74
1:A:399:ILE:O	1:A:400:LEU:HD13	1.98	0.63
1:A:268:LYS:HD2	3:A:524:HOH:O	1.98	0.63
1:A:399:ILE:O	1:A:399:ILE:HG22	1.98	0.62
2:B:1731:VAL:HG23	2:B:1732:SER:N	2.16	0.60
1:A:223:ILE:HD11	3:A:505:HOH:O	2.05	0.56
1:A:384:GLU:O	1:A:387:GLN:HG2	2.06	0.54
2:B:1714:GLU:HG3	3:B:1927:HOH:O	2.10	0.52
2:B:1771:SER:OG	2:B:1803:MET:HG2	2.10	0.51
1:A:344:THR:OG1	1:A:346:GLU:HG3	2.12	0.50
1:A:334:LYS:O	1:A:353:LEU:HD22	2.13	0.49
2:B:1696:GLU:OE1	2:B:1824:VAL:N	2.42	0.48
2:B:1736:GLN:O	2:B:1740:PRO:HD2	2.13	0.48
2:B:1724:ALA:HB2	2:B:1784:GLY:HA3	1.96	0.48
1:A:289:MET:HG3	1:A:293:GLU:HG2	1.94	0.48
1:A:310:VAL:HG22	1:A:333:THR:HG21	1.96	0.48
1:A:399:ILE:C	1:A:400:LEU:HD22	2.39	0.48
1:A:219:ALA:O	1:A:223:ILE:HG23	2.14	0.47
2:B:1731:VAL:CG2	2:B:1732:SER:N	2.78	0.46
2:B:1676:ASP:OD1	2:B:1766:LYS:NZ	2.46	0.46
1:A:266:LEU:HD11	1:A:279:ILE:HD11	1.98	0.46
1:A:275:GLY:HA2	1:A:278:LYS:HE2	1.97	0.45
1:A:357:LYS:HE2	1:A:371:GLY:HA3	1.99	0.44
1:A:326:VAL:HG22	1:A:327:PRO:O	2.17	0.43
1:A:335:GLU:HG2	1:A:336:CYS:N	2.33	0.43
2:B:1802:MET:HG2	3:B:1915:HOH:O	2.19	0.43
2:B:1702:MET:HE1	2:B:1761:LEU:HD12	2.00	0.43
2:B:1706:VAL:HG21	2:B:1810:LEU:HD13	2.01	0.43
2:B:1666:ALA:O	3:B:1949:HOH:O	2.21	0.43
2:B:1668:LEU:HD12	2:B:1776:LEU:CD2	2.49	0.42
2:B:1774:GLN:HE21	2:B:1778:THR:HG23	1.84	0.42
1:A:297:ARG:NH2	3:A:521:HOH:O	2.53	0.41
2:B:1736:GLN:HG2	2:B:1737:TYR:N	2.35	0.41
1:A:398:ILE:C	1:A:400:LEU:H	2.28	0.41
2:B:1658:GLN:HG3	2:B:1780:LYS:HE2	2.03	0.41
1:A:256:LYS:HG3	3:A:539:HOH:O	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:GLU:O	1:A:241:PHE:HD2	2.05	0.40
1:A:348:ILE:HD13	1:A:348:ILE:HG21	1.77	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	190/222 (86%)	181 (95%)	7 (4%)	2 (1%)	11	5
2	B	168/216 (78%)	165 (98%)	3 (2%)	0	100	100
All	All	358/438 (82%)	346 (97%)	10 (3%)	2 (1%)	21	13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	334	LYS
1	A	399	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/196 (86%)	147 (88%)	21 (12%)	4	1
2	B	129/168 (77%)	112 (87%)	17 (13%)	4	1

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	297/364 (82%)	259 (87%)	38 (13%)	4 1

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	212	LEU
1	A	214	LEU
1	A	215	LEU
1	A	223	ILE
1	A	254	LYS
1	A	256	LYS
1	A	263	LYS
1	A	268	LYS
1	A	272	LYS
1	A	278	LYS
1	A	292	ILE
1	A	293	GLU
1	A	296	VAL
1	A	303	ARG
1	A	306	LYS
1	A	334	LYS
1	A	335	GLU
1	A	342	GLU
1	A	343	LYS
1	A	387	GLN
1	A	391	LEU
2	B	1703	LEU
2	B	1723	GLU
2	B	1725	SER
2	B	1732	SER
2	B	1743	LEU
2	B	1753	LEU
2	B	1761	LEU
2	B	1762	LEU
2	B	1775	LEU
2	B	1776	LEU
2	B	1781	GLU
2	B	1788	GLN
2	B	1794	GLU
2	B	1802	MET
2	B	1810	LEU
2	B	1814	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	1824	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	242	GLN
1	A	246	GLN
1	A	283	HIS
1	A	355	ASN
1	A	390	GLN
2	B	1685	GLN
2	B	1701	GLN
2	B	1755	HIS
2	B	1774	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	192/222 (86%)	0.09	8 (4%)	40 40	35, 65, 108, 133	0
2	B	170/216 (78%)	-0.10	2 (1%)	76 78	30, 53, 97, 107	0
All	All	362/438 (82%)	-0.00	10 (2%)	55 57	30, 60, 101, 133	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1655	ALA	3.5
1	A	209	PRO	3.1
1	A	210	VAL	3.1
1	A	271	VAL	2.5
1	A	286	CYS	2.3
1	A	372	ASP	2.3
1	A	400	LEU	2.1
1	A	262	LEU	2.1
2	B	1737	TYR	2.1
1	A	296	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.