



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:32 AM UTC

PDB ID : 4F7S / pdb_00004f7s
Title : Crystal structure of human CDK8/CYCC in the DMG-in conformation
Authors : Schneider, E.V.; Boettcher, J.; Huber, R.; Maskos, K.
Deposited on : 2012-05-16
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	405	10% 77% 6% 16%
2	B	287	4% 87% 6% 7%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	339	2811	1812	491	490	18	129	2	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ASP	-	expression tag	UNP P49336
A	0	LYS	-	expression tag	UNP P49336


- Molecule 2 is a protein called Cyclin-C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	267	2225	1460	363	385	17	61	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	ASP	-	expression tag	UNP P24863
B	-2	ASP	-	expression tag	UNP P24863
B	-1	LYS	-	expression tag	UNP P24863
B	0	ALA	-	expression tag	UNP P24863

- Molecule 3 is N-(2-phenylethyl)quinazolin-4-amine (CCD ID: 0SW) (formula: C₁₆H₁₅N₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			19	16	3		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is FORMIC ACID (CCD ID: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			3	1	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	128	Total 128	O 128	0	0
6	B	81	Total 81	O 81	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cyclin-dependent kinase 8



- Molecule 2: Cyclin-C



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.50Å 70.77Å 170.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	85.07 – 2.20 85.07 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.5 (85.07-2.20) 99.5 (85.07-2.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.193 , 0.222 0.205 , 0.235	Depositor DCC
R_{free} test set	1468 reflections (3.33%)	wwPDB-VP
Wilson B-factor (Å ²)	38.4	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.035 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5319	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, OSW, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	1/2889 (0.0%)	0.83	3/3891 (0.1%)
2	B	0.69	0/2284	0.85	0/3094
All	All	0.67	1/5173 (0.0%)	0.84	3/6985 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	43	GLY	C-N	-8.14	1.22	1.33

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	43	GLY	O-C-N	-8.63	111.47	122.70
1	A	122	GLN	N-CA-CB	-7.28	98.12	110.50
1	A	123	LEU	N-CA-CB	5.19	115.32	110.39

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2811	0	2809	14	0
2	B	2225	0	2261	18	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	19	0	15	0	0
4	A	24	0	36	0	0
4	B	28	0	42	0	0
5	B	3	0	1	0	0
6	A	128	0	0	0	0
6	B	81	0	0	0	0
All	All	5319	0	5164	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:ILE:HD13	2:B:143:LEU:HD22	1.76	0.67
2:B:72:PHE:CD1	2:B:145:LEU:HD23	2.31	0.64
2:B:76:TYR:CD2	2:B:145:LEU:HD22	2.37	0.59
2:B:76:TYR:CE2	2:B:145:LEU:HD13	2.40	0.57
2:B:40:LEU:HD23	2:B:250:MET:HE1	1.88	0.54
2:B:72:PHE:HA	2:B:145:LEU:HD23	1.92	0.51
2:B:163:VAL:HG21	2:B:173:LEU:HD13	1.93	0.51
1:A:62:SER:HA	1:A:177:ALA:HB2	1.93	0.51
1:A:108:ILE:HD13	1:A:226:GLU:HG2	1.93	0.51
1:A:146:TRP:CZ3	2:B:2:ALA:HB3	2.46	0.50
2:B:72:PHE:HA	2:B:145:LEU:CD2	2.42	0.49
2:B:172:LEU:HD23	2:B:209:GLN:OE1	2.14	0.48
1:A:315:LEU:HD23	1:A:324:ILE:CG2	2.43	0.48
2:B:76:TYR:CE2	2:B:145:LEU:HD22	2.49	0.47
2:B:40:LEU:HD21	2:B:254:LEU:HD23	1.97	0.47
2:B:76:TYR:HD2	2:B:145:LEU:HD22	1.80	0.46
2:B:92:PHE:CD1	2:B:108:LEU:HD22	2.51	0.46
2:B:76:TYR:HE2	2:B:145:LEU:HD13	1.79	0.45
1:A:279:LEU:HD12	1:A:283:PHE:CD2	2.52	0.45
1:A:79:ILE:HD13	1:A:158:LEU:HD12	1.99	0.45
1:A:152:LEU:HD22	1:A:157:ILE:HD11	1.99	0.44
1:A:59:ILE:HG22	1:A:64[A]:CYS:SG	2.59	0.43
1:A:132:LEU:HD22	1:A:224:PHE:CD1	2.53	0.43
1:A:59:ILE:HD13	2:B:143:LEU:CD2	2.48	0.43
1:A:297:MET:HE3	1:A:302:VAL:HG21	2.00	0.43
2:B:217:PHE:HD1	2:B:224:MET:HE1	1.84	0.42
1:A:233:ILE:HG23	1:A:234:PHE:CD2	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:40:LEU:HD23	2:B:250:MET:CE	2.52	0.40
1:A:139:ILE:HG13	1:A:220:ILE:HD11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/405 (82%)	325 (98%)	8 (2%)	0	100	100
2	B	265/287 (92%)	263 (99%)	2 (1%)	0	100	100
All	All	598/692 (86%)	588 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/360 (84%)	303 (100%)	1 (0%)	86	93
2	B	244/261 (94%)	243 (100%)	1 (0%)	84	92
All	All	548/621 (88%)	546 (100%)	2 (0%)	84	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	ILE
2	B	160	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	GLN
1	A	88	HIS
1	A	210	HIS
1	A	310	HIS
1	A	331	GLN
2	B	26	GLN
2	B	203	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	B	307	-	3,3,3	0.49	0	2,2,2	0.31	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	507	-	3,3,3	0.47	0	2,2,2	0.09	0
4	EDO	B	304	-	3,3,3	0.51	0	2,2,2	0.18	0
4	EDO	A	503	-	3,3,3	0.28	0	2,2,2	0.43	0
4	EDO	B	305	-	3,3,3	0.42	0	2,2,2	0.27	0
3	0SW	A	501	-	21,21,21	2.10	5 (23%)	25,27,27	2.38	5 (20%)
4	EDO	A	505	-	3,3,3	0.47	0	2,2,2	0.25	0
4	EDO	A	502	-	3,3,3	0.53	0	2,2,2	0.08	0
4	EDO	B	303	-	3,3,3	0.43	0	2,2,2	0.31	0
4	EDO	B	301	-	3,3,3	0.44	0	2,2,2	0.38	0
4	EDO	A	506	-	3,3,3	0.51	0	2,2,2	0.32	0
4	EDO	B	302	-	3,3,3	0.46	0	2,2,2	0.24	0
5	FMT	B	308	-	2,2,2	0.73	0	1,1,1	0.49	0
4	EDO	A	504	-	3,3,3	0.51	0	2,2,2	0.25	0
4	EDO	B	306	-	3,3,3	0.47	0	2,2,2	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	307	-	-	1/1/1/1	-
4	EDO	A	507	-	-	1/1/1/1	-
4	EDO	B	304	-	-	1/1/1/1	-
4	EDO	A	503	-	-	1/1/1/1	-
4	EDO	B	305	-	-	0/1/1/1	-
3	0SW	A	501	-	-	2/6/6/6	0/3/3/3
4	EDO	A	505	-	-	1/1/1/1	-
4	EDO	A	502	-	-	1/1/1/1	-
4	EDO	B	303	-	-	0/1/1/1	-
4	EDO	B	301	-	-	0/1/1/1	-
4	EDO	A	506	-	-	1/1/1/1	-
4	EDO	B	302	-	-	1/1/1/1	-
4	EDO	A	504	-	-	1/1/1/1	-
4	EDO	B	306	-	-	0/1/1/1	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	0SW	C10-N11	5.12	1.42	1.34
3	A	501	0SW	C10-N8	4.17	1.39	1.34
3	A	501	0SW	C7-N9	4.00	1.38	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	0SW	C10-C2	-3.46	1.40	1.44
3	A	501	0SW	C7-N8	2.73	1.38	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	0SW	C7-N8-C10	7.39	122.34	116.60
3	A	501	0SW	N9-C7-N8	-6.62	119.69	128.67
3	A	501	0SW	C10-C2-C1	3.89	118.16	115.86
3	A	501	0SW	C2-C10-N8	-3.58	118.70	121.35
3	A	501	0SW	C7-N9-C1	3.03	118.87	115.43

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	304	EDO	O1-C1-C2-O2
4	A	505	EDO	O1-C1-C2-O2
4	B	307	EDO	O1-C1-C2-O2
3	A	501	0SW	C12-C13-C14-C17
3	A	501	0SW	C12-C13-C14-C19
4	A	504	EDO	O1-C1-C2-O2
4	A	507	EDO	O1-C1-C2-O2
4	A	502	EDO	O1-C1-C2-O2
4	A	506	EDO	O1-C1-C2-O2
4	B	302	EDO	O1-C1-C2-O2
4	A	503	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	338/405 (83%)	0.52	40 (11%) 9 7	23, 45, 75, 94	35 (10%)
2	B	267/287 (93%)	0.26	11 (4%) 41 38	24, 44, 66, 94	19 (7%)
All	All	605/692 (87%)	0.40	51 (8%) 17 14	23, 44, 70, 94	54 (8%)

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	43	GLY	6.4
1	A	359	LEU	4.7
1	A	358	PHE	4.6
1	A	33	GLY	4.3
1	A	186	PRO	4.3
1	A	243	THR	4.2
1	A	32	TYR	4.1
2	B	260	PRO	3.8
1	A	350	GLN	3.8
1	A	357	GLU	3.7
1	A	122	GLN	3.7
1	A	244	SER	3.5
1	A	42	ASP	3.5
2	B	264	PRO	3.5
1	A	248[A]	HIS	3.4
1	A	31	THR	3.3
1	A	45	ASP	3.3
1	A	196	THR	3.1
2	B	75	ARG	3.1
2	B	145	LEU	2.8
1	A	115	LYS	2.8
1	A	28	GLY	2.8
2	B	221	SER	2.8
1	A	55	GLU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	114	SER	2.7
1	A	124	PRO	2.7
1	A	30	GLY	2.6
1	A	123	LEU	2.6
1	A	59	ILE	2.6
1	A	343	SER	2.6
2	B	-2	ASP	2.6
1	A	1	MET	2.5
1	A	113	ALA	2.5
1	A	34	HIS	2.4
2	B	171	MET	2.4
2	B	98	GLU	2.4
1	A	46	ASP	2.3
1	A	179	LEU	2.3
1	A	47	LYS	2.3
1	A	40	ARG	2.3
1	A	355	LYS	2.2
2	B	263	PRO	2.2
1	A	125	ARG	2.2
1	A	112	ARG	2.1
2	B	170	ASP	2.1
1	A	348	GLY	2.1
1	A	237	ARG	2.1
1	A	91	ARG	2.1
1	A	176	PHE	2.0
1	A	146	TRP	2.0
2	B	41	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	B	307	4/4	0.76	0.20	59,64,65,70	0
4	EDO	A	506	4/4	0.77	0.24	58,59,63,67	0
4	EDO	B	305	4/4	0.78	0.21	65,67,67,67	0
4	EDO	A	507	4/4	0.78	0.26	72,73,75,78	0
4	EDO	A	504	4/4	0.79	0.28	98,99,99,99	0
4	EDO	B	304	4/4	0.79	0.22	63,66,67,68	0
4	EDO	A	502	4/4	0.81	0.27	66,68,69,69	0
4	EDO	B	301	4/4	0.85	0.21	58,59,61,67	0
4	EDO	B	306	4/4	0.85	0.15	59,63,65,67	0
3	OSW	A	501	19/19	0.85	0.17	53,64,67,67	0
4	EDO	B	302	4/4	0.87	0.20	57,65,66,66	0
4	EDO	A	505	4/4	0.88	0.20	75,76,76,77	0
4	EDO	A	503	4/4	0.91	0.13	42,43,45,47	0
5	FMT	B	308	3/3	0.92	0.17	61,61,61,62	0
4	EDO	B	303	4/4	0.94	0.14	54,54,55,58	0

6.5 Other polymers

There are no such residues in this entry.