



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:01 AM UTC

PDB ID : 5F73 / pdb_00005f73
Title : Crystal structure of Mutant S12T of Adenosine/Methylthioadenosine Phosphorylase in APO form
Authors : Torini, J.R.S.; Brandao-Neto, J.; DeMarco, R.; Pereira, H.M.
Deposited on : 2015-12-07
Resolution : 2.06 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	3% 83% 6% 10%
1	B	320	2% 81% 8% 11%
1	C	320	0% 84% 12%
1	D	320	2% 84% 13%
1	E	320	2% 84% 5% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	320	 A horizontal bar chart showing the quality of chain F. The bar is divided into three segments: a small red segment at the beginning labeled '2%', a large green segment labeled '78%', and a small yellow segment at the end labeled '7%'. A grey segment at the far right is labeled '15%'. The total length of the bar represents 100%.

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13806 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methylthioadenosine phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	288	Total	C	N	O	S	0	0	0
			2151	1364	370	401	16			
1	B	286	Total	C	N	O	S	0	0	0
			2163	1370	372	404	17			
1	C	281	Total	C	N	O	S	0	2	0
			2143	1358	370	397	18			
1	D	279	Total	C	N	O	S	0	0	0
			2087	1322	361	388	16			
1	E	286	Total	C	N	O	S	0	0	0
			2139	1356	371	395	17			
1	F	272	Total	C	N	O	S	0	0	0
			2081	1319	362	385	15			

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP I0B503
A	-19	GLY	-	expression tag	UNP I0B503
A	-18	SER	-	expression tag	UNP I0B503
A	-17	SER	-	expression tag	UNP I0B503
A	-16	HIS	-	expression tag	UNP I0B503
A	-15	HIS	-	expression tag	UNP I0B503
A	-14	HIS	-	expression tag	UNP I0B503
A	-13	HIS	-	expression tag	UNP I0B503
A	-12	HIS	-	expression tag	UNP I0B503
A	-11	HIS	-	expression tag	UNP I0B503
A	-10	SER	-	expression tag	UNP I0B503
A	-9	SER	-	expression tag	UNP I0B503
A	-8	GLY	-	expression tag	UNP I0B503
A	-7	LEU	-	expression tag	UNP I0B503
A	-6	VAL	-	expression tag	UNP I0B503
A	-5	PRO	-	expression tag	UNP I0B503
A	-4	ARG	-	expression tag	UNP I0B503

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP I0B503
A	-2	SER	-	expression tag	UNP I0B503
A	-1	HIS	-	expression tag	UNP I0B503
A	0	MET	-	expression tag	UNP I0B503
A	12	THR	SER	engineered mutation	UNP I0B503
B	-20	MET	-	initiating methionine	UNP I0B503
B	-19	GLY	-	expression tag	UNP I0B503
B	-18	SER	-	expression tag	UNP I0B503
B	-17	SER	-	expression tag	UNP I0B503
B	-16	HIS	-	expression tag	UNP I0B503
B	-15	HIS	-	expression tag	UNP I0B503
B	-14	HIS	-	expression tag	UNP I0B503
B	-13	HIS	-	expression tag	UNP I0B503
B	-12	HIS	-	expression tag	UNP I0B503
B	-11	HIS	-	expression tag	UNP I0B503
B	-10	SER	-	expression tag	UNP I0B503
B	-9	SER	-	expression tag	UNP I0B503
B	-8	GLY	-	expression tag	UNP I0B503
B	-7	LEU	-	expression tag	UNP I0B503
B	-6	VAL	-	expression tag	UNP I0B503
B	-5	PRO	-	expression tag	UNP I0B503
B	-4	ARG	-	expression tag	UNP I0B503
B	-3	GLY	-	expression tag	UNP I0B503
B	-2	SER	-	expression tag	UNP I0B503
B	-1	HIS	-	expression tag	UNP I0B503
B	0	MET	-	expression tag	UNP I0B503
B	12	THR	SER	engineered mutation	UNP I0B503
C	-20	MET	-	initiating methionine	UNP I0B503
C	-19	GLY	-	expression tag	UNP I0B503
C	-18	SER	-	expression tag	UNP I0B503
C	-17	SER	-	expression tag	UNP I0B503
C	-16	HIS	-	expression tag	UNP I0B503
C	-15	HIS	-	expression tag	UNP I0B503
C	-14	HIS	-	expression tag	UNP I0B503
C	-13	HIS	-	expression tag	UNP I0B503
C	-12	HIS	-	expression tag	UNP I0B503
C	-11	HIS	-	expression tag	UNP I0B503
C	-10	SER	-	expression tag	UNP I0B503
C	-9	SER	-	expression tag	UNP I0B503
C	-8	GLY	-	expression tag	UNP I0B503
C	-7	LEU	-	expression tag	UNP I0B503
C	-6	VAL	-	expression tag	UNP I0B503

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-5	PRO	-	expression tag	UNP I0B503
C	-4	ARG	-	expression tag	UNP I0B503
C	-3	GLY	-	expression tag	UNP I0B503
C	-2	SER	-	expression tag	UNP I0B503
C	-1	HIS	-	expression tag	UNP I0B503
C	0	MET	-	expression tag	UNP I0B503
C	12	THR	SER	engineered mutation	UNP I0B503
D	-20	MET	-	initiating methionine	UNP I0B503
D	-19	GLY	-	expression tag	UNP I0B503
D	-18	SER	-	expression tag	UNP I0B503
D	-17	SER	-	expression tag	UNP I0B503
D	-16	HIS	-	expression tag	UNP I0B503
D	-15	HIS	-	expression tag	UNP I0B503
D	-14	HIS	-	expression tag	UNP I0B503
D	-13	HIS	-	expression tag	UNP I0B503
D	-12	HIS	-	expression tag	UNP I0B503
D	-11	HIS	-	expression tag	UNP I0B503
D	-10	SER	-	expression tag	UNP I0B503
D	-9	SER	-	expression tag	UNP I0B503
D	-8	GLY	-	expression tag	UNP I0B503
D	-7	LEU	-	expression tag	UNP I0B503
D	-6	VAL	-	expression tag	UNP I0B503
D	-5	PRO	-	expression tag	UNP I0B503
D	-4	ARG	-	expression tag	UNP I0B503
D	-3	GLY	-	expression tag	UNP I0B503
D	-2	SER	-	expression tag	UNP I0B503
D	-1	HIS	-	expression tag	UNP I0B503
D	0	MET	-	expression tag	UNP I0B503
D	12	THR	SER	engineered mutation	UNP I0B503
E	-20	MET	-	initiating methionine	UNP I0B503
E	-19	GLY	-	expression tag	UNP I0B503
E	-18	SER	-	expression tag	UNP I0B503
E	-17	SER	-	expression tag	UNP I0B503
E	-16	HIS	-	expression tag	UNP I0B503
E	-15	HIS	-	expression tag	UNP I0B503
E	-14	HIS	-	expression tag	UNP I0B503
E	-13	HIS	-	expression tag	UNP I0B503
E	-12	HIS	-	expression tag	UNP I0B503
E	-11	HIS	-	expression tag	UNP I0B503
E	-10	SER	-	expression tag	UNP I0B503
E	-9	SER	-	expression tag	UNP I0B503
E	-8	GLY	-	expression tag	UNP I0B503

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-7	LEU	-	expression tag	UNP I0B503
E	-6	VAL	-	expression tag	UNP I0B503
E	-5	PRO	-	expression tag	UNP I0B503
E	-4	ARG	-	expression tag	UNP I0B503
E	-3	GLY	-	expression tag	UNP I0B503
E	-2	SER	-	expression tag	UNP I0B503
E	-1	HIS	-	expression tag	UNP I0B503
E	0	MET	-	expression tag	UNP I0B503
E	12	THR	SER	engineered mutation	UNP I0B503
F	-20	MET	-	initiating methionine	UNP I0B503
F	-19	GLY	-	expression tag	UNP I0B503
F	-18	SER	-	expression tag	UNP I0B503
F	-17	SER	-	expression tag	UNP I0B503
F	-16	HIS	-	expression tag	UNP I0B503
F	-15	HIS	-	expression tag	UNP I0B503
F	-14	HIS	-	expression tag	UNP I0B503
F	-13	HIS	-	expression tag	UNP I0B503
F	-12	HIS	-	expression tag	UNP I0B503
F	-11	HIS	-	expression tag	UNP I0B503
F	-10	SER	-	expression tag	UNP I0B503
F	-9	SER	-	expression tag	UNP I0B503
F	-8	GLY	-	expression tag	UNP I0B503
F	-7	LEU	-	expression tag	UNP I0B503
F	-6	VAL	-	expression tag	UNP I0B503
F	-5	PRO	-	expression tag	UNP I0B503
F	-4	ARG	-	expression tag	UNP I0B503
F	-3	GLY	-	expression tag	UNP I0B503
F	-2	SER	-	expression tag	UNP I0B503
F	-1	HIS	-	expression tag	UNP I0B503
F	0	MET	-	expression tag	UNP I0B503
F	12	THR	SER	engineered mutation	UNP I0B503

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	F	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	126	Total	O	0	0
			126	126		
3	B	176	Total	O	0	0
			176	176		
3	C	221	Total	O	0	0
			221	221		
3	D	158	Total	O	0	0
			158	158		
3	E	150	Total	O	0	0
			150	150		
3	F	181	Total	O	0	0
			181	181		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Methylthioadenosine phosphorylase




• Molecule 1: Methylthioadenosine phosphorylase




• Molecule 1: Methylthioadenosine phosphorylase




• Molecule 1: Methylthioadenosine phosphorylase








● Molecule 1: Methylthioadenosine phosphorylase



● Molecule 1: Methylthioadenosine phosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.44Å 82.16Å 150.91Å 90.00° 101.50° 90.00°	Depositor
Resolution (Å)	39.58 – 2.06 39.58 – 2.06	Depositor EDS
% Data completeness (in resolution range)	98.6 (39.58-2.06) 98.6 (39.58-2.06)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 2.05Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.175 , 0.216 0.176 , 0.216	Depositor DCC
R_{free} test set	6075 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtrriage
Anisotropy	0.348	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13806	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.66 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.2844e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/2195	0.43	0/2986
1	B	0.30	0/2206	0.47	0/2999
1	C	0.36	0/2190	0.50	0/2971
1	D	0.30	0/2127	0.45	0/2891
1	E	0.28	0/2182	0.45	0/2968
1	F	0.31	0/2119	0.46	0/2870
All	All	0.31	0/13019	0.46	0/17685

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2151	0	2092	9	0
1	B	2163	0	2124	12	0
1	C	2143	0	2140	7	0
1	D	2087	0	2040	5	0
1	E	2139	0	2088	9	0
1	F	2081	0	2094	14	0
2	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	126	0	0	0	0
3	B	176	0	0	4	0
3	C	221	0	0	0	0
3	D	158	0	0	0	0
3	E	150	0	0	2	0
3	F	181	0	0	3	0
All	All	13806	0	12578	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:131:HIS:HD2	1:C:289:GLN:HG3	1.42	0.85
3:E:529:HOH:O	1:F:191:CYS:SG	2.39	0.79
3:B:542:HOH:O	1:C:191:CYS:SG	2.45	0.73
1:F:131:HIS:HD2	1:F:289:GLN:HG3	1.58	0.69
1:E:162:THR:HG23	1:E:163:MET:HG3	1.75	0.69
1:B:132:MET:HE2	1:B:216:ARG:HG2	1.75	0.67
1:E:131:HIS:HD2	1:E:289:GLN:HG3	1.59	0.67
1:B:18:ASN:O	1:B:258:ARG:NH1	2.30	0.64
1:B:2:SER:N	3:B:403:HOH:O	2.31	0.64
1:F:26:ARG:NH2	1:F:41:GLU:OE2	2.32	0.63
1:F:180:VAL:HB	1:F:201:LEU:HD13	1.80	0.62
1:E:132:MET:HE2	1:E:216:ARG:HG2	1.80	0.62
1:F:132:MET:HE2	1:F:216:ARG:HG2	1.82	0.62
1:F:271:GLU:OE1	3:F:401:HOH:O	2.16	0.61
1:B:27:GLN:NE2	3:B:404:HOH:O	2.34	0.61
1:E:227:ILE:HD11	1:E:257:VAL:HG11	1.82	0.60
1:B:17:PRO:HB2	1:B:19:LEU:HD13	1.85	0.59
1:E:253:SER:O	1:E:257:VAL:HG12	2.03	0.59
1:B:104:LEU:HD21	1:B:203:ILE:HD11	1.85	0.58
1:C:131:HIS:CD2	1:C:289:GLN:HG3	2.32	0.57
1:B:5:LYS:NZ	1:B:41:GLU:OE2	2.30	0.56
1:F:76:LYS:HD2	1:F:218:ALA:HB1	1.88	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:164:ASP:OD1	1:F:164:ASP:N	2.24	0.55
1:D:239:GLU:O	1:D:243:VAL:HG23	2.08	0.54
1:A:169:ILE:O	1:A:172:CYS:HB3	2.06	0.54
1:D:132:MET:HE3	1:D:133:PRO:HD2	1.87	0.54
1:A:132:MET:HE3	1:A:133:PRO:HD2	1.92	0.52
1:B:108:MET:HE1	1:B:199:MET:SD	2.49	0.52
1:C:104:LEU:HD21	1:C:203:ILE:HD11	1.91	0.52
1:A:162:THR:HG23	1:A:163:MET:HG3	1.93	0.50
1:F:89:CYS:SG	1:F:203:ILE:HD12	2.52	0.50
1:A:164:ASP:OD2	1:A:166:SER:OG	2.24	0.49
1:E:143:ARG:NH2	1:E:175:ALA:O	2.46	0.48
1:F:132:MET:HE1	1:F:283:LEU:HG	1.95	0.48
1:C:89[A]:CYS:SG	1:C:203:ILE:HD12	2.54	0.47
1:B:180:VAL:HB	1:B:201:LEU:HD13	1.97	0.46
1:A:162:THR:OG1	1:E:258:ARG:NE	2.48	0.46
1:D:132:MET:HE2	1:D:132:MET:HB3	1.70	0.46
1:E:145:ILE:HG21	1:E:268:ILE:HG13	1.98	0.46
1:F:57:LYS:NZ	3:F:413:HOH:O	2.48	0.45
1:A:141:ARG:O	1:A:145:ILE:HG12	2.17	0.45
1:C:17:PRO:HG2	1:C:51:VAL:HG11	1.98	0.45
1:F:121:SER:HA	1:F:127:LYS:NZ	2.32	0.45
1:F:104:LEU:HD21	1:F:203:ILE:HD11	1.99	0.44
1:C:108:MET:HB3	1:C:201:LEU:HD21	2.00	0.44
1:D:93:GLN:OE1	1:D:197:LYS:NZ	2.51	0.43
1:B:241:VAL:HG23	1:B:245:MET:HE2	2.00	0.43
1:B:169:ILE:O	1:B:172:CYS:HB3	2.19	0.43
1:A:251:ARG:O	1:A:254:VAL:HG22	2.19	0.43
1:B:124:ASP:OD2	3:B:401:HOH:O	2.21	0.43
1:E:258:ARG:HD2	3:E:533:HOH:O	2.18	0.42
1:A:227:ILE:HD11	1:A:257:VAL:HG21	2.00	0.42
1:F:258:ARG:NH2	3:F:419:HOH:O	2.53	0.42
1:D:132:MET:HE1	1:D:283:LEU:HG	2.03	0.40
1:A:195:ILE:HG22	1:A:199:MET:HE2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/320 (89%)	281 (98%)	4 (1%)	1 (0%)	36	30
1	B	282/320 (88%)	279 (99%)	2 (1%)	1 (0%)	30	23
1	C	279/320 (87%)	277 (99%)	2 (1%)	0	100	100
1	D	273/320 (85%)	271 (99%)	2 (1%)	0	100	100
1	E	282/320 (88%)	277 (98%)	5 (2%)	0	100	100
1	F	266/320 (83%)	265 (100%)	1 (0%)	0	100	100
All	All	1668/1920 (87%)	1650 (99%)	16 (1%)	2 (0%)	48	44

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	162	THR
1	A	21	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/274 (83%)	222 (98%)	5 (2%)	45	43
1	B	234/274 (85%)	227 (97%)	7 (3%)	36	32
1	C	235/274 (86%)	233 (99%)	2 (1%)	70	74
1	D	222/274 (81%)	219 (99%)	3 (1%)	59	60
1	E	227/274 (83%)	222 (98%)	5 (2%)	45	43

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	228/274 (83%)	225 (99%)	3 (1%)	61	62
All	All	1373/1644 (84%)	1348 (98%)	25 (2%)	51	51

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	LEU
1	A	22	LYS
1	A	25	VAL
1	A	50	VAL
1	A	132	MET
1	B	4	VAL
1	B	27	GLN
1	B	52	LEU
1	B	77	ASP
1	B	97	VAL
1	B	241	VAL
1	B	275	LYS
1	C	108	MET
1	C	233	CYS
1	D	50	VAL
1	D	132	MET
1	D	205	ASN
1	E	77	ASP
1	E	97	VAL
1	E	164	ASP
1	E	205	ASN
1	E	257	VAL
1	F	77	ASP
1	F	164	ASP
1	F	285	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	148	GLN
1	C	256	HIS
1	D	156	ASN
1	E	59	HIS
1	E	249	GLN
1	F	256	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	301	-	4,4,4	0.21	0	6,6,6	0.26	0
2	SO4	D	301	-	4,4,4	0.30	0	6,6,6	0.28	0
2	SO4	F	301	-	4,4,4	0.40	0	6,6,6	0.13	0
2	SO4	A	301	-	4,4,4	0.31	0	6,6,6	0.23	0
2	SO4	C	301	-	4,4,4	0.32	0	6,6,6	0.25	0
2	SO4	E	301	-	4,4,4	0.34	0	6,6,6	0.24	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	288/320 (90%)	-0.10	9 (3%)	51	52	19, 34, 70, 92	0
1	B	286/320 (89%)	-0.22	6 (2%)	63	66	20, 30, 63, 79	0
1	C	281/320 (87%)	-0.34	4 (1%)	73	75	16, 26, 49, 79	2 (0%)
1	D	279/320 (87%)	-0.28	7 (2%)	58	60	17, 30, 56, 89	0
1	E	286/320 (89%)	-0.07	8 (2%)	55	56	21, 34, 68, 83	0
1	F	272/320 (85%)	-0.31	5 (1%)	67	69	18, 28, 49, 99	0
All	All	1692/1920 (88%)	-0.22	39 (2%)	61	63	16, 30, 61, 99	2 (0%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	25	VAL	4.9
1	E	291	LEU	4.4
1	B	291	LEU	4.0
1	A	23	VAL	3.4
1	B	241	VAL	3.4
1	D	237	GLU	3.4
1	E	241	VAL	3.2
1	A	17	PRO	3.1
1	B	240	HIS	3.1
1	D	240	HIS	3.1
1	D	234	TRP	3.0
1	C	17	PRO	3.0
1	D	124	ASP	3.0
1	C	233	CYS	2.9
1	E	2	SER	2.7
1	F	27	GLN	2.7
1	D	243	VAL	2.7
1	E	236	SER	2.6
1	C	18	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	167	ALA	2.6
1	E	162	THR	2.6
1	A	166	SER	2.6
1	E	169	ILE	2.5
1	B	235	LYS	2.4
1	A	19	LEU	2.4
1	D	241	VAL	2.4
1	B	2	SER	2.3
1	B	18	ASN	2.3
1	F	243	VAL	2.3
1	D	232	ASP	2.3
1	F	15	ASP	2.3
1	A	22	LYS	2.2
1	A	291	LEU	2.2
1	E	49	CYS	2.1
1	E	245	MET	2.1
1	C	232	ASP	2.0
1	A	240	HIS	2.0
1	A	234	TRP	2.0
1	F	232	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	301	5/5	0.98	0.06	25,25,29,29	0
2	SO4	C	301	5/5	0.98	0.05	20,20,22,24	0
2	SO4	E	301	5/5	0.98	0.05	26,27,29,30	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	F	301	5/5	0.98	0.05	24,24,27,27	0
2	SO4	A	301	5/5	0.99	0.05	21,21,27,28	0
2	SO4	D	301	5/5	0.99	0.04	21,22,22,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.