



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 09:36 PM UTC

PDB ID : 5F7W / pdb_00005f7w
Title : Blood group antigen binding adhesin BabA of Helicobacter pylori strain 17875
in complex with blood group B Lewis b heptasaccharide
Authors : Moonens, K.; Gideonsson, P.; Subedi, S.; Romao, E.; Oscarson, S.; Muylder-
mans, S.; Boren, T.; Remaut, H.
Deposited on : 2015-12-08
Resolution : 2.81 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4591 (2.84-2.80)
Clashscore	190562	5010 (2.84-2.80)
Ramachandran outliers	187476	4916 (2.84-2.80)
Sidechain outliers	187428	4918 (2.84-2.80)
RSRZ outliers	180081	4594 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	466	82% 8% 10%
1	B	466	80% 9% 10%
2	C	120	89% 5% 6%
2	D	120	87% 8% 5%
3	E	7	14% 86%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	7	 71% 29%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8176 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adhesin binding fucosylated histo-blood group antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	419	Total	C	N	O	S	0	0	0
			3114	1920	537	644	13			
1	B	419	Total	C	N	O	S	0	0	0
			3120	1924	539	644	13			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	ALA	-	expression tag	UNP O52269
A	4	SER	-	expression tag	UNP O52269
A	5	TRP	-	expression tag	UNP O52269
A	6	SER	-	expression tag	UNP O52269
A	7	HIS	-	expression tag	UNP O52269
A	8	PRO	-	expression tag	UNP O52269
A	9	GLN	-	expression tag	UNP O52269
A	10	PHE	-	expression tag	UNP O52269
A	11	GLU	-	expression tag	UNP O52269
A	12	LYS	-	expression tag	UNP O52269
A	13	SER	-	expression tag	UNP O52269
A	14	GLY	-	expression tag	UNP O52269
A	15	GLY	-	expression tag	UNP O52269
A	16	GLY	-	expression tag	UNP O52269
A	17	GLY	-	expression tag	UNP O52269
A	18	GLY	-	expression tag	UNP O52269
A	19	LEU	-	expression tag	UNP O52269
A	20	VAL	-	expression tag	UNP O52269
A	21	PRO	-	expression tag	UNP O52269
A	22	ARG	-	expression tag	UNP O52269
A	23	GLY	-	expression tag	UNP O52269
A	24	SER	-	expression tag	UNP O52269
A	461	GLY	-	expression tag	UNP O52269
A	462	SER	-	expression tag	UNP O52269
A	463	HIS	-	expression tag	UNP O52269

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	464	HIS	-	expression tag	UNP O52269
A	465	HIS	-	expression tag	UNP O52269
A	466	HIS	-	expression tag	UNP O52269
A	467	HIS	-	expression tag	UNP O52269
A	468	HIS	-	expression tag	UNP O52269
B	3	ALA	-	expression tag	UNP O52269
B	4	SER	-	expression tag	UNP O52269
B	5	TRP	-	expression tag	UNP O52269
B	6	SER	-	expression tag	UNP O52269
B	7	HIS	-	expression tag	UNP O52269
B	8	PRO	-	expression tag	UNP O52269
B	9	GLN	-	expression tag	UNP O52269
B	10	PHE	-	expression tag	UNP O52269
B	11	GLU	-	expression tag	UNP O52269
B	12	LYS	-	expression tag	UNP O52269
B	13	SER	-	expression tag	UNP O52269
B	14	GLY	-	expression tag	UNP O52269
B	15	GLY	-	expression tag	UNP O52269
B	16	GLY	-	expression tag	UNP O52269
B	17	GLY	-	expression tag	UNP O52269
B	18	GLY	-	expression tag	UNP O52269
B	19	LEU	-	expression tag	UNP O52269
B	20	VAL	-	expression tag	UNP O52269
B	21	PRO	-	expression tag	UNP O52269
B	22	ARG	-	expression tag	UNP O52269
B	23	GLY	-	expression tag	UNP O52269
B	24	SER	-	expression tag	UNP O52269
B	461	GLY	-	expression tag	UNP O52269
B	462	SER	-	expression tag	UNP O52269
B	463	HIS	-	expression tag	UNP O52269
B	464	HIS	-	expression tag	UNP O52269
B	465	HIS	-	expression tag	UNP O52269
B	466	HIS	-	expression tag	UNP O52269
B	467	HIS	-	expression tag	UNP O52269
B	468	HIS	-	expression tag	UNP O52269

- Molecule 2 is a protein called Nanobody Nb-ER19.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	113	Total	C	N	O	S	0	0	0
			863	540	155	163	5			
2	D	114	Total	C	N	O	S	0	0	0
			873	546	158	164	5			

- Molecule 3 is an oligosaccharide called alpha-L-fucopyranose-(1-2)-[alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	7	Total	C	N	O	0	0	0
			79	44	1	34			
3	F	7	Total	C	N	O	0	0	0
			79	44	1	34			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	19	Total	O	0	0
			19	19		
4	B	14	Total	O	0	0
			14	14		
4	C	3	Total	O	0	0
			3	3		
4	D	12	Total	O	0	0
			12	12		



- Molecule 3: alpha-L-fucopyranose-(1-2)-[alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain E: 14% 86%



- Molecule 3: alpha-L-fucopyranose-(1-2)-[alpha-D-galactopyranose-(1-3)]beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-beta-D-galactopyranose-(1-4)-beta-D-glucopyranose

Chain F: 71% 29%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.09Å 132.44Å 123.51Å 90.00° 94.93° 90.00°	Depositor
Resolution (Å)	43.20 – 2.81 43.20 – 2.81	Depositor EDS
% Data completeness (in resolution range)	99.6 (43.20-2.81) 99.7 (43.20-2.81)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.171 , 0.208 (Not available) , 0.204	Depositor DCC
R_{free} test set	2026 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	70.0	Xtriage
Anisotropy	0.100	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8176	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, GAL, GLA, NAG, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.02	0/3160	1.01	2/4295 (0.0%)
1	B	0.94	2/3166 (0.1%)	0.99	4/4302 (0.1%)
2	C	0.97	0/882	0.97	0/1195
2	D	1.01	0/893	1.00	0/1210
All	All	0.98	2/8101 (0.0%)	1.00	6/11002 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	218	ILE	CA-C	5.44	1.59	1.52
1	B	277	CYS	C-O	-5.36	1.21	1.23

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	79	ASN	N-CA-C	5.59	120.09	113.16
1	A	79	ASN	N-CA-C	5.06	119.44	113.16
1	B	273	ILE	CB-CA-C	-5.06	105.40	111.88
1	A	105	GLN	N-CA-CB	5.01	117.63	110.06
1	B	288	GLY	CA-C-N	5.00	126.09	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3114	0	3044	14	0
1	B	3120	0	3047	19	0
2	C	863	0	845	1	0
2	D	873	0	852	3	0
3	E	79	0	69	0	0
3	F	79	0	69	2	0
4	A	19	0	0	0	0
4	B	14	0	0	0	0
4	C	3	0	0	0	0
4	D	12	0	0	1	0
All	All	8176	0	7926	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 37 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:219:ASP:O	1:B:221:LYS:N	2.21	0.74
1:A:35:ASN:OD1	1:A:36:LEU:N	2.21	0.73
1:B:35:ASN:OD1	1:B:36:LEU:N	2.25	0.70
1:A:35:ASN:OD1	1:A:35:ASN:C	2.38	0.67
1:B:35:ASN:OD1	1:B:35:ASN:C	2.41	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	415/466 (89%)	394 (95%)	17 (4%)	4 (1%)	12 36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	413/466 (89%)	391 (95%)	18 (4%)	4 (1%)	12	36
2	C	111/120 (92%)	106 (96%)	5 (4%)	0	100	100
2	D	112/120 (93%)	106 (95%)	6 (5%)	0	100	100
All	All	1051/1172 (90%)	997 (95%)	46 (4%)	8 (1%)	16	41

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	338	ASN
1	A	288	GLY
1	B	338	ASN
1	B	240	THR
1	B	288	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/381 (91%)	331 (96%)	14 (4%)	27	60
1	B	346/381 (91%)	332 (96%)	14 (4%)	28	61
2	C	92/99 (93%)	88 (96%)	4 (4%)	26	58
2	D	93/99 (94%)	88 (95%)	5 (5%)	20	50
All	All	876/960 (91%)	839 (96%)	37 (4%)	26	59

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	14	GLN
2	D	103	ARG
2	C	29	ILE
2	D	29	ILE
1	A	394	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	396	ASN
2	D	40	GLN
1	B	260	GLN
1	B	265	GLN
1	B	274	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	BGC	E	1	3	12,12,12	0.59	0	17,17,17	1.32	3 (17%)
3	GAL	E	2	3	11,11,12	0.47	0	15,15,17	1.55	3 (20%)
3	NAG	E	3	3	14,14,15	0.72	0	17,19,21	1.36	4 (23%)
3	GAL	E	4	3	11,11,12	1.09	2 (18%)	15,15,17	1.97	5 (33%)
3	FUC	E	5	3	10,10,11	0.87	0	14,14,16	1.02	1 (7%)
3	GLA	E	6	3	11,11,12	0.66	0	15,15,17	1.91	4 (26%)
3	FUC	E	7	3	10,10,11	0.83	0	14,14,16	1.12	0
3	BGC	F	1	3	12,12,12	0.49	0	17,17,17	1.13	1 (5%)
3	GAL	F	2	3	11,11,12	0.76	0	15,15,17	1.89	4 (26%)
3	NAG	F	3	3	14,14,15	0.66	0	17,19,21	1.53	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GAL	F	4	3	11,11,12	0.91	0	15,15,17	1.64	3 (20%)
3	FUC	F	5	3	10,10,11	0.77	0	14,14,16	1.22	2 (14%)
3	GLA	F	6	3	11,11,12	0.86	0	15,15,17	1.27	1 (6%)
3	FUC	F	7	3	10,10,11	0.65	0	14,14,16	1.33	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	E	1	3	-	2/2/22/22	0/1/1/1
3	GAL	E	2	3	-	2/2/19/22	0/1/1/1
3	NAG	E	3	3	-	0/6/23/26	0/1/1/1
3	GAL	E	4	3	-	2/2/19/22	0/1/1/1
3	FUC	E	5	3	-	-	0/1/1/1
3	GLA	E	6	3	-	2/2/19/22	0/1/1/1
3	FUC	E	7	3	-	-	0/1/1/1
3	BGC	F	1	3	-	0/2/22/22	0/1/1/1
3	GAL	F	2	3	-	0/2/19/22	0/1/1/1
3	NAG	F	3	3	-	0/6/23/26	0/1/1/1
3	GAL	F	4	3	-	2/2/19/22	0/1/1/1
3	FUC	F	5	3	-	-	0/1/1/1
3	GLA	F	6	3	-	1/2/19/22	0/1/1/1
3	FUC	F	7	3	-	-	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	4	GAL	O5-C1	-2.26	1.39	1.43
3	E	4	GAL	O5-C5	-2.06	1.39	1.43

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	6	GLA	C1-O5-C5	5.12	119.05	112.19
3	E	4	GAL	C1-O5-C5	-4.98	105.51	112.19
3	F	2	GAL	C1-C2-C3	4.20	115.75	109.64
3	F	4	GAL	C1-O5-C5	4.19	117.81	112.19
3	E	2	GAL	C1-O5-C5	3.58	116.98	112.19

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	2	GAL	C4-C5-C6-O6
3	E	4	GAL	C4-C5-C6-O6
3	F	4	GAL	O5-C5-C6-O6
3	F	4	GAL	C4-C5-C6-O6
3	E	2	GAL	O5-C5-C6-O6


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	5	FUC	1	0
3	F	6	GLA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain E





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	419/466 (89%)	-0.29	5 (1%) 76 68	43, 66, 119, 143	0
1	B	419/466 (89%)	-0.16	6 (1%) 73 64	49, 75, 129, 153	0
2	C	113/120 (94%)	-0.45	2 (1%) 67 58	49, 68, 93, 121	0
2	D	114/120 (95%)	-0.46	1 (0%) 81 74	46, 59, 83, 114	0
All	All	1065/1172 (90%)	-0.27	14 (1%) 75 66	43, 69, 120, 153	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	3	VAL	4.9
2	C	3	VAL	4.9
1	B	218	ILE	4.2
1	A	218	ILE	3.2
1	A	398	LYS	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BGC	F	1	12/12	0.55	0.15	68,75,76,77	0
3	GAL	F	4	11/12	0.64	0.16	40,43,45,45	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GLA	F	6	11/12	0.64	0.15	41,44,45,46	0
3	BGC	E	1	12/12	0.65	0.14	60,74,83,87	0
3	GAL	F	2	11/12	0.67	0.14	49,59,63,65	0
3	GAL	E	2	11/12	0.71	0.12	56,63,68,71	0
3	FUC	F	7	10/11	0.71	0.19	40,43,45,45	0
3	NAG	F	3	14/15	0.74	0.14	35,45,46,46	0
3	GAL	E	4	11/12	0.78	0.13	41,43,45,45	0
3	FUC	F	5	10/11	0.79	0.13	39,41,44,46	0
3	FUC	E	7	10/11	0.81	0.15	36,41,44,44	0
3	GLA	E	6	11/12	0.88	0.10	45,48,49,51	0
3	NAG	E	3	14/15	0.89	0.11	42,46,50,52	0
3	FUC	E	5	10/11	0.94	0.09	30,34,37,38	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.