



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:43 PM UTC

PDB ID : 4F8R / pdb_00004f8r
Title : Bacillus DNA Polymerase I Large Fragment complex 7
Authors : Wang, W.; Beese, L.S.
Deposited on : 2012-05-17
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	1141 (1.64-1.64)
Clashscore	190562	1171 (1.64-1.64)
Ramachandran outliers	187476	1151 (1.64-1.64)
Sidechain outliers	187428	1150 (1.64-1.64)
RSRZ outliers	180081	1141 (1.64-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	592	2% 93% 5%
1	D	592	2% 88% 5% 7%
2	B	9	11% 44% 56%
2	E	9	67% 33%
3	C	13	54% 23% 8% 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	13	 92%8%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 21034 atoms, of which 9790 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	579	Total	C	H	N	O	S	0	6	0
			9416	2968	4744	814	873	17			
1	D	553	Total	C	H	N	O	S	0	18	0
			9124	2884	4598	781	845	16			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	598	ALA	ASP	engineered mutation	UNP Q5KWC1
A	710	TYR	PHE	engineered mutation	UNP Q5KWC1
A	823	HIS	ARG	SEE REMARK 999	UNP Q5KWC1
D	598	ALA	ASP	engineered mutation	UNP Q5KWC1
D	710	TYR	PHE	engineered mutation	UNP Q5KWC1
D	823	HIS	ARG	SEE REMARK 999	UNP Q5KWC1


- Molecule 2 is a DNA chain called DNA (5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(DDG))-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	9	Total	C	H	N	O	P	0	0	0
			276	86	100	31	51	8			
2	E	9	Total	C	H	N	O	P	0	0	0
			277	86	100	31	52	8			

- Molecule 3 is a DNA chain called DNA (5'-D(*CP*AP*TP*TP*CP*GP*AP*GP*TP*CP*AP*GP*G)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	11	Total	C	H	N	O	P	0	0	0
			341	108	113	42	67	11			
3	F	12	Total	C	H	N	O	P	0	0	0
			381	118	135	47	70	11			

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	406	Total	O	0	0
			406	406		
5	D	620	Total	O	0	0
			620	620		
5	B	29	Total	O	0	0
			29	29		
5	C	59	Total	O	0	0
			59	59		
5	E	26	Total	O	0	0
			26	26		
5	F	54	Total	O	0	0
			54	54		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase




- Molecule 1: DNA polymerase




- Molecule 2: DNA (5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(DDG))-3')



- Molecule 2: DNA (5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(DDG))-3')



- Molecule 3: DNA (5'-D(*CP*AP*TP*TP*CP*GP*AP*GP*TP*CP*AP*GP*G)-3')



- Molecule 3: DNA (5'-D(*CP*AP*TP*TP*CP*GP*AP*GP*TP*CP*AP*GP*G)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	93.67Å 109.37Å 150.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.98 – 1.64 30.98 – 1.64	Depositor EDS
% Data completeness (in resolution range)	87.7 (30.98-1.64) 87.7 (30.98-1.64)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.60 (at 1.64Å)	Xtriage
Refinement program	PHENIX dev_1026	Depositor
R, R_{free}	0.181 , 0.203 0.185 , 0.206	Depositor DCC
R_{free} test set	7474 reflections (4.52%)	wwPDB-VP
Wilson B-factor (Å ²)	20.0	Xtriage
Anisotropy	0.223	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 50.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	21034	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DDG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/4785	0.75	0/6466
1	D	0.74	0/4672	0.83	0/6313
2	B	0.47	0/173	1.22	1/264 (0.4%)
2	E	0.53	0/173	1.32	1/264 (0.4%)
3	C	0.53	0/255	1.47	5/392 (1.3%)
3	F	0.59	0/276	1.29	0/425
All	All	0.65	0/10334	0.86	7/14124 (0.0%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	5	DG	OP2-P-O3'	-5.88	90.35	108.00
2	B	26	DC	N1-C1'-C2'	-5.69	104.97	113.50
3	C	3	DT	C4'-C3'-O3'	-5.66	101.51	110.00
3	C	5	DG	C4'-C3'-O3'	5.49	118.23	110.00
3	C	4	DC	N1-C1'-C2'	-5.46	105.31	113.50
3	C	5	DG	O5'-C5'-C4'	5.17	118.55	110.80
2	E	23	DT	N1-C1'-C2'	5.10	121.15	113.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4672	4744	4706	22	0
1	D	4526	4598	4511	22	0
2	B	176	100	103	3	0
2	E	177	100	103	1	0
3	C	228	113	125	4	0
3	F	246	135	137	0	0
4	A	15	0	0	0	0
4	D	10	0	0	0	0
5	A	406	0	0	8	0
5	B	29	0	0	0	0
5	C	59	0	0	1	0
5	D	620	0	0	10	1
5	E	26	0	0	0	0
5	F	54	0	0	0	0
All	All	11244	9790	9685	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:502:GLN:NE2	5:D:1603:HOH:O	2.10	0.84
1:A:845:CYS:SG	5:A:1386:HOH:O	2.36	0.83
1:A:408:ASP:OD1	5:A:1097:HOH:O	1.96	0.81
1:A:466:ARG:NH2	5:A:1403:HOH:O	2.13	0.81
1:D:517:ARG:NH1	5:D:1144:HOH:O	2.20	0.74
1:A:435:ARG:NH1	5:A:1229:HOH:O	2.19	0.74
1:D:607:ASN:OD1	5:D:1378:HOH:O	2.07	0.72
1:A:607[A]:ASN:OD1	5:A:1094:HOH:O	2.07	0.71
1:D:309:GLU:OE2	5:D:1598:HOH:O	2.12	0.67
1:D:873:TYR:O	1:D:876:LYS:HD3	1.95	0.65
1:D:386:GLU:OE1	5:D:1243:HOH:O	2.14	0.65
3:C:5:DG:H2'	5:C:150:HOH:O	1.99	0.62
1:A:365:THR:O	5:A:1338:HOH:O	2.17	0.58
1:A:665:ILE:HG22	1:A:749:TYR:CE1	2.39	0.57
3:C:5:DG:C2'	3:C:6:DA:OP2	2.51	0.56
1:D:730:LYS:NZ	1:D:734:GLU:OE2	2.38	0.56
1:A:405:GLN:HG3	1:A:416:MET:HE3	1.92	0.51
1:D:764[B]:THR:CG2	1:D:768:HIS:HA	2.41	0.50
1:D:665:ILE:HG22	1:D:749[A]:TYR:CE1	2.47	0.50
1:A:411:ALA:O	1:A:415:LYS:HG2	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:764[B]:THR:HG21	5:D:1396:HOH:O	2.13	0.49
1:A:559:ASP:OD1	1:A:563:LYS:NZ	2.45	0.48
1:D:725:LEU:O	1:D:726:ASN:HB2	2.13	0.48
1:D:459:ARG:NH1	5:D:1328:HOH:O	2.38	0.47
1:D:708:VAL:HG12	1:D:712:ILE:CD1	2.45	0.47
1:A:754:VAL:O	1:A:758:LYS:HG3	2.16	0.46
1:D:645:SER:O	1:D:646:ASP:HB2	2.15	0.46
1:A:500:LEU:HD21	1:A:591:LEU:HD23	1.98	0.45
1:D:500:LEU:HD11	1:D:591:LEU:HD23	1.98	0.45
2:B:21:DC:H2'	2:B:22:DC:C6	2.51	0.45
1:A:578:ARG:NH2	2:B:25:DA:H5''	2.32	0.45
1:A:613:THR:CG2	3:C:5:DG:H4'	2.47	0.45
1:A:848:VAL:HB	1:A:849:PRO:HD3	1.99	0.45
1:A:725:LEU:HB2	1:A:727:ILE:HG12	1.99	0.45
1:D:712:ILE:HA	1:D:716:ILE:HG22	1.98	0.44
1:D:848:VAL:HB	1:D:849:PRO:HD3	2.00	0.43
1:D:708:VAL:HG12	1:D:712:ILE:HD11	1.99	0.43
1:D:873:TYR:CE2	1:D:876:LYS:HE2	2.53	0.43
2:B:22:DC:H2'	2:B:23:DT:H72	2.00	0.42
1:A:550:THR:CG2	5:A:1245:HOH:O	2.67	0.42
1:D:515:GLU:HG2	1:D:519:TYR:CE2	2.54	0.42
1:A:467[B]:ARG:HA	5:D:1501:HOH:O	2.19	0.42
1:A:467[A]:ARG:HA	5:D:1501:HOH:O	2.19	0.41
1:D:839[A]:GLU:OE2	5:D:1512:HOH:O	2.21	0.41
1:D:582:LYS:HE2	2:E:26:DC:O2	2.20	0.41
1:A:613:THR:HG21	3:C:5:DG:H4'	2.03	0.40
1:A:515:GLU:HG2	1:A:519:TYR:CE2	2.57	0.40
1:A:665:ILE:CG2	1:A:749:TYR:CE1	3.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:D:1474:HOH:O	5:D:1565:HOH:O[4_545]	2.16	0.04

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	583/592 (98%)	571 (98%)	12 (2%)	0	100	100
1	D	567/592 (96%)	557 (98%)	9 (2%)	1 (0%)	43	27
All	All	1150/1184 (97%)	1128 (98%)	21 (2%)	1 (0%)	48	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	628	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	501/507 (99%)	498 (99%)	3 (1%)	78	66
1	D	490/507 (97%)	487 (99%)	3 (1%)	78	66
All	All	991/1014 (98%)	985 (99%)	6 (1%)	78	66

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	325	GLU
1	A	477	LEU
1	A	544	LEU
1	D	303	LEU
1	D	309	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	708	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	724	ASN
1	D	405	GLN
1	D	418	GLN
1	D	755	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DDG	B	29	2,3	19,22,24	0.80	0	27,31,36	0.75	0
2	DDG	E	29	2,3	20,23,24	0.84	0	27,33,36	1.12	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DDG	B	29	2,3	-	0/6/17/19	0/3/3/3
2	DDG	E	29	2,3	-	0/7/18/19	0/3/3/3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	E	29	DDG	O4'-C1'-N9	2.87	112.95	107.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	902	-	4,4,4	0.34	0	6,6,6	0.23	0
4	SO4	A	903	-	4,4,4	0.28	0	6,6,6	0.08	0
4	SO4	D	901	-	4,4,4	0.23	0	6,6,6	0.36	0
4	SO4	D	902	-	4,4,4	0.28	0	6,6,6	0.26	0
4	SO4	A	901	-	4,4,4	0.41	0	6,6,6	0.12	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	579/592 (97%)	0.32	14 (2%) 59 65	12, 38, 65, 82	3 (0%)
1	D	553/592 (93%)	-0.24	11 (1%) 65 71	8, 25, 49, 62	10 (1%)
2	B	8/9 (88%)	0.18	1 (12%) 8 9	23, 35, 54, 70	0
2	E	8/9 (88%)	0.13	0 100 100	21, 36, 46, 55	0
3	C	11/13 (84%)	0.14	0 100 100	23, 33, 44, 56	0
3	F	12/13 (92%)	0.02	0 100 100	19, 30, 65, 78	0
All	All	1171/1228 (95%)	0.05	26 (2%) 62 68	8, 33, 60, 82	13 (1%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	712	ILE	4.7
1	D	708	VAL	3.7
1	D	735[A]	PHE	3.3
1	D	713	VAL	3.2
1	D	502	GLN	2.8
1	A	816	LYS	2.7
1	A	844	LEU	2.7
1	D	710	TYR	2.7
1	A	742[A]	SER	2.6
1	A	819	ARG	2.6
1	A	647	TRP	2.5
1	D	298	LYS	2.5
1	A	433	ALA	2.5
1	D	681	ILE	2.4
1	A	836	ALA	2.3
1	D	680	ASP	2.3
1	A	695	ASP	2.2
1	A	698	THR	2.2
1	D	782	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	727	ILE	2.2
1	A	298	LYS	2.2
1	A	355	PRO	2.2
1	A	459	ARG	2.1
1	D	781	PHE	2.1
2	B	21	DC	2.0
1	A	679	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DDG	B	29	20/22	0.90	0.09	19,23,29,35	0
2	DDG	E	29	21/22	0.97	0.06	14,20,25,27	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	D	902	5/5	0.79	0.12	70,73,78,81	0
4	SO4	A	903	5/5	0.81	0.10	53,62,70,75	0
4	SO4	A	902	5/5	0.89	0.10	41,48,58,65	0
4	SO4	A	901	5/5	0.91	0.10	36,39,48,50	0
4	SO4	D	901	5/5	0.95	0.08	37,38,43,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.