



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 10:28 AM UTC

PDB ID : 1F93 / pdb_00001f93
Title : CRYSTAL STRUCTURE OF A COMPLEX BETWEEN THE DIMERIZATION DOMAIN OF HNF-1 ALPHA AND THE COACTIVATOR DCOH
Authors : Rose, R.B.; Bayle, J.H.; Endrizzi, J.A.; Cronk, J.D.; Crabtree, G.R.; Alber, T.
Deposited on : 2000-07-06
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	104	3% 80% 18% ..
1	B	104	% 81% 13% 5%
1	C	104	15% 55% 37% 5%
1	D	104	48% 49% 38% 7% ..
2	E	32	16% 47% 50% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	32	
2	G	32	
2	H	32	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4106 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DIMERIZATION COFACTOR OF HEPATOCYTE NUCLEAR FACTOR 1-ALPHA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	103	Total	C	N	O	S	Se	0	0	0
			831	526	151	151	1	2			
1	B	99	Total	C	N	O	S	Se	0	0	0
			799	509	144	143	1	2			
1	C	99	Total	C	N	O	S	Se	0	0	0
			806	510	146	147	1	2			
1	D	100	Total	C	N	O	S	Se	0	0	0
			797	506	145	143	1	2			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP P61459
A	50	MSE	MET	modified residue	UNP P61459
A	103	MSE	MET	modified residue	UNP P61459
B	1	MSE	MET	modified residue	UNP P61459
B	50	MSE	MET	modified residue	UNP P61459
B	103	MSE	MET	modified residue	UNP P61459
C	1	MSE	MET	modified residue	UNP P61459
C	50	MSE	MET	modified residue	UNP P61459
C	103	MSE	MET	modified residue	UNP P61459
D	1	MSE	MET	modified residue	UNP P61459
D	50	MSE	MET	modified residue	UNP P61459
D	103	MSE	MET	modified residue	UNP P61459

- Molecule 2 is a protein called HEPATOCYTE NUCLEAR FACTOR 1-ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	31	Total	C	N	O	S	0	0	0
			226	144	36	45	1			
2	F	27	Total	C	N	O		0	0	0
			196	125	31	40				

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	G	27	Total	C	N	O	0	0	0
			197	126	31	40			
2	H	26	Total	C	N	O	0	0	0
			192	123	30	39			



- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	27	Total	O	0	0
			27	27		
3	B	19	Total	O	0	0
			19	19		
3	C	10	Total	O	0	0
			10	10		
3	D	2	Total	O	0	0
			2	2		
3	E	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DIMERIZATION COFACTOR OF HEPATOCYTE NUCLEAR FACTOR 1-ALPHA




- Molecule 1: DIMERIZATION COFACTOR OF HEPATOCYTE NUCLEAR FACTOR 1-ALPHA




- Molecule 1: DIMERIZATION COFACTOR OF HEPATOCYTE NUCLEAR FACTOR 1-ALPHA




- Molecule 1: DIMERIZATION COFACTOR OF HEPATOCYTE NUCLEAR FACTOR 1-ALPHA



- Molecule 2: HEPATOCYTE NUCLEAR FACTOR 1-ALPHA





• Molecule 2: HEPATOCYTE NUCLEAR FACTOR 1-ALPHA



• Molecule 2: HEPATOCYTE NUCLEAR FACTOR 1-ALPHA



• Molecule 2: HEPATOCYTE NUCLEAR FACTOR 1-ALPHA



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.48Å 82.75Å 70.64Å 90.00° 97.83° 90.00°	Depositor
Resolution (Å)	20.10 – 2.60 20.10 – 2.60	Depositor EDS
% Data completeness (in resolution range)	95.1 (20.10-2.60) 94.9 (20.10-2.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.81 (at 2.60Å)	Xtriage
Refinement program	CNS 0.4	Depositor
R, R_{free}	0.257 , 0.299 0.256 , 0.296	Depositor DCC
R_{free} test set	1296 reflections (7.82%)	wwPDB-VP
Wilson B-factor (Å ²)	43.3	Xtriage
Anisotropy	0.079	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 72.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4106	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	3/849 (0.4%)	0.86	2/1142 (0.2%)
1	B	0.56	1/817 (0.1%)	0.90	2/1101 (0.2%)
1	C	0.48	1/824 (0.1%)	0.81	1/1109 (0.1%)
1	D	0.45	1/815 (0.1%)	0.79	1/1100 (0.1%)
2	E	0.32	0/225	0.72	0/301
2	F	0.30	0/195	0.81	0/262
2	G	0.28	0/196	0.76	0/264
2	H	0.27	0/191	0.74	0/257
All	All	0.50	6/4112 (0.1%)	0.82	6/5536 (0.1%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	50	MSE	SE-CE	-10.10	1.65	1.95
1	B	103	MSE	SE-CE	-5.53	1.78	1.95
1	C	50	MSE	SE-CE	-5.36	1.79	1.95
1	A	50	MSE	CG-SE	-5.29	1.79	1.95
1	D	103	MSE	SE-CE	-5.06	1.80	1.95

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	24	GLY	N-CA-C	6.23	123.02	114.92
1	D	48	GLY	N-CA-C	-6.11	104.96	112.77
1	B	24	GLY	N-CA-C	6.01	121.94	114.37
1	C	48	GLY	N-CA-C	-5.97	105.13	112.77
1	A	48	GLY	N-CA-C	-5.96	105.14	112.77

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	831	0	791	20	0
1	B	799	0	759	20	0
1	C	806	0	758	40	0
1	D	797	0	745	49	0
2	E	226	0	251	23	0
2	F	196	0	212	16	0
2	G	197	0	211	12	0
2	H	192	0	209	15	0
3	A	27	0	0	0	0
3	B	19	0	0	0	0
3	C	10	0	0	1	0
3	D	2	0	0	1	0
3	E	4	0	0	0	0
All	All	4106	0	3936	155	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 155 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:TRP:HE1	1:B:68:ASN:HD22	1.08	0.95
1:A:68:ASN:HD22	1:B:66:TRP:HE1	1.15	0.94
1:D:55:LEU:HD13	2:H:19:SER:HB3	1.52	0.89
2:E:23:LYS:HD3	2:F:31:GLY:HA2	1.60	0.82
1:A:50:MSE:HE2	1:A:73:VAL:HG22	1.64	0.79

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	101/104 (97%)	93 (92%)	8 (8%)	0	100	100
1	B	97/104 (93%)	93 (96%)	4 (4%)	0	100	100
1	C	97/104 (93%)	82 (84%)	10 (10%)	5 (5%)	1	2
1	D	98/104 (94%)	79 (81%)	12 (12%)	7 (7%)	1	1
2	E	29/32 (91%)	23 (79%)	5 (17%)	1 (3%)	3	4
2	F	25/32 (78%)	22 (88%)	3 (12%)	0	100	100
2	G	25/32 (78%)	21 (84%)	4 (16%)	0	100	100
2	H	24/32 (75%)	21 (88%)	3 (12%)	0	100	100
All	All	496/544 (91%)	434 (88%)	49 (10%)	13 (3%)	4	7

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	80	HIS
1	D	81	GLU
1	D	82	CYS
1	C	32	ASP
1	D	8	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/85 (100%)	85 (100%)	0	100	100
1	B	82/85 (96%)	81 (99%)	1 (1%)	63	83
1	C	83/85 (98%)	82 (99%)	1 (1%)	63	83
1	D	80/85 (94%)	76 (95%)	4 (5%)	22	46
2	E	25/26 (96%)	25 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	21/26 (81%)	21 (100%)	0	100	100
2	G	21/26 (81%)	21 (100%)	0	100	100
2	H	21/26 (81%)	21 (100%)	0	100	100
All	All	418/444 (94%)	412 (99%)	6 (1%)	59	81

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	81	GLU
1	D	89	ASP
1	D	103	MSE
1	C	39	HIS
1	B	7	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	80	HIS
2	E	28	GLN
2	G	9	GLN
2	E	7	GLN
1	C	63	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	101/104 (97%)	-0.09	3 (2%) 52 47	13, 29, 60, 87	0
1	B	97/104 (93%)	-0.16	1 (1%) 79 76	15, 29, 51, 68	0
1	C	97/104 (93%)	1.08	16 (16%) 4 3	20, 61, 90, 103	0
1	D	98/104 (94%)	2.13	50 (51%) 0 0	60, 90, 103, 117	0
2	E	31/32 (96%)	1.28	5 (16%) 4 3	39, 67, 91, 96	0
2	F	27/32 (84%)	1.15	5 (18%) 3 3	51, 68, 91, 99	0
2	G	27/32 (84%)	1.61	9 (33%) 1 1	75, 90, 102, 106	0
2	H	26/32 (81%)	1.57	9 (34%) 1 1	56, 98, 110, 111	0
All	All	504/544 (92%)	0.88	98 (19%) 3 2	13, 59, 102, 117	0

The worst 5 of 98 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	33	ALA	7.3
1	C	44	ASN	4.5
1	C	70	TYR	4.5
1	C	71	ASN	4.5
1	D	101	VAL	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.