



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:17 AM UTC

PDB ID : 4F9M / pdb_00004f9m
Title : Crystal structure of the PPARgamma-LBD complexed with a cercosporamide derivative modulator
Authors : Matsui, Y.; Hanzawa, H.
Deposited on : 2012-05-19
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	283	
2	C	15	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2374 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisome proliferator-activated receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	261	Total	C	N	O	S	0	0	0
			2096	1355	340	391	10			


There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	195	MET	-	expression tag	UNP P37231
A	196	ARG	-	expression tag	UNP P37231
A	197	GLY	-	expression tag	UNP P37231
A	198	SER	-	expression tag	UNP P37231
A	199	HIS	-	expression tag	UNP P37231
A	200	HIS	-	expression tag	UNP P37231
A	201	HIS	-	expression tag	UNP P37231
A	202	HIS	-	expression tag	UNP P37231
A	203	HIS	-	expression tag	UNP P37231
A	204	HIS	-	expression tag	UNP P37231
A	205	GLY	-	expression tag	UNP P37231

- Molecule 2 is a protein called peptide from Nuclear receptor coactivator 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	11	Total	C	N	O	0	0	0
			94	60	20	14			

- Molecule 3 is (9aS)-8-acetyl-N-[(2-ethyl-4-fluoronaphthalen-1-yl)methyl]-1,7-dihydroxy-3-methoxy-9a-methyl-9-oxo-9,9a-dihydrodibenzo[b,d]furan-4-carboxamide (CCD ID: FCM) (formula: C₃₀H₂₆FNO₇).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			39	30	1	1	7		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	142	Total	O	0	0
			142	142		
4	C	3	Total	O	0	0
			3	3		

- Molecule 1: Peroxisome proliferator-activated receptor gamma



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	38.69Å 54.17Å 66.22Å 90.00° 91.38° 90.00°	Depositor
Resolution (Å)	19.37 – 1.90 19.37 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.4 (19.37-1.90) 99.3 (19.37-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.47 (at 1.85Å)	Xtriage
Refinement program	CNX 2000.1	Depositor
R, R_{free}	0.228 , 0.254 0.225 , 0.252	Depositor DCC
R_{free} test set	2364 reflections (10.14%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.288	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.033 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2374	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FCM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/2131	0.85	9/2868 (0.3%)
2	C	0.44	0/95	0.90	0/125
All	All	0.35	0/2226	0.86	9/2993 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	403	VAL	N-CA-C	6.71	116.84	110.53
1	A	335	ASN	N-CA-C	-6.39	100.11	109.18
1	A	226	PHE	CA-C-N	6.09	125.58	119.24
1	A	226	PHE	C-N-CA	6.09	125.58	119.24
1	A	322	VAL	N-CA-C	5.79	116.44	110.36
1	A	431	LEU	N-CA-C	5.65	117.12	111.07
1	A	425	HIS	CA-C-N	5.62	125.99	119.47
1	A	425	HIS	C-N-CA	5.62	125.99	119.47
1	A	457	LYS	N-CA-C	-5.13	106.29	113.37

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	222	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2096	0	2157	59	0
2	C	94	0	100	9	0
3	A	39	0	25	4	0
4	A	142	0	0	0	0
4	C	3	0	0	0	0
All	All	2374	0	2282	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:696:GLU:HG2	2:C:697:GLY:H	1.33	0.90
1:A:262:ILE:HG13	1:A:263:LYS:HG3	1.61	0.82
1:A:471:GLU:CD	2:C:687:HIS:HA	2.11	0.74
1:A:465:LEU:HD23	1:A:470:GLN:HG2	1.70	0.74
1:A:275:LYS:HB3	1:A:357:ARG:HE	1.54	0.73
1:A:325:ILE:HD12	1:A:388:ILE:HG23	1.69	0.73
1:A:279:ILE:HG23	1:A:463:MET:HE1	1.76	0.67
1:A:264:PHE:HZ	1:A:345:GLN:HE22	1.43	0.66
1:A:443:ARG:O	1:A:447:THR:HG23	1.96	0.66
2:C:696:GLU:HG2	2:C:697:GLY:N	2.11	0.64
1:A:452:LEU:O	1:A:455:VAL:HG12	1.98	0.64
1:A:450:VAL:O	1:A:454:GLN:HG3	1.97	0.64
1:A:319:LYS:HD2	2:C:687:HIS:NE2	2.13	0.63
1:A:244:LYS:HD2	1:A:244:LYS:H	1.64	0.63
1:A:463:MET:HE3	1:A:463:MET:HA	1.81	0.61
1:A:275:LYS:CB	1:A:357:ARG:HE	2.14	0.60
1:A:276:GLU:HG2	1:A:280:ARG:HH21	1.67	0.60
1:A:457:LYS:HE2	1:A:463:MET:O	2.03	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:457:LYS:O	1:A:457:LYS:HD3	2.02	0.58
1:A:319:LYS:HZ3	2:C:687:HIS:CE1	2.23	0.57
1:A:456:ILE:C	1:A:458:LYS:H	2.11	0.57
1:A:468:LEU:HD13	2:C:689:ILE:CG2	2.35	0.56
1:A:446:VAL:O	1:A:450:VAL:HG23	2.06	0.54
1:A:327:TYR:CZ	1:A:367:LYS:HE3	2.42	0.54
1:A:230:LYS:O	1:A:234:ARG:HG3	2.06	0.54
1:A:430:GLN:O	1:A:434:LYS:HG3	2.09	0.53
1:A:259:GLU:OE1	1:A:263:LYS:HD2	2.07	0.53
1:A:424:ASN:C	1:A:426:PRO:HD3	2.34	0.52
1:A:447:THR:O	1:A:451:GLN:HG3	2.09	0.52
1:A:318:LEU:O	1:A:322:VAL:HG22	2.10	0.51
1:A:421:LEU:HD11	1:A:435:LEU:HD12	1.93	0.51
1:A:425:HIS:N	1:A:426:PRO:HD3	2.26	0.51
1:A:335:ASN:OD1	1:A:337:ASP:N	2.44	0.50
1:A:360:PHE:O	1:A:363:PHE:HB3	2.11	0.49
1:A:418:GLU:O	1:A:422:LYS:HD3	2.12	0.49
1:A:363:PHE:HA	1:A:452:LEU:HD23	1.95	0.48
1:A:275:LYS:HG3	1:A:278:ALA:H	1.78	0.48
1:A:468:LEU:HD13	2:C:689:ILE:HG21	1.96	0.48
1:A:459:THR:O	1:A:459:THR:HG22	2.14	0.48
1:A:285:CYS:SG	3:A:501:FCM:H22	2.55	0.47
1:A:335:ASN:OD1	1:A:335:ASN:C	2.57	0.47
1:A:276:GLU:CG	1:A:280:ARG:HH21	2.27	0.47
1:A:324:GLU:HG2	1:A:446:VAL:HG21	1.97	0.47
1:A:263:LYS:HE2	3:A:501:FCM:O14	2.15	0.47
2:C:688:LYS:O	2:C:692:ARG:HG3	2.15	0.47
1:A:379:LEU:HD21	1:A:435:LEU:HD21	1.96	0.46
1:A:365:GLU:HB3	1:A:366:PRO:HD3	1.98	0.46
1:A:468:LEU:HD13	2:C:689:ILE:HG22	1.98	0.46
1:A:322:VAL:HG21	1:A:472:ILE:HD13	1.97	0.45
1:A:325:ILE:HG22	1:A:329:MET:HE2	1.99	0.45
1:A:457:LYS:O	1:A:457:LYS:CD	2.65	0.44
1:A:454:GLN:O	1:A:458:LYS:HB2	2.17	0.44
1:A:241:THR:HG21	1:A:243:ASP:OD2	2.18	0.44
1:A:348:MET:SD	1:A:353:LEU:HD21	2.58	0.44
1:A:367:LYS:HE2	3:A:501:FCM:F38	2.07	0.44
1:A:453:LEU:HD13	1:A:473:TYR:CZ	2.53	0.44
3:A:501:FCM:O18	3:A:501:FCM:H6	2.18	0.43
1:A:456:ILE:C	1:A:458:LYS:N	2.74	0.42
1:A:452:LEU:O	1:A:456:ILE:HG13	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:463:MET:HA	1:A:463:MET:CE	2.49	0.41
1:A:244:LYS:H	1:A:244:LYS:CD	2.32	0.41
1:A:241:THR:HG22	1:A:242:THR:N	2.36	0.41
1:A:245:SER:HB2	1:A:246:PRO:HD2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/283 (91%)	252 (98%)	5 (2%)	0	100	100
2	C	9/15 (60%)	9 (100%)	0	0	100	100
All	All	266/298 (89%)	261 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/254 (92%)	232 (99%)	2 (1%)	70	73
2	C	10/14 (71%)	9 (90%)	1 (10%)	7	3
All	All	244/268 (91%)	241 (99%)	3 (1%)	63	63

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	207	GLU
1	A	237	LEU
2	C	691	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	253	ASN
1	A	286	GLN
1	A	312	ASN
1	A	345	GLN
1	A	412	ASN
1	A	437	GLN
2	C	691	HIS
2	C	695	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FCM	A	501	-	42,43,43	1.53	11 (26%)	63,66,66	2.16	16 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FCM	A	501	-	-	9/17/51/51	0/5/5/5

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	FCM	C1-C2	3.46	1.48	1.41
3	A	501	FCM	C3-C4	3.42	1.48	1.36
3	A	501	FCM	C28-C29	-2.67	1.37	1.41
3	A	501	FCM	C30-C29	2.38	1.48	1.43
3	A	501	FCM	C10-C21	-2.34	1.47	1.51
3	A	501	FCM	C27-C28	2.33	1.40	1.36
3	A	501	FCM	C3-C2	2.25	1.47	1.42
3	A	501	FCM	C31-C29	-2.04	1.38	1.42
3	A	501	FCM	C25-C30	-2.03	1.39	1.43
3	A	501	FCM	O7-C4	-2.01	1.35	1.39
3	A	501	FCM	C33-C34	2.00	1.41	1.36

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	FCM	O7-C4-C3	7.17	135.66	124.70
3	A	501	FCM	C27-C28-C29	-6.06	120.92	124.15
3	A	501	FCM	C9-C5-C6	-5.40	112.03	120.36
3	A	501	FCM	C36-O35-C11	-5.23	109.84	117.51
3	A	501	FCM	O7-C4-C5	-4.05	104.31	110.44
3	A	501	FCM	C8-O7-C4	3.77	110.65	105.89
3	A	501	FCM	C19-C5-C6	3.20	109.73	105.44
3	A	501	FCM	O35-C11-C10	2.96	120.04	115.84
3	A	501	FCM	F38-C28-C29	2.72	122.61	117.25
3	A	501	FCM	O7-C8-C10	2.71	129.12	123.39
3	A	501	FCM	C10-C8-C9	-2.71	120.06	124.78
3	A	501	FCM	C5-C9-C8	-2.56	104.34	107.35
3	A	501	FCM	C5-C4-C3	-2.46	120.01	124.79
3	A	501	FCM	C9-C5-C4	2.42	103.42	99.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	FCM	C5-C9-C13	2.28	136.09	132.24
3	A	501	FCM	O35-C11-C12	-2.18	120.32	124.08

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	501	FCM	C6-C1-C15-C16
3	A	501	FCM	C6-C1-C15-O17
3	A	501	FCM	C8-C10-C21-N23
3	A	501	FCM	C11-C10-C21-O22
3	A	501	FCM	C11-C10-C21-N23
3	A	501	FCM	C2-C1-C15-C16
3	A	501	FCM	C2-C1-C15-O17
3	A	501	FCM	C8-C10-C21-O22
3	A	501	FCM	C12-C11-O35-C36

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	FCM	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	261/283 (92%)	0.53	19 (7%) 21 22	14, 28, 49, 56	0
2	C	11/15 (73%)	1.85	4 (36%) 1 0	22, 35, 47, 49	0
All	All	272/298 (91%)	0.58	23 (8%) 16 17	14, 29, 49, 56	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	477	TYR	6.2
2	C	697	GLY	5.9
1	A	462	ASP	5.7
1	A	238	THR	3.8
1	A	455	VAL	3.4
2	C	687	HIS	3.4
1	A	458	LYS	3.4
1	A	264	PHE	2.9
1	A	457	LYS	2.9
1	A	450	VAL	2.8
1	A	456	ILE	2.8
2	C	696	GLU	2.7
1	A	241	THR	2.6
1	A	463	MET	2.6
1	A	242	THR	2.6
1	A	244	LYS	2.4
1	A	461	THR	2.3
1	A	327	TYR	2.2
1	A	252	MET	2.2
1	A	239	GLY	2.2
2	C	695	GLN	2.1
1	A	276	GLU	2.1
1	A	452	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FCM	A	501	39/39	0.88	0.10	28,32,40,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.