



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 07:00 PM UTC

PDB ID : 6F9P / pdb_00006f9p
Title : Crystal structure of the Fe(II)/alpha-ketoglutarate dependent dioxygenase KDO5 with Re(II)
Authors : Isabet, T.; Stura, E.; Legrand, P.; Zaparucha, A.; Bastard, K.
Deposited on : 2017-12-15
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	2% 86% 7% 7%
1	B	372	2% 67% 10% 22%
1	C	372	4% 79% 10% 10%
1	D	372	0% 84% 6% 9%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10635 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-lysine 4-hydroxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	347	Total	C	N	O	S	0	1	0
			2756	1752	471	521	12			
1	B	289	Total	C	N	O	S	0	2	0
			2334	1498	397	428	11			
1	C	335	Total	C	N	O	S	0	1	0
			2663	1692	457	502	12			
1	D	339	Total	C	N	O	S	0	0	0
			2684	1706	460	506	12			

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		
2	B	1	Total	Cl	0	0
			1	1		
2	C	1	Total	Cl	0	0
			1	1		
2	D	1	Total	Cl	0	0
			1	1		

- Molecule 3 is RHENIUM (CCD ID: RE) (formula: Re).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Re	0	0
			1	1		
3	B	1	Total	Re	0	0
			1	1		
3	C	1	Total	Re	0	0
			1	1		
3	D	1	Total	Re	0	0
			1	1		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	60	Total	O	0	0
			60	60		
6	B	31	Total	O	0	0
			31	31		
6	C	32	Total	O	0	0
			32	32		
6	D	46	Total	O	0	0
			46	46		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: L-lysine 4-hydroxylase



• Molecule 1: L-lysine 4-hydroxylase





• Molecule 1: L-lysine 4-hydroxylase





● Molecule 1: L-lysine 4-hydroxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.20Å 98.87Å 165.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.43 – 2.40 49.43 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.43-2.40) 100.0 (49.43-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.14 (at 2.39Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.194 , 0.213 0.193 , 0.213	Depositor DCC
R_{free} test set	2925 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	59.2	Xtriage
Anisotropy	0.633	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 55.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10635	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL, RE, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.87	1/2818 (0.0%)	1.22	5/3813 (0.1%)
1	B	0.81	2/2386 (0.1%)	1.27	10/3221 (0.3%)
1	C	0.82	2/2724 (0.1%)	1.27	6/3686 (0.2%)
1	D	0.82	2/2745 (0.1%)	1.24	5/3715 (0.1%)
All	All	0.83	7/10673 (0.1%)	1.25	26/14435 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	213	MET	SD-CE	-8.98	1.57	1.79
1	D	155	ILE	CG1-CD1	7.19	1.79	1.51
1	B	213	MET	SD-CE	-5.40	1.66	1.79
1	B	20	GLU	C-N	5.14	1.39	1.33
1	C	134	VAL	CA-C	5.12	1.56	1.52
1	C	213	MET	SD-CE	5.11	1.92	1.79
1	D	134	VAL	CA-C	5.06	1.57	1.52

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	318	ILE	CA-C-N	7.08	132.20	122.07
1	C	318	ILE	C-N-CA	7.08	132.20	122.07
1	D	372	ASP	CA-CB-CG	6.99	119.59	112.60
1	C	145	ARG	CG-CD-NE	6.62	126.57	112.00
1	D	314	ARG	N-CA-CB	6.05	120.75	110.71
1	B	314	ARG	N-CA-C	-5.98	104.98	112.93
1	D	177	THR	N-CA-C	-5.89	100.95	110.32
1	A	177	THR	N-CA-C	-5.86	101.01	110.32
1	A	157	ASP	CA-CB-CG	5.83	118.44	112.60
1	B	267	ASN	CA-CB-CG	5.76	118.36	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	315	SER	N-CA-C	-5.70	105.20	111.82
1	C	177	THR	N-CA-C	-5.60	101.41	110.32
1	D	333	ARG	CA-C-N	5.54	128.72	120.90
1	D	333	ARG	C-N-CA	5.54	128.72	120.90
1	A	325	ASN	CA-CB-CG	5.49	118.09	112.60
1	B	120	LEU	N-CA-C	-5.28	105.69	111.82
1	A	273	THR	N-CA-CB	-5.25	102.85	110.36
1	C	267	ASN	CA-CB-CG	5.25	117.85	112.60
1	B	177	THR	N-CA-C	-5.24	102.00	110.32
1	A	370	ASP	CA-CB-CG	5.23	117.83	112.60
1	B	41	VAL	CA-C-N	5.22	125.74	119.94
1	B	41	VAL	C-N-CA	5.22	125.74	119.94
1	B	315	SER	CA-C-N	5.16	131.40	121.54
1	B	315	SER	C-N-CA	5.16	131.40	121.54
1	C	157	ASP	CA-CB-CG	5.09	117.69	112.60
1	B	317	PHE	CA-CB-CG	5.02	118.82	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2756	0	2703	14	0
1	B	2334	0	2303	23	0
1	C	2663	0	2599	25	0
1	D	2684	0	2625	19	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	5	0	0	0	0
5	D	6	0	8	3	0
6	A	60	0	0	0	0
6	B	31	0	0	0	0
6	C	32	0	0	0	0
6	D	46	0	0	0	0
All	All	10635	0	10238	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:155:ILE:CD1	1:D:155:ILE:CG1	1.79	1.55
1:D:155:ILE:O	1:D:155:ILE:HD12	1.58	1.03
1:C:173:LEU:HB2	1:C:314:ARG:NH1	1.76	1.01
1:B:213:MET:HE2	1:B:221:PHE:HE2	1.29	0.97
1:A:213:MET:HE2	1:A:221:PHE:HE2	1.34	0.89
1:A:273:THR:HG22	1:A:276:ALA:H	1.38	0.86
1:A:213:MET:CE	1:A:221:PHE:HE2	1.97	0.77
1:C:173:LEU:HB2	1:C:314:ARG:HH12	1.47	0.76
1:D:155:ILE:CD1	1:D:155:ILE:O	2.33	0.76
1:B:213:MET:CE	1:B:221:PHE:HE2	1.98	0.75
1:D:112:TRP:H	1:D:155:ILE:HD11	1.51	0.75
1:B:213:MET:HE2	1:B:221:PHE:CE2	2.20	0.72
1:D:111:ASN:HA	1:D:155:ILE:HD13	1.73	0.70
1:D:174:PHE:HA	5:D:403:GOL:C3	2.22	0.69
1:C:226:GLN:HB2	1:C:272:GLN:OE1	1.95	0.67
1:C:260:ASP:H	1:C:264:GLN:HE21	1.43	0.66
1:C:32:GLN:OE1	1:C:35:ARG:NH1	2.29	0.65
1:A:213:MET:HE2	1:A:221:PHE:CE2	2.25	0.65
1:B:194:LEU:HB2	1:B:335:GLN:HG3	1.80	0.64
1:D:112:TRP:N	1:D:155:ILE:HD11	2.13	0.63
1:B:334:ARG:HH12	1:B:336:MET:HE2	1.63	0.63
1:C:314:ARG:HE	1:C:318:ILE:HD11	1.66	0.61
1:C:169:SER:O	1:C:316:ALA:HB1	2.00	0.61
1:D:174:PHE:HA	5:D:403:GOL:H32	1.84	0.60
1:B:174:PHE:N	1:B:315:SER:HG	1.99	0.60
1:A:273:THR:CG2	1:A:276:ALA:H	2.14	0.56
1:A:314:ARG:HH21	1:A:314:ARG:HG2	1.70	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:173:LEU:HB2	1:C:314:ARG:HH11	1.70	0.55
1:D:112:TRP:N	1:D:155:ILE:CD1	2.71	0.54
1:B:63:LEU:O	1:B:67:GLN:HG2	2.08	0.53
1:B:217[B]:MET:HE1	1:B:283:PHE:CE1	2.44	0.52
1:A:109:PRO:HG2	1:A:155:ILE:HD13	1.91	0.52
1:A:213:MET:CE	1:A:221:PHE:CE2	2.87	0.52
1:B:217[B]:MET:HE1	1:B:283:PHE:HE1	1.73	0.52
1:C:109:PRO:HG2	1:C:155:ILE:HD13	1.92	0.52
1:B:213:MET:CE	1:B:221:PHE:CE2	2.87	0.52
1:D:192:LEU:HD23	1:D:337:LEU:HD12	1.93	0.51
1:C:105:LEU:HD22	1:C:195:ARG:CZ	2.41	0.50
1:A:192:LEU:HD23	1:A:337:LEU:HD12	1.94	0.49
1:B:192:LEU:HD23	1:B:337:LEU:HD12	1.95	0.49
1:B:214:ASN:OD1	1:B:217[B]:MET:HG2	2.12	0.49
1:C:171:THR:O	1:C:316:ALA:HB2	2.13	0.49
1:C:217:MET:HE3	1:C:286:GLU:OE2	2.13	0.48
1:B:38:LEU:HD22	1:B:126:ILE:HG12	1.95	0.48
1:C:200:VAL:HG21	1:C:318:ILE:HG12	1.96	0.48
1:D:20:GLU:O	1:D:20:GLU:HG2	2.15	0.47
1:A:324:GLU:HB3	1:A:329:ILE:CD1	2.44	0.47
1:A:192:LEU:HB2	1:A:302:LEU:HD12	1.97	0.46
1:C:201:PRO:HD2	1:C:315:SER:HB2	1.97	0.46
1:D:62:HIS:O	1:D:66:PHE:HD2	1.98	0.46
1:D:174:PHE:HA	5:D:403:GOL:H31	1.98	0.45
1:C:315:SER:O	1:C:316:ALA:HB3	2.16	0.45
1:B:195:ARG:HB2	1:B:335:GLN:HG2	1.97	0.45
1:D:192:LEU:HB2	1:D:302:LEU:HD12	1.99	0.45
1:D:197:GLU:HB3	1:D:331:CYS:SG	2.57	0.44
1:D:195:ARG:HD3	1:D:332:GLU:HB3	1.98	0.44
1:D:229:LYS:HD3	1:D:364:HIS:HB2	1.99	0.44
1:B:334:ARG:NH1	1:B:336:MET:HE2	2.29	0.44
1:D:155:ILE:CD1	1:D:155:ILE:CB	2.81	0.44
1:A:63:LEU:HD13	1:B:74:THR:HG21	1.99	0.44
1:C:53:GLU:HA	1:C:58:ILE:HD11	1.99	0.43
1:B:174:PHE:HD2	1:B:284:TRP:HH2	1.64	0.43
1:B:201:PRO:HG2	1:B:318:ILE:HG23	2.00	0.43
1:C:318:ILE:HG22	1:C:319:ALA:N	2.33	0.43
1:C:260:ASP:H	1:C:264:GLN:NE2	2.14	0.43
1:B:181:PHE:HB2	1:B:364:HIS:HD2	1.84	0.43
1:C:192:LEU:HD23	1:C:337:LEU:HD12	2.00	0.43
1:C:216:VAL:HG12	1:C:217:MET:HE2	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:GLN:NE2	1:C:33:GLN:HG3	2.34	0.43
1:C:314:ARG:HH11	1:C:314:ARG:HG3	1.84	0.43
1:C:192:LEU:HB2	1:C:302:LEU:HD12	2.01	0.42
1:D:35:ARG:HD3	1:D:99:GLU:HG3	2.01	0.42
1:B:281:MET:HE3	1:B:281:MET:HA	2.01	0.42
1:A:181:PHE:HB2	1:A:364:HIS:HD2	1.84	0.42
1:B:192:LEU:HB2	1:B:302:LEU:HD12	2.01	0.41
1:B:88:GLN:O	1:B:208:ARG:HG3	2.21	0.41
1:C:181:PHE:CG	1:C:229:LYS:HA	2.56	0.41
1:B:154:VAL:HG11	1:B:336:MET:HE3	2.02	0.40
1:C:181:PHE:HB2	1:C:364:HIS:HD2	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	344/372 (92%)	338 (98%)	6 (2%)	0	100	100
1	B	281/372 (76%)	275 (98%)	6 (2%)	0	100	100
1	C	330/372 (89%)	321 (97%)	8 (2%)	1 (0%)	36	50
1	D	333/372 (90%)	326 (98%)	7 (2%)	0	100	100
All	All	1288/1488 (87%)	1260 (98%)	27 (2%)	1 (0%)	48	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	318	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/322 (93%)	292 (97%)	8 (3%)	39	62
1	B	256/322 (80%)	249 (97%)	7 (3%)	39	62
1	C	290/322 (90%)	285 (98%)	5 (2%)	53	74
1	D	292/322 (91%)	284 (97%)	8 (3%)	39	62
All	All	1138/1288 (88%)	1110 (98%)	28 (2%)	42	64

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	LEU
1	A	41	VAL
1	A	119	LYS
1	A	159	LYS
1	A	229	LYS
1	A	273	THR
1	A	275	GLU
1	A	372	ASP
1	B	71	GLU
1	B	174	PHE
1	B	195	ARG
1	B	197	GLU
1	B	198	GLU
1	B	203	THR
1	B	275	GLU
1	C	203	THR
1	C	229	LYS
1	C	244	THR
1	C	274	SER
1	C	314	ARG
1	D	20	GLU
1	D	203	THR
1	D	230	ASP
1	D	240	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	275	GLU
1	D	314	ARG
1	D	332	GLU
1	D	372	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	43	ASN
1	A	67	GLN
1	A	79	GLN
1	B	40	ASN
1	B	79	GLN
1	B	184	ASN
1	B	267	ASN
1	C	43	ASN
1	C	79	GLN
1	C	264	GLN
1	C	267	ASN
1	D	40	ASN
1	D	43	ASN
1	D	79	GLN
1	D	267	ASN
1	D	321	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	403	-	4,4,4	0.45	0	6,6,6	0.30	0
4	SO4	D	404	-	4,4,4	0.33	0	6,6,6	0.20	0
5	GOL	D	403	-	5,5,5	0.16	0	5,5,5	0.35	0
4	SO4	A	404	-	4,4,4	0.30	0	6,6,6	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	D	403	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	403	GOL	3	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	347/372 (93%)	-0.17	8 (2%) 61 57	38, 67, 112, 139	1 (0%)
1	B	289/372 (77%)	0.04	9 (3%) 51 47	33, 76, 119, 175	2 (0%)
1	C	335/372 (90%)	0.17	16 (4%) 35 32	40, 87, 166, 189	1 (0%)
1	D	339/372 (91%)	-0.02	4 (1%) 76 73	55, 78, 141, 170	0
All	All	1310/1488 (88%)	0.00	37 (2%) 55 51	33, 76, 137, 189	4 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	318	ILE	4.4
1	B	316	ALA	4.4
1	B	147	GLY	4.2
1	C	165	THR	3.5
1	A	231	ALA	3.4
1	C	266	PHE	3.4
1	C	317	PHE	3.3
1	C	284	TRP	3.2
1	B	317	PHE	3.1
1	C	173	LEU	3.0
1	B	174	PHE	3.0
1	C	171	THR	3.0
1	B	243	PRO	3.0
1	C	319	ALA	2.9
1	D	231	ALA	2.8
1	C	217	MET	2.8
1	A	174	PHE	2.8
1	C	174	PHE	2.8
1	C	160	MET	2.8
1	B	154	VAL	2.7
1	C	318	ILE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	213	MET	2.6
1	C	161	ALA	2.6
1	A	241	SER	2.5
1	D	162	ALA	2.5
1	A	161	ALA	2.4
1	A	168	GLY	2.3
1	C	166	GLY	2.2
1	B	347	HIS	2.2
1	A	242	GLY	2.2
1	D	270	ALA	2.1
1	A	17	LEU	2.1
1	C	371	LEU	2.1
1	A	166	GLY	2.1
1	D	18	THR	2.1
1	B	266	PHE	2.0
1	C	316	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	C	401	1/1	0.62	0.34	116,116,116,116	1
2	CL	B	401	1/1	0.77	0.31	92,92,92,92	1
2	CL	D	401	1/1	0.79	0.32	103,103,103,103	1
4	SO4	D	404	5/5	0.86	0.08	134,135,136,137	0
4	SO4	A	404	5/5	0.87	0.18	156,158,158,159	0
5	GOL	D	403	6/6	0.87	0.13	103,108,108,109	0
4	SO4	A	403	5/5	0.88	0.09	111,113,115,115	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	A	401	1/1	0.88	0.23	92,92,92,92	0
3	RE	D	402	1/1	0.97	0.07	108,108,108,108	1
3	RE	B	402	1/1	0.98	0.11	101,101,101,101	1
3	RE	C	402	1/1	0.99	0.09	119,119,119,119	1
3	RE	A	402	1/1	0.99	0.15	76,76,76,76	1

6.5 Other polymers [i](#)

There are no such residues in this entry.