



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:08 AM UTC

PDB ID : 9F9L / pdb_00009f9l
Title : Crystal structure of MUS81-EME1 bound by compound 16.
Authors : Collie, G.W.
Deposited on : 2024-05-07
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	13299 (2.04-2.00)
Clashscore	190562	1022 (2.02-2.02)
Ramachandran outliers	187476	1014 (2.02-2.02)
Sidechain outliers	187428	1014 (2.02-2.02)
RSRZ outliers	180081	13314 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	4% 57% 6% • 37%
1	C	308	0% 57% • 40%
2	B	326	7% 37% 5% 57%
2	D	326	5% 39% 5% 56%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EOH	A	603	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5417 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Crossover junction endonuclease MUS81.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	0	0
			1509	953	274	278	4			
1	C	185	Total	C	N	O	S	0	0	0
			1466	928	267	267	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	244	GLY	-	expression tag	UNP Q96NY9
A	245	SER	-	expression tag	UNP Q96NY9
C	244	GLY	-	expression tag	UNP Q96NY9
C	245	SER	-	expression tag	UNP Q96NY9

- Molecule 2 is a protein called Crossover junction endonuclease EME1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	139	Total	C	N	O	S	0	0	0
			1048	666	173	201	8			
2	D	145	Total	C	N	O	S	0	0	0
			1081	692	178	203	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	245	GLY	-	expression tag	UNP Q96AY2
D	245	GLY	-	expression tag	UNP Q96AY2

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	2	Total	Mg	0	0
			2	2		

- Molecule 4 is ETHANOL (CCD ID: EOH) (formula: C_2H_6O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			3	2	1		

- Molecule 5 is 2-[2-[4-(cyanomethyl)phenyl]phenyl]-5-oxidanyl-6-oxidanylidene-1H-pyrimidine-4-carboxylic acid (CCD ID: A1IA8) (formula: $C_{19}H_{13}N_3O_4$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			26	19	3	4		
5	C	1	Total	C	N	O	0	0
			26	19	3	4		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	95	Total	O	0	0
			95	95		
6	B	30	Total	O	0	0
			30	30		
6	C	96	Total	O	0	0
			96	96		
6	D	33	Total	O	0	0
			33	33		

- Molecule 2: Crossover junction endonuclease EME1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.57Å 75.14Å 83.01Å 90.00° 90.20° 90.00°	Depositor
Resolution (Å)	55.71 – 2.02 55.71 – 2.02	Depositor EDS
% Data completeness (in resolution range)	99.6 (55.71-2.02) 99.6 (55.71-2.02)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 2.01Å)	Xtriage
Refinement program	BUSTER 2.11.8	Depositor
R, R_{free}	0.225 , 0.262 0.215 , 0.246	Depositor DCC
R_{free} test set	2967 reflections (5.20%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtriage
Anisotropy	0.388	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.016 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5417	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1IA8, EOH, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/1538	1.04	1/2090 (0.0%)
1	C	0.72	0/1494	1.01	2/2025 (0.1%)
2	B	0.64	0/1059	0.96	0/1439
2	D	0.71	0/1094	0.96	0/1488
All	All	0.70	0/5185	1.00	3/7042 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	269	VAL	N-CA-CB	7.86	120.40	111.21
1	C	393	VAL	N-CA-C	5.95	116.12	110.53
1	C	300	VAL	N-CA-CB	5.03	118.09	111.25

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1509	0	1489	9	0
1	C	1466	0	1460	4	0
2	B	1048	0	1055	7	0
2	D	1081	0	1083	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	2	0	0	0	0
3	C	2	0	0	0	0
4	A	3	0	6	2	0
5	A	26	0	0	0	0
5	C	26	0	0	0	0
6	A	95	0	0	0	0
6	B	30	0	0	0	0
6	C	96	0	0	0	0
6	D	33	0	0	0	0
All	All	5417	0	5093	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:313:LEU:HD11	2:D:362:VAL:HG23	1.73	0.69
2:D:315:LEU:HD11	2:D:364:VAL:HG23	1.76	0.66
2:B:263:MET:HE2	2:B:429:TRP:HH2	1.66	0.61
1:A:426:HIS:HD2	4:A:603:EOH:H22	1.67	0.59
1:C:287:LEU:HB2	1:C:405:ILE:HD11	1.84	0.59
2:D:347:VAL:HG21	2:D:415:LEU:HD11	1.86	0.57
1:C:317:PRO:HG3	1:C:323:PRO:HB3	1.90	0.53
1:A:426:HIS:HD2	4:A:603:EOH:C2	2.20	0.53
1:A:431:ARG:HG3	1:A:452:SER:OG	2.10	0.52
2:D:313:LEU:HD11	2:D:362:VAL:CG2	2.40	0.52
1:C:334:ARG:NH2	1:C:405:ILE:HD13	2.23	0.52
2:D:263:MET:HE2	2:D:317:ARG:HE	1.76	0.51
1:C:287:LEU:HD13	1:C:405:ILE:HD11	1.94	0.50
2:D:276:MET:HE3	2:D:278:CYS:SG	2.52	0.50
2:B:260:LEU:HB2	2:B:289:CYS:HA	1.94	0.49
2:B:312:VAL:HG11	2:B:354:THR:HG21	1.94	0.48
1:A:354:PHE:CZ	1:A:358:ARG:HD2	2.49	0.48
2:B:360:SER:HA	2:B:422:GLN:HB2	1.96	0.48
1:A:400:LYS:HG3	1:A:411:TYR:CZ	2.50	0.46
2:B:261:LEU:HD11	2:B:282:ILE:HD12	1.98	0.46
2:B:261:LEU:CD1	2:B:282:ILE:HD12	2.48	0.44
1:A:287:LEU:HB2	1:A:405:ILE:HD11	2.00	0.44
1:A:278:THR:HB	1:A:285:PRO:HG2	2.01	0.43
1:A:400:LYS:HG3	1:A:411:TYR:CE1	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:261:LEU:CD1	2:D:282:ILE:HD12	2.49	0.42
1:A:341:CYS:O	1:A:345:ILE:HG12	2.20	0.41
2:D:322:VAL:O	2:D:325:ILE:HG13	2.20	0.41
2:B:264:GLU:CD	2:B:264:GLU:H	2.27	0.41
2:D:263:MET:CE	2:D:317:ARG:HE	2.34	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	188/308 (61%)	187 (100%)	1 (0%)	0	100	100
1	C	179/308 (58%)	178 (99%)	1 (1%)	0	100	100
2	B	131/326 (40%)	127 (97%)	3 (2%)	1 (1%)	16	8
2	D	137/326 (42%)	135 (98%)	2 (2%)	0	100	100
All	All	635/1268 (50%)	627 (99%)	7 (1%)	1 (0%)	43	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	355	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/260 (62%)	155 (96%)	7 (4%)	26	19
1	C	159/260 (61%)	157 (99%)	2 (1%)	61	63
2	B	113/275 (41%)	107 (95%)	6 (5%)	20	13
2	D	114/275 (42%)	110 (96%)	4 (4%)	32	27
All	All	548/1070 (51%)	529 (96%)	19 (4%)	32	27

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	269	VAL
1	A	288	LEU
1	A	297	THR
1	A	300	VAL
1	A	337	LEU
1	A	348	ARG
1	A	405	ILE
2	B	252	ILE
2	B	264	GLU
2	B	285	GLN
2	B	287	VAL
2	B	325	ILE
2	B	344	GLN
1	C	288	LEU
1	C	300	VAL
2	D	253	ILE
2	D	281	VAL
2	D	319	GLU
2	D	325	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	322	ASN
1	A	330	HIS
1	A	426	HIS
1	A	448	ASN
2	B	424	GLN
1	C	283	HIS
1	C	352	GLN
2	D	285	GLN
2	D	424	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EOH	A	603	-	2,2,2	0.44	0	1,1,1	0.60	0
5	A1IA8	A	604	3	26,28,28	0.42	0	32,39,39	0.39	0
5	A1IA8	C	603	3	26,28,28	0.33	0	32,39,39	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	A1IA8	A	604	3	-	1/14/15/15	0/3/3/3
5	A1IA8	C	603	3	-	3/14/15/15	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	603	A1IA8	O-C-C1-C2
5	C	603	A1IA8	O-C-C1-N1
5	A	604	A1IA8	O-C-C1-N1
5	C	603	A1IA8	N-C4-C5-C10


There are no ring outliers.

1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	603	EOH	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand A1IA8 A 604




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	194/308 (62%)	0.34	11 (5%) 29 28	17, 30, 50, 59	0
1	C	185/308 (60%)	0.15	4 (2%) 62 62	16, 26, 44, 53	0
2	B	139/326 (42%)	1.06	22 (15%) 5 4	25, 42, 64, 75	0
2	D	145/326 (44%)	0.92	17 (11%) 9 8	21, 37, 56, 71	0
All	All	663/1268 (52%)	0.57	54 (8%) 18 17	16, 33, 56, 75	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	325	ILE	5.0
2	D	356	GLY	4.3
2	B	249	LEU	4.3
1	A	448	ASN	4.0
2	B	325	ILE	3.9
2	B	446	ALA	3.8
2	D	328	GLY	3.8
2	D	326	ASP	3.7
2	D	249	LEU	3.6
2	B	427	GLN	3.4
1	A	444	MET	3.3
2	D	306	TRP	3.2
2	B	306	TRP	3.1
2	D	366	GLN	3.0
1	A	279	ARG	3.0
2	B	342	THR	3.0
1	A	318	ARG	3.0
2	B	307	VAL	3.0
2	D	343	LEU	3.0
2	B	315	LEU	2.9
1	A	280	GLY	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	377	LEU	2.8
2	D	248	CYS	2.8
2	B	349	ASP	2.8
2	D	451	LEU	2.8
2	B	248	CYS	2.8
1	C	285	PRO	2.7
1	A	379	LEU	2.7
2	D	342	THR	2.6
2	D	406	VAL	2.6
2	B	365	ASP	2.6
2	B	295	ARG	2.6
2	B	310	PRO	2.5
2	B	445	GLU	2.5
1	C	446	SER	2.4
1	C	346	ASP	2.4
1	C	283	HIS	2.4
1	A	446	SER	2.4
2	B	404	SER	2.4
2	B	343	LEU	2.3
1	A	442	GLY	2.3
2	B	366	GLN	2.3
2	D	252	ILE	2.3
2	B	348	THR	2.2
2	B	355	ALA	2.2
2	D	251	HIS	2.2
2	D	307	VAL	2.2
2	B	251	HIS	2.1
1	A	341	CYS	2.1
2	B	323	SER	2.1
2	D	404	SER	2.1
1	A	286	GLU	2.1
2	B	354	THR	2.1
2	D	264	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EOH	A	603	3/3	0.92	0.14	22,22,22,22	0
5	A1IA8	C	603	26/26	0.94	0.09	24,28,37,39	0
5	A1IA8	A	604	26/26	0.95	0.06	26,27,30,32	0
3	MG	A	601	1/1	0.98	0.03	20,20,20,20	0
3	MG	A	602	1/1	0.99	0.02	25,25,25,25	0
3	MG	C	601	1/1	0.99	0.02	17,17,17,17	0
3	MG	C	602	1/1	0.99	0.04	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A1IA8 C 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A1IA8 A 604:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.