



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:57 PM UTC

PDB ID : 1FAX / pdb_00001fax
Title : COAGULATION FACTOR XA INHIBITOR COMPLEX
Authors : Brandstetter, H.; Engh, R.A.
Deposited on : 1996-08-23
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	254	 69% 21% 7%
2	L	96	 38% 18% 43%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2842 atoms, of which 542 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FACTOR XA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	235	Total	C	H	N	O	S	463	0	0
			2295	1168	440	326	347	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	74	ALA	GLU	conflict	UNP P00742
A	75	ALA	GLN	conflict	UNP P00742

- Molecule 2 is a protein called FACTOR XA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	L	55	Total	C	H	N	O	S	129	0	0
			507	246	96	74	84	7			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	105	ALA	ASN	conflict	UNP P00742

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is (2S)-3-(7-carbamimidoylnaphthalen-2-yl)-2-[4-({(3R)-1-[(1Z)-ethanimidoyl]pyrrolidin-3-yl}oxy)phenyl]propanoic acid (CCD ID: DX9) (formula: C₂₆H₂₈N₄O₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	6	0
			39	26	6	4	3		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: FACTOR XA



• Molecule 2: FACTOR XA



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.93Å 73.17Å 79.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 3.00	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-3.00)	Depositor
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.197 , 0.261	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2842	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, DX9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/1893	1.28	10/2550 (0.4%)
2	L	0.57	0/417	1.37	4/561 (0.7%)
All	All	0.57	0/2310	1.30	14/3111 (0.5%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	144	THR	CA-CB-OG1	-7.40	98.50	109.60
1	A	35	ASN	CA-CB-CG	6.07	118.67	112.60
1	A	185(A)	ASP	CA-CB-CG	5.96	118.56	112.60
1	A	144	THR	CA-CB-CG2	5.95	120.61	110.50
2	L	120	ASN	CA-CB-CG	5.94	118.54	112.60
1	A	243	LYS	CA-C-N	5.64	132.31	121.54
1	A	243	LYS	C-N-CA	5.64	132.31	121.54
1	A	76	GLU	N-CA-CB	-5.51	103.81	110.53
1	A	127	TRP	CA-C-O	5.22	127.97	120.51
2	L	97	ASP	CA-CB-CG	5.20	117.80	112.60
2	L	105	ALA	CA-C-O	5.19	124.23	118.52
2	L	85	THR	CA-CB-OG1	-5.06	102.02	109.60
1	A	127	TRP	CA-C-N	5.03	131.16	121.54
1	A	127	TRP	C-N-CA	5.03	131.16	121.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1855	440	1817	30	0
2	L	411	96	381	5	0
3	A	1	0	0	0	0
4	A	33	6	25	4	0
All	All	2300	542	2223	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:GLU:HG3	1:A:232:THR:HG21	1.70	0.72
1:A:23:LYS:HA	1:A:71:ARG:HH21	1.68	0.58
1:A:183:ALA:HB3	1:A:228:TYR:CE2	2.39	0.58
1:A:177:THR:H	1:A:180:MET:HE3	1.69	0.56
2:L:89:CYS:O	2:L:93:ASN:HA	2.06	0.56
2:L:100:CYS:SG	2:L:107:VAL:HG13	2.46	0.56
1:A:70:ASP:HA	1:A:80:GLU:HG2	1.92	0.52
1:A:218:GLY:O	4:A:265:DX9:H9	2.09	0.52
1:A:113:THR:HB	1:A:115:ARG:HH21	1.75	0.52
1:A:137:ILE:HD13	1:A:159:GLU:HG3	1.93	0.49
1:A:192:GLN:HE21	4:A:265:DX9:H141	1.77	0.49
1:A:183:ALA:HB3	1:A:228:TYR:HE2	1.76	0.48
1:A:97:GLU:O	4:A:265:DX9:N32	2.46	0.48
1:A:142:GLY:HA2	1:A:151:GLN:HG2	1.96	0.47
1:A:213:VAL:HA	1:A:228:TYR:HD1	1.79	0.47
1:A:46:ILE:HA	1:A:52:ILE:HD13	1.97	0.46
2:L:103:GLU:HB2	2:L:108:VAL:HG21	1.97	0.46
2:L:111:CYS:HB3	2:L:115:TYR:HB2	1.98	0.46
1:A:57:HIS:HB3	1:A:94:PHE:CZ	2.51	0.46
1:A:128:ALA:HA	1:A:131(A):LEU:HB2	1.98	0.45
1:A:192:GLN:NE2	4:A:265:DX9:H141	2.32	0.44
1:A:213:VAL:HA	1:A:228:TYR:CD1	2.53	0.44
1:A:128:ALA:O	1:A:131(B):MET:HG2	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:ILE:CD1	1:A:159:GLU:HG3	2.49	0.43
1:A:158:LEU:HD13	1:A:160:VAL:HG13	2.01	0.43
1:A:49:GLU:O	1:A:112:ILE:HG22	2.19	0.42
1:A:67:ARG:HG2	1:A:70:ASP:HB2	2.02	0.42
1:A:237:TRP:O	1:A:240:ARG:HB2	2.20	0.42
1:A:42:CYS:SG	1:A:61:GLN:NE2	2.93	0.41
1:A:25:GLY:O	1:A:28:PRO:HD3	2.21	0.41
1:A:122:CYS:SG	2:L:132:CYS:C	3.05	0.40
1:A:16:ILE:HG21	1:A:158:LEU:HB2	2.04	0.40
1:A:16:ILE:N	1:A:142:GLY:O	2.55	0.40
1:A:104:ALA:HA	1:A:237:TRP:HH2	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	233/254 (92%)	201 (86%)	24 (10%)	8 (3%)	3	17
2	L	53/96 (55%)	41 (77%)	9 (17%)	3 (6%)	1	8
All	All	286/350 (82%)	242 (85%)	33 (12%)	11 (4%)	2	15

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	128	ALA
1	A	244	THR
2	L	91	LEU
1	A	56	ALA
1	A	92	ASN
1	A	147	GLU
1	A	151	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	127	TRP
2	L	104	GLN
2	L	108	VAL
1	A	131	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/214 (92%)	186 (94%)	12 (6%)	17	49
2	L	46/81 (57%)	41 (89%)	5 (11%)	6	26
All	All	244/295 (83%)	227 (93%)	17 (7%)	14	44

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	GLU
1	A	61	GLN
1	A	77	GLU
1	A	144	THR
1	A	145	HIS
1	A	158	LEU
1	A	159	GLU
1	A	170	LEU
1	A	212	ILE
1	A	227	ILE
1	A	229	THR
1	A	235	LEU
2	L	90	SER
2	L	103	GLU
2	L	106	SER
2	L	116	THR
2	L	138	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
2	L	93	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	DX9	A	265	-	35,36,36	1.41	5 (14%)	41,51,51	1.72	8 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DX9	A	265	-	-	13/24/33/33	0/4/4/4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	265	DX9	C4-C2	-4.30	1.39	1.47
4	A	265	DX9	O25-C19	3.45	1.44	1.38
4	A	265	DX9	C31-N32	2.26	1.32	1.27
4	A	265	DX9	O24-C22	-2.16	1.23	1.30
4	A	265	DX9	C9-C8	-2.07	1.37	1.42

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	265	DX9	C16-C15-C22	5.52	120.13	109.67
4	A	265	DX9	C30-C29-N28	-3.81	98.04	102.93
4	A	265	DX9	C19-O25-C26	3.54	125.59	119.18
4	A	265	DX9	C13-C14-C15	3.32	117.30	112.69
4	A	265	DX9	C29-N28-C27	2.32	115.09	112.38
4	A	265	DX9	C14-C15-C16	-2.32	108.88	112.34
4	A	265	DX9	O25-C26-C30	2.22	116.34	108.71
4	A	265	DX9	C17-C16-C21	2.09	120.90	118.30

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	265	DX9	C16-C15-C22-O23
4	A	265	DX9	C16-C15-C22-O24
4	A	265	DX9	C20-C19-O25-C26
4	A	265	DX9	C27-C26-O25-C19
4	A	265	DX9	C18-C19-O25-C26
4	A	265	DX9	N1-C2-C4-C5
4	A	265	DX9	N1-C2-C4-C9
4	A	265	DX9	C11-C13-C14-C15
4	A	265	DX9	C14-C15-C22-O23
4	A	265	DX9	C14-C15-C22-O24
4	A	265	DX9	N2-C2-C4-C5
4	A	265	DX9	N2-C2-C4-C9
4	A	265	DX9	C12-C13-C14-C15

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	265	DX9	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.