



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 11:29 PM UTC

PDB ID : 5FA0 / pdb\_00005fa0  
Title : The structure of the beta-3-deoxy-D-manno-oct-2-ulosonic acid transferase domain from WbbB  
Authors : Mallette, E.; Ovchinnikova, O.G.; Whitfield, C.; Kimber, M.S.  
Deposited on : 2015-12-10  
Resolution : 2.30 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	410	
1	B	410	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6801 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative N-acetyl glucosaminyl transferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	394	3140	2012	519	596	7	6	0	1	0
1	B	388	3098	1991	505	590	7	5	0	5	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MSE	-	initiating methionine	UNP Q6U8B0
A	1	GLY	-	expression tag	UNP Q6U8B0
A	402	LEU	-	expression tag	UNP Q6U8B0
A	403	GLU	-	expression tag	UNP Q6U8B0
A	404	HIS	-	expression tag	UNP Q6U8B0
A	405	HIS	-	expression tag	UNP Q6U8B0
A	406	HIS	-	expression tag	UNP Q6U8B0
A	407	HIS	-	expression tag	UNP Q6U8B0
A	408	HIS	-	expression tag	UNP Q6U8B0
A	409	HIS	-	expression tag	UNP Q6U8B0
B	0	MSE	-	initiating methionine	UNP Q6U8B0
B	1	GLY	-	expression tag	UNP Q6U8B0
B	402	LEU	-	expression tag	UNP Q6U8B0
B	403	GLU	-	expression tag	UNP Q6U8B0
B	404	HIS	-	expression tag	UNP Q6U8B0
B	405	HIS	-	expression tag	UNP Q6U8B0
B	406	HIS	-	expression tag	UNP Q6U8B0
B	407	HIS	-	expression tag	UNP Q6U8B0
B	408	HIS	-	expression tag	UNP Q6U8B0
B	409	HIS	-	expression tag	UNP Q6U8B0

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	306	Total O 306 306	0	0
3	B	256	Total O 256 256	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative N-acetyl glucosaminyl transferase



- Molecule 1: Putative N-acetyl glucosaminyl transferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.93Å 82.93Å 120.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.47 – 2.30 41.47 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (41.47-2.30) 100.0 (41.47-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.32 (at 1.88Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.185 , 0.229 0.187 , 0.230	Depositor DCC
$R_{free}$ test set	2011 reflections (3.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.8	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 51.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.034 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6801	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.49 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.5948e-03.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/3215	0.68	0/4355
1	B	0.28	0/3183	0.68	2/4315 (0.0%)
All	All	0.28	0/6398	0.68	2/8670 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	110	ILE	CA-C-N	5.37	124.83	119.24
1	B	110	ILE	C-N-CA	5.37	124.83	119.24

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3140	0	3025	29	0
1	B	3098	0	3010	29	0
2	A	1	0	0	1	0
3	A	306	0	0	9	1
3	B	256	0	0	10	1
All	All	6801	0	6035	55	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:ASP:OD1	3:A:601:HOH:O	1.93	0.85
1:B:97:ASP:OD2	3:B:501:HOH:O	1.98	0.81
2:A:501:CL:CL	3:A:867:HOH:O	2.38	0.79
1:A:255:TYR:OH	1:A:294:GLU:OE2	2.01	0.76
1:B:368:GLU:OE1	3:B:502:HOH:O	2.10	0.68
1:B:67:GLU:OE2	3:B:503:HOH:O	2.13	0.66
1:A:125:LYS:NZ	3:A:607:HOH:O	2.26	0.65
1:B:182:THR:O	3:B:505:HOH:O	2.14	0.64
1:B:51:ARG:NH1	3:B:506:HOH:O	2.19	0.64
1:A:99:ILE:HG23	1:A:236:ILE:HD12	1.82	0.61
1:B:163:ARG:NH2	3:B:515:HOH:O	2.30	0.61
1:B:370:VAL:HG21	1:B:392:LEU:HD21	1.82	0.61
1:B:58:PRO:HA	1:B:61:MSE:HE2	1.83	0.58
1:A:96:HIS:ND1	3:A:603:HOH:O	2.17	0.57
1:B:202:GLY:O	1:B:206:ASN:ND2	2.37	0.56
1:A:148:ARG:NH2	3:A:618:HOH:O	2.38	0.56
1:A:382:PRO:HB3	1:A:387:GLN:HG2	1.88	0.56
1:A:210:GLU:OE1	3:A:605:HOH:O	2.18	0.55
1:A:202:GLY:O	1:A:206:ASN:ND2	2.32	0.53
1:A:271:ARG:HH11	1:B:271:ARG:NE	2.07	0.52
1:A:12:ARG:NH2	1:A:231:ASP:OD1	2.42	0.52
1:A:378:SER:OG	3:A:602:HOH:O	2.09	0.52
1:A:122:ASP:OD1	1:A:125:LYS:NZ	2.41	0.51
1:B:343:SER:OG	1:B:366:GLU:OE1	2.29	0.49
1:A:381:MSE:SE	1:A:382:PRO:HD2	2.63	0.49
1:A:407:HIS:N	3:A:622:HOH:O	2.45	0.48
1:B:263:ARG:NE	3:B:504:HOH:O	2.14	0.48
1:B:158:GLU:HB3	1:B:178:VAL:HG21	1.95	0.48
1:B:112:ASP:O	3:B:508:HOH:O	2.20	0.47
1:B:155:MSE:HE3	1:B:361:LEU:HD13	1.95	0.47
1:B:182:THR:HG21	1:B:320:SER:HB3	1.96	0.46
1:B:159:ILE:CG2	1:B:163:ARG:HD2	2.46	0.46
1:B:199:ALA:HB1	1:B:203:ASP:HB2	1.97	0.45
1:A:271:ARG:NH1	1:B:271:ARG:HH21	2.15	0.45
1:A:343:SER:OG	1:A:366:GLU:OE1	2.35	0.45
1:B:354[B]:GLN:HG2	1:B:358:LYS:HE2	1.98	0.45
1:A:124:LYS:O	1:A:128:ILE:HG12	2.16	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:TYR:CD1	1:A:137:ILE:HD13	2.53	0.44
1:B:12:ARG:NH2	1:B:231:ASP:OD1	2.51	0.44
1:A:282:SER:HB3	1:A:287:ALA:HB3	2.00	0.44
1:A:225:VAL:HG22	1:A:251:VAL:HG21	1.99	0.43
1:A:271:ARG:HH11	1:B:271:ARG:CZ	2.32	0.43
1:B:124:LYS:O	1:B:128:ILE:HG12	2.18	0.43
1:A:8:PRO:HB3	1:A:38:GLY:HA3	2.01	0.43
1:B:161:PRO:HD3	1:B:172:TYR:CZ	2.54	0.43
1:B:235:LEU:O	3:B:509:HOH:O	2.22	0.42
1:A:158:GLU:HB3	1:A:178:VAL:HG21	2.02	0.42
1:A:271:ARG:NH1	1:B:271:ARG:HE	2.17	0.42
1:A:209:LEU:HD21	1:A:286:LEU:HD11	2.01	0.42
1:B:33:ILE:O	1:B:77:HIS:HB3	2.20	0.41
1:A:254:ARG:NH1	3:A:631:HOH:O	2.53	0.41
1:B:226:VAL:O	3:B:510:HOH:O	2.22	0.41
1:A:195:PHE:CE1	1:A:392:LEU:HD21	2.56	0.41
1:A:53:ARG:HB3	1:A:55:GLU:OE1	2.21	0.41
1:B:159:ILE:HG23	1:B:163:ARG:HD2	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:855:HOH:O	3:B:749:HOH:O[1_655]	1.89	0.31

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	389/410 (95%)	375 (96%)	11 (3%)	3 (1%)	16	20
1	B	387/410 (94%)	376 (97%)	11 (3%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	776/820 (95%)	751 (97%)	22 (3%)	3 (0%)	30 38

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	381	MSE
1	A	274	ASP
1	A	211	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	338/342 (99%)	333 (98%)	5 (2%)	57 75
1	B	336/342 (98%)	334 (99%)	2 (1%)	78 89
All	All	674/684 (98%)	667 (99%)	7 (1%)	68 82

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LEU
1	A	210	GLU
1	A	211	VAL
1	A	212	LEU
1	A	387	GLN
1	B	398	GLU
1	B	400	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	GLN
1	A	363	HIS
1	A	390	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	406	HIS
1	B	145	HIS
1	B	240	HIS
1	B	363	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	388/410 (94%)	-0.02	13 (3%) 48 50	9, 28, 58, 99	1 (0%)
1	B	383/410 (93%)	0.22	20 (5%) 33 34	10, 31, 69, 84	5 (1%)
All	All	771/820 (94%)	0.10	33 (4%) 40 41	9, 29, 65, 99	6 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	379	ALA	5.7
1	A	212	LEU	4.9
1	A	382	PRO	4.1
1	A	407	HIS	4.0
1	B	402	LEU	3.9
1	A	406	HIS	3.8
1	A	328	ALA	3.4
1	B	330	ALA	3.1
1	A	222	ALA	3.1
1	B	193	ALA	3.1
1	B	390	HIS	2.9
1	B	258	GLU	2.9
1	B	362	GLY	2.8
1	B	281	ASP	2.8
1	B	222	ALA	2.7
1	A	258	GLU	2.6
1	B	329	GLY	2.5
1	A	223	ALA	2.4
1	B	328	ALA	2.4
1	B	212	LEU	2.3
1	A	389	ALA	2.3
1	B	327	LEU	2.3
1	B	283	ALA	2.2
1	A	275	ASP	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	380	ASP	2.1
1	B	401	SER	2.1
1	B	216	GLU	2.1
1	B	275	ASP	2.1
1	B	198	LYS	2.1
1	A	290	ASN	2.1
1	B	363	HIS	2.0
1	B	365	GLU	2.0
1	A	387	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CL	A	501	1/1	0.97	0.15	42,42,42,42	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.