



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:57 AM UTC

PDB ID : 7FA0 / pdb_00007fa0
Title : The crystal structure of VyPAL2-C214A, a dead mutant of VyPAL2 peptide asparaginyl ligase in form II
Authors : Hu, S.; Sahili, A.; Lescar, J.
Deposited on : 2021-07-05
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	 3% 93% 5% ..
1	B	282	 4% 92% 5% .
2	C	2	 100%
3	D	2	 50% 50%

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4889 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptide Asparaginyl Ligases.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	279	2185	1394	355	425	11	0	4	0
1	B	274	2125	1356	344	414	11	0	1	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ASP	deletion	UNP A0A509GV09
A	172	HD0	HIS	conflict	UNP A0A509GV09
A	214	ALA	CYS	engineered mutation	UNP A0A509GV09
B	?	-	ASP	deletion	UNP A0A509GV09
B	172	HD0	HIS	conflict	UNP A0A509GV09
B	214	ALA	CYS	engineered mutation	UNP A0A509GV09

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	2	24	14	1	9	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	2	28	16	2	10	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0
4	B	1	14	8	1	5	0	0
4	B	1	14	8	1	5	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	257	Total 257 O 257	0	0
5	B	214	Total 214 O 214	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Peptide Asparaginyl Ligases



- Molecule 1: Peptide Asparaginyl Ligases



- Molecule 2: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	76.50Å 78.55Å 92.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.33 – 1.80 46.33 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.1 (46.33-1.80) 99.1 (46.33-1.80)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 1.79Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.164 , 0.194 0.164 , 0.190	Depositor DCC
R_{free} test set	2618 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	27.3	Xtrriage
Anisotropy	0.057	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.010 for k,h,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4889	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HD0, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/2234	0.95	4/3035 (0.1%)
1	B	0.72	0/2164	0.98	3/2941 (0.1%)
All	All	0.72	0/4398	0.96	7/5976 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	102	ASN	CA-CB-CG	-8.59	104.02	112.60
1	B	73	ASP	CA-CB-CG	8.24	120.84	112.60
1	A	73	ASP	CA-CB-CG	7.44	120.04	112.60
1	A	67	ASN	CA-C-N	5.29	128.75	120.82
1	A	67	ASN	C-N-CA	5.29	128.75	120.82
1	A	277	ASN	N-CA-C	-5.21	100.60	108.67
1	B	158	ASP	CA-CB-CG	5.03	117.63	112.60

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	172	HD0	Mainchain,Peptide
1	B	172	HD0	Mainchain,Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2185	0	2090	6	0
1	B	2125	0	2026	4	0
2	C	24	0	22	0	0
3	D	28	0	25	0	0
4	A	28	0	26	0	0
4	B	28	0	26	0	0
5	A	257	0	0	0	0
5	B	214	0	0	0	0
All	All	4889	0	4215	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:VAL:HG11	1:A:150:ILE:HD12	1.86	0.56
1:B:58:LEU:HB3	1:B:74:VAL:HG13	1.95	0.48
1:B:282:THR:HA	1:B:312:SER:HA	1.97	0.46
1:A:282:THR:HA	1:A:312:SER:HA	1.98	0.46
1:B:164:HIS:HD2	1:B:206:SER:OG	2.00	0.44
1:A:143:LEU:HD11	1:A:197:GLN:HB2	2.02	0.41
1:A:107:PHE:HB3	1:A:110:ILE:HD12	2.02	0.41
1:A:143:LEU:HD12	1:A:143:LEU:HA	1.91	0.41
1:A:92:ILE:HG13	1:A:155:LYS:HD2	2.03	0.41
1:B:305:ASP:OD1	1:B:307:PRO:HD2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	280/282 (99%)	276 (99%)	4 (1%)	0	100	100
1	B	272/282 (96%)	267 (98%)	5 (2%)	0	100	100
All	All	552/564 (98%)	543 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/233 (100%)	231 (99%)	3 (1%)	61	54
1	B	226/233 (97%)	223 (99%)	3 (1%)	61	54
All	All	460/466 (99%)	454 (99%)	6 (1%)	61	54

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	ILE
1	A	143	LEU
1	A	205	LYS
1	B	51	ILE
1	B	150	ILE
1	B	250	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	66	HIS
1	A	162	ASN
1	A	164	HIS
1	A	215	ASN
1	A	302	GLN
1	B	66	HIS
1	B	162	ASN
1	B	164	HIS
1	B	215	ASN
1	B	276	HIS
1	B	302	GLN
1	B	325	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	HD0	A	172	1	16,18,19	2.05	3 (18%)	18,25,27	2.32	8 (44%)
1	HD0	B	172	1	16,18,19	2.16	4 (25%)	18,25,27	2.13	6 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HD0	A	172	1	-	1/9/26/28	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HD0	B	172	1	-	0/9/26/28	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	172	HD0	C1-NAS	-6.52	1.38	1.47
1	A	172	HD0	C1-NAS	-6.43	1.38	1.47
1	B	172	HD0	CAX-NAS	-3.53	1.33	1.39
1	B	172	HD0	CAU-NAS	-3.16	1.33	1.39
1	A	172	HD0	CAU-NAS	-3.15	1.33	1.39
1	A	172	HD0	CAX-NAS	-3.08	1.33	1.39
1	B	172	HD0	CB-C1	2.42	1.55	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	172	HD0	CAW-CAX-NAS	4.15	110.90	107.59
1	A	172	HD0	OAY-CAX-CAW	-4.02	121.40	127.28
1	B	172	HD0	CA-CAU-NAS	3.75	111.36	107.45
1	A	172	HD0	CA-CAU-NAS	3.70	111.31	107.45
1	B	172	HD0	OAY-CAX-CAW	-3.43	122.26	127.28
1	A	172	HD0	CAW-CAX-NAS	3.39	110.30	107.59
1	B	172	HD0	CD2-CG-ND1	3.01	109.21	105.65
1	A	172	HD0	OAY-CAX-NAS	3.00	128.30	124.36
1	A	172	HD0	OAV-CAU-CA	-2.97	121.96	126.65
1	A	172	HD0	CD2-CG-ND1	2.88	109.05	105.65
1	B	172	HD0	OAV-CAU-CA	-2.81	122.21	126.65
1	A	172	HD0	C1-NAS-CAX	2.74	128.06	123.65
1	A	172	HD0	CAW-CA-CAU	-2.49	101.49	104.45
1	B	172	HD0	CAW-CA-CAU	-2.35	101.65	104.45

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	172	HD0	CB-C1-NAS-CAU

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates i

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	1,2	14,14,15	0.33	0	17,19,21	0.63	0
2	FUC	C	2	2	10,10,11	0.26	0	14,14,16	0.57	0
3	NAG	D	1	1,3	14,14,15	0.29	0	17,19,21	1.12	2 (11%)
3	NAG	D	2	3	14,14,15	0.24	0	17,19,21	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	FUC	C	2	2	-	-	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	NAG	O4-C4-C5	-2.30	103.66	109.32
3	D	1	NAG	O4-C4-C3	-2.29	104.98	110.38

There are no chirality outliers.


All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	402	1	14,14,15	0.26	0	17,19,21	0.36	0
4	NAG	A	401	1	14,14,15	0.32	0	17,19,21	0.55	0
4	NAG	B	402	1	14,14,15	0.27	0	17,19,21	0.73	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	B	401	1	14,14,15	0.30	0	17,19,21	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	402	1	-	0/6/23/26	0/1/1/1
4	NAG	A	401	1	-	3/6/23/26	0/1/1/1
4	NAG	B	402	1	-	1/6/23/26	0/1/1/1
4	NAG	B	401	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	402	NAG	O5-C1-C2	-2.11	108.02	111.29

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	401	NAG	O5-C5-C6-O6
4	A	401	NAG	C4-C5-C6-O6
4	B	402	NAG	O5-C5-C6-O6
4	A	401	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	278/282 (98%)	-0.24	9 (3%) 50 50	17, 28, 48, 69	4 (1%)
1	B	273/282 (96%)	-0.14	10 (3%) 45 45	20, 31, 49, 75	1 (0%)
All	All	551/564 (97%)	-0.19	19 (3%) 48 48	17, 29, 49, 75	5 (0%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	249	GLY	3.1
1	B	143	LEU	3.1
1	A	249	GLY	3.0
1	A	251	PRO	2.9
1	A	250	THR	2.8
1	B	151	GLY	2.7
1	B	252	ASP	2.6
1	B	250	THR	2.6
1	B	251	PRO	2.6
1	B	319	THR	2.6
1	A	253	PHE	2.5
1	A	242	SER	2.4
1	B	322	ALA	2.3
1	A	254	PRO	2.3
1	B	150	ILE	2.3
1	B	51	ILE	2.2
1	A	329	VAL	2.2
1	A	50	SER	2.2
1	A	51	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	HD0	B	172	17/18	0.93	0.12	21,27,45,45	0
1	HD0	A	172	17/18	0.96	0.08	18,21,38,38	0


6.3 Carbohydrates [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	D	2	14/15	0.71	0.14	50,52,53,54	0
2	FUC	C	2	10/11	0.76	0.13	52,53,53,53	0
3	NAG	D	1	14/15	0.88	0.10	42,45,46,48	0
2	NAG	C	1	14/15	0.89	0.09	47,49,50,51	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	B	402	14/15	0.44	0.17	65,67,68,68	0
4	NAG	A	402	14/15	0.78	0.12	48,50,52,53	0
4	NAG	A	401	14/15	0.80	0.14	46,47,48,48	0
4	NAG	B	401	14/15	0.83	0.10	46,49,51,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.