



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:17 AM UTC

PDB ID : 8FA5 / pdb_00008fa5
Title : Crystal structure of Xanthomonas campestris GH35 beta-galactosidase
Authors : Godoy, A.S.; Polikarpov, I.
Deposited on : 2022-11-25
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	591	7% 91% 7% ..
1	B	591	% 94% 5% .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10606 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-galactosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	585	Total	C	N	O	S	0	5	0
			4628	2941	824	850	13			
1	A	585	Total	C	N	O	S	0	0	0
			4582	2913	813	843	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	initiating methionine	UNP Q8P6S0
A	1	MET	-	initiating methionine	UNP Q8P6S0

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	864	Total O 864 864	0	0
3	A	472	Total O 472 472	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-galactosidase



- Molecule 1: Beta-galactosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.86Å 84.77Å 99.81Å 90.00° 96.61° 90.00°	Depositor
Resolution (Å)	42.79 – 1.70 42.79 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.6 (42.79-1.70) 97.6 (42.79-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 1.70Å)	Xtriage
Refinement program	PHENIX v1	Depositor
R, R_{free}	0.170 , 0.203 (Not available) , 0.178	Depositor DCC
R_{free} test set	7177 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	20.6	Xtriage
Anisotropy	0.498	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 51.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10606	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	3/4715 (0.1%)	0.80	4/6439 (0.1%)
1	B	0.90	7/4770 (0.1%)	0.93	11/6509 (0.2%)
All	All	0.80	10/9485 (0.1%)	0.87	15/12948 (0.1%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	440	GLY	C-N	-7.39	1.22	1.33
1	B	76	ALA	C-O	-7.30	1.17	1.24
1	B	161	VAL	C-O	-7.00	1.16	1.24
1	B	162	GLU	C-O	-6.21	1.16	1.24
1	B	163	ASN	C-O	-6.19	1.16	1.23
1	B	277	LEU	C-O	-5.65	1.17	1.24
1	B	222	SER	C-O	-5.64	1.17	1.24
1	B	222	SER	N-CA	5.51	1.52	1.46
1	A	442	HIS	C-O	-5.14	1.17	1.23
1	A	294	PHE	C-O	-5.14	1.18	1.23

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	221[A]	LYS	CA-C-N	11.54	135.44	120.44
1	B	221[A]	LYS	C-N-CA	11.54	135.44	120.44
1	B	221[B]	LYS	CA-C-N	11.54	135.44	120.44
1	B	221[B]	LYS	C-N-CA	11.54	135.44	120.44
1	B	579[A]	ARG	CA-C-N	9.77	140.56	121.41
1	B	579[A]	ARG	C-N-CA	9.77	140.56	121.41
1	B	579[B]	ARG	CA-C-N	9.77	140.56	121.41
1	B	579[B]	ARG	C-N-CA	9.77	140.56	121.41
1	A	441	GLN	O-C-N	-8.89	112.84	123.16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	578	VAL	O-C-N	-5.93	116.32	123.02
1	A	104	ALA	N-CA-C	5.90	120.05	112.86
1	B	490	SER	CA-C-N	-5.46	114.19	119.87
1	B	490	SER	C-N-CA	-5.46	114.19	119.87
1	A	206	LEU	CA-C-N	-5.01	114.60	119.76
1	A	206	LEU	C-N-CA	-5.01	114.60	119.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4582	0	4382	25	0
1	B	4628	0	4444	16	0
2	A	16	0	24	0	0
2	B	44	0	66	7	0
3	A	472	0	0	4	0
3	B	864	0	0	3	0
All	All	10606	0	8916	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:441:GLN:OE1	3:A:701:HOH:O	1.78	0.99
1:A:524:MET:CE	1:A:551:LEU:HD12	2.06	0.85
1:B:489:ARG:HH11	2:B:609:EDO:H22	1.45	0.81
1:A:524:MET:HE3	1:A:551:LEU:HD12	1.66	0.77
1:B:396:THR:H	2:B:610:EDO:H22	1.56	0.69
1:A:524:MET:HE2	1:A:551:LEU:HD12	1.75	0.69
1:B:201:LEU:H	2:B:604:EDO:H21	1.56	0.68
1:B:16[A]:GLN:NE2	3:B:701:HOH:O	2.30	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:434:GLU:OE2	3:A:702:HOH:O	2.16	0.62
1:A:524:MET:HE1	1:A:532:ALA:CB	2.32	0.59
1:B:49:ARG:HG3	2:B:611:EDO:H22	1.85	0.58
1:A:88:ALA:HB2	1:A:155:PRO:HG3	1.88	0.56
1:A:85:ARG:HD3	3:A:948:HOH:O	2.06	0.55
1:A:411:VAL:HG22	1:A:425:GLU:HG2	1.91	0.53
1:A:554:PRO:HB2	1:A:556:PRO:HD2	1.91	0.53
1:A:99:GLY:HA3	1:A:100:PRO:C	2.34	0.53
1:A:282:GLY:HA3	1:A:309:TYR:O	2.10	0.51
1:B:457:GLY:O	1:B:460:MET:HG2	2.12	0.49
1:B:233:ASP:OD1	3:B:702:HOH:O	2.20	0.49
1:A:281:ILE:HB	1:A:315:LEU:HB2	1.96	0.48
1:B:13:GLN:O	1:B:16[A]:GLN:HG2	2.15	0.47
1:A:479:THR:O	1:A:480:ASN:HB3	2.15	0.47
1:B:525:ARG:HD3	1:B:585:VAL:HG11	1.97	0.46
1:A:231:GLN:HG2	1:A:234:GLN:HB2	1.96	0.46
1:A:229:LYS:HE3	1:A:229:LYS:HB2	1.70	0.45
1:B:201:LEU:HD12	2:B:604:EDO:H12	1.98	0.45
1:A:294:PHE:CG	1:A:459:ARG:HD2	2.52	0.45
1:B:99:GLY:HA3	1:B:100:PRO:C	2.41	0.44
1:A:237:MET:N	3:A:717:HOH:O	2.47	0.44
1:A:398:PRO:HA	1:A:437:ILE:O	2.17	0.44
1:A:51:LEU:HD23	1:A:53:LEU:HD13	1.99	0.44
1:B:221[B]:LYS:HD2	3:B:781:HOH:O	2.17	0.43
1:A:40:TYR:O	1:A:44:ARG:HG2	2.19	0.43
1:B:524:MET:HE2	1:B:542:HIS:CD2	2.54	0.43
1:A:278:TYR:HA	1:A:279:MET:HA	1.73	0.43
1:A:390:LEU:HB3	1:A:484:PHE:HB2	2.02	0.42
1:B:201:LEU:N	2:B:604:EDO:H21	2.29	0.42
1:A:354:ASP:HA	1:A:579:ARG:HB3	2.01	0.42
1:B:11:ALA:HB1	2:B:606:EDO:H22	2.02	0.42
1:B:160:GLN:HB2	1:B:193:PHE:CZ	2.56	0.41
1:A:101:TYR:HB2	1:A:111:TYR:OH	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	583/591 (99%)	564 (97%)	19 (3%)	0	100	100
1	B	588/591 (100%)	570 (97%)	18 (3%)	0	100	100
All	All	1171/1182 (99%)	1134 (97%)	37 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	466/478 (98%)	463 (99%)	3 (1%)	78	72
1	B	473/478 (99%)	472 (100%)	1 (0%)	87	84
All	All	939/956 (98%)	935 (100%)	4 (0%)	84	80

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	77	ASN
1	A	53	LEU
1	A	77	ASN
1	A	212	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	295	GLN
1	B	511	ASN
1	A	6	GLN
1	A	234	GLN
1	A	339	GLN
1	A	418	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	601	-	3,3,3	0.43	0	2,2,2	0.49	0
2	EDO	B	610	-	3,3,3	0.43	0	2,2,2	0.42	0
2	EDO	A	602	-	3,3,3	0.38	0	2,2,2	0.60	0
2	EDO	B	603	-	3,3,3	0.39	0	2,2,2	0.60	0
2	EDO	B	605	-	3,3,3	0.57	0	2,2,2	0.26	0
2	EDO	B	601	-	3,3,3	0.69	0	2,2,2	0.41	0
2	EDO	B	607	-	3,3,3	0.40	0	2,2,2	0.43	0
2	EDO	B	608	-	3,3,3	0.33	0	2,2,2	0.63	0
2	EDO	B	602	-	3,3,3	0.41	0	2,2,2	0.68	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	A	603	-	3,3,3	0.49	0	2,2,2	0.37	0
2	EDO	A	604	-	3,3,3	0.39	0	2,2,2	0.23	0
2	EDO	B	604	-	3,3,3	0.35	0	2,2,2	0.41	0
2	EDO	B	611	-	3,3,3	0.36	0	2,2,2	0.72	0
2	EDO	B	609	-	3,3,3	0.51	0	2,2,2	0.15	0
2	EDO	B	606	-	3,3,3	0.38	0	2,2,2	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	601	-	-	1/1/1/1	-
2	EDO	B	610	-	-	1/1/1/1	-
2	EDO	A	602	-	-	0/1/1/1	-
2	EDO	B	603	-	-	0/1/1/1	-
2	EDO	B	605	-	-	0/1/1/1	-
2	EDO	B	601	-	-	0/1/1/1	-
2	EDO	B	607	-	-	0/1/1/1	-
2	EDO	B	608	-	-	0/1/1/1	-
2	EDO	B	602	-	-	0/1/1/1	-
2	EDO	A	603	-	-	0/1/1/1	-
2	EDO	A	604	-	-	1/1/1/1	-
2	EDO	B	604	-	-	1/1/1/1	-
2	EDO	B	611	-	-	0/1/1/1	-
2	EDO	B	609	-	-	0/1/1/1	-
2	EDO	B	606	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	604	EDO	O1-C1-C2-O2
2	B	610	EDO	O1-C1-C2-O2
2	A	604	EDO	O1-C1-C2-O2
2	A	601	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	610	EDO	1	0
2	B	604	EDO	3	0
2	B	611	EDO	1	0
2	B	609	EDO	1	0
2	B	606	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	221[A]:LYS	C	222:SER	N	1.17

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	585/591 (98%)	0.66	41 (7%)	22 24	20, 46, 70, 86	0
1	B	585/591 (98%)	-0.39	4 (0%)	84 86	11, 21, 44, 73	5 (0%)
All	All	1170/1182 (98%)	0.13	45 (3%)	44 48	11, 32, 66, 86	5 (0%)

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	156	ILE	3.2
1	A	7	TRP	3.0
1	A	159	VAL	2.9
1	A	18	VAL	2.8
1	A	342	ALA	2.8
1	A	338	VAL	2.8
1	A	158	ALA	2.8
1	A	335	VAL	2.8
1	A	269	LEU	2.7
1	A	208	GLY	2.6
1	A	243	ALA	2.6
1	A	331	VAL	2.5
1	A	24	TYR	2.5
1	A	95	ILE	2.5
1	A	348	ALA	2.5
1	A	233	ASP	2.5
1	A	583	GLN	2.4
1	A	26	VAL	2.4
1	A	347	ILE	2.4
1	A	345	ALA	2.4
1	A	80	VAL	2.4
1	A	350	ALA	2.4
1	A	157	ILE	2.4
1	A	151	HIS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	89	ALA	2.3
1	A	332	ILE	2.3
1	A	153	GLY	2.3
1	B	590	LYS	2.3
1	A	154	GLY	2.2
1	A	9	THR	2.2
1	A	149	LEU	2.2
1	A	14	GLY	2.2
1	A	515	GLY	2.2
1	A	155	PRO	2.2
1	A	53	LEU	2.1
1	B	6	GLN	2.1
1	A	325	PHE	2.1
1	A	21	GLY	2.1
1	A	191	LEU	2.1
1	A	343	LEU	2.1
1	A	501	THR	2.1
1	A	70	GLY	2.0
1	B	501	THR	2.0
1	A	262	THR	2.0
1	B	474	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	B	610	4/4	0.71	0.18	50,53,55,57	0
2	EDO	B	603	4/4	0.78	0.17	38,39,40,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	B	611	4/4	0.79	0.17	44,44,48,54	0
2	EDO	B	609	4/4	0.80	0.15	53,55,56,58	0
2	EDO	B	608	4/4	0.84	0.17	51,52,56,61	0
2	EDO	B	607	4/4	0.87	0.14	39,40,41,42	0
2	EDO	A	603	4/4	0.90	0.10	37,39,39,40	0
2	EDO	B	604	4/4	0.91	0.12	32,37,38,39	0
2	EDO	A	601	4/4	0.91	0.12	28,32,36,39	0
2	EDO	B	606	4/4	0.91	0.10	45,46,48,53	0
2	EDO	A	604	4/4	0.94	0.10	32,35,38,38	0
2	EDO	A	602	4/4	0.95	0.07	29,32,33,35	0
2	EDO	B	601	4/4	0.97	0.05	17,18,20,21	0
2	EDO	B	602	4/4	0.98	0.04	17,17,19,20	0
2	EDO	B	605	4/4	0.98	0.06	21,23,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.