



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 06:10 AM UTC

PDB ID : 1FBH / pdb_00001fbh
Title : CRYSTALLOGRAPHIC STUDIES OF THE CATALYTIC MECHANISM OF
THE NEUTRAL FORM OF FRUCTOSE-1,6-BISPHOSPHATASE
Authors : Zhang, Y.; Liang, J.-Y.; Huang, S.; Ke, H.; Lipscomb, W.N.
Deposited on : 1992-10-16
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	335	 75% 18% • 5%
1	B	335	 73% 19% • • 5%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6366 atoms, of which 1317 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FRUCTOSE 1,6-BISPHOSPHATASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	317	Total	C	H	N	O	S	0	0	1
			2962	1540	538	409	460	15			
1	B	318	Total	C	H	N	O	S	0	0	1
			2974	1549	539	410	461	15			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	20	GLN	GLU	conflict	UNP P00636
A	96	THR	SER	conflict	UNP P00636
A	199	ASN	ASP	conflict	UNP P00636
B	20	GLN	GLU	conflict	UNP P00636
B	96	THR	SER	conflict	UNP P00636
B	199	ASN	ASP	conflict	UNP P00636

- Molecule 2 is 1,6-di-O-phosphono-alpha-D-fructofuranose (CCD ID: AFP) (formula: $C_6H_{14}O_{12}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	O	P	0	1
			25	6	5	12	2		
2	B	1	Total	C	H	O	P	0	1
			25	6	5	12	2		

- Molecule 3 is 1,6-di-O-phosphono-beta-D-fructofuranose (CCD ID: FBP) (formula: $C_6H_{14}O_{12}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	O	P	0	1
			25	6	5	12	2		
3	B	1	Total	C	H	O	P	0	1
			25	6	5	12	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	56	Total	H	O	0	0
			168	112	56		
4	B	54	Total	H	O	0	0
			162	108	54		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: FRUCTOSE 1,6-BISPHOSPHATASE



• Molecule 1: FRUCTOSE 1,6-BISPHOSPHATASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	131.20Å 131.20Å 69.30Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.50	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.193 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6366	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AFP, FBP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	5/2463 (0.2%)	1.55	23/3330 (0.7%)
1	B	0.76	6/2475 (0.2%)	1.60	27/3346 (0.8%)
All	All	0.78	11/4938 (0.2%)	1.57	50/6676 (0.7%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	253	HIS	CD2-NE2	-7.00	1.30	1.37
1	A	311	HIS	CD2-NE2	-6.97	1.30	1.37
1	B	311	HIS	CD2-NE2	-6.91	1.30	1.37
1	B	54	ALA	C-N	-6.74	1.24	1.33
1	A	54	ALA	C-N	-6.33	1.24	1.33
1	A	334	HIS	CD2-NE2	-6.19	1.31	1.37
1	A	253	HIS	CD2-NE2	-6.06	1.31	1.37
1	B	334	HIS	CD2-NE2	-5.73	1.31	1.37
1	B	82	ILE	CA-CB	5.71	1.61	1.54
1	A	311	HIS	CG-ND1	-5.34	1.32	1.38
1	B	311	HIS	CG-ND1	-5.28	1.32	1.38

All (50) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	179	ASN	OD1-CG-ND2	-9.36	113.24	122.60
1	B	179	ASN	CA-CB-CG	9.32	121.92	112.60
1	A	199	ASN	OD1-CG-ND2	-8.86	113.74	122.60
1	B	179	ASN	CB-CG-ND2	8.30	128.85	116.40
1	B	198	ARG	N-CA-C	8.28	123.51	113.41
1	B	220	ASP	CA-CB-CG	8.25	120.85	112.60
1	A	267	ASN	CA-CB-CG	7.82	120.42	112.60
1	A	282	ASN	OD1-CG-ND2	-7.67	114.92	122.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	145	ASP	CA-CB-CG	7.62	120.22	112.60
1	B	298	GLY	N-CA-C	-7.61	104.89	113.79
1	B	278	LEU	N-CA-C	7.07	119.84	111.71
1	A	35	ASN	CA-CB-CG	-7.04	105.56	112.60
1	A	112	LYS	N-CA-C	6.94	121.58	113.18
1	B	20	GLN	OE1-CD-NE2	-6.90	115.70	122.60
1	A	203	LYS	CA-CB-CG	6.53	127.17	114.10
1	B	112	LYS	N-CA-C	6.45	120.82	113.15
1	A	79	ASP	CA-CB-CG	6.43	119.03	112.60
1	A	267	ASN	OD1-CG-ND2	-6.32	116.28	122.60
1	A	199	ASN	CB-CG-ND2	6.28	125.82	116.40
1	B	74	ASP	CA-CB-CG	6.24	118.84	112.60
1	B	113	TYR	N-CA-C	6.22	119.84	109.95
1	B	51	ALA	N-CA-C	6.20	118.83	111.33
1	A	233	PRO	CA-C-N	6.07	125.29	118.97
1	A	233	PRO	C-N-CA	6.07	125.29	118.97
1	A	228	GLN	OE1-CD-NE2	-5.86	116.74	122.60
1	B	281	CYS	N-CA-C	5.86	117.33	111.07
1	B	282	ASN	OD1-CG-ND2	-5.82	116.78	122.60
1	A	282	ASN	CB-CG-ND2	5.75	125.03	116.40
1	B	121	ASP	CA-C-O	-5.73	114.11	120.70
1	A	118	ASP	N-CA-C	-5.72	99.60	109.15
1	B	268	LYS	CA-CB-CG	5.69	125.47	114.10
1	A	317	ILE	N-CA-C	-5.59	99.44	107.78
1	B	267	ASN	CA-CB-CG	5.54	118.14	112.60
1	B	78	ASN	OD1-CG-ND2	-5.46	117.14	122.60
1	A	118	ASP	CA-CB-CG	5.43	118.03	112.60
1	B	154	GLN	OE1-CD-NE2	-5.42	117.18	122.60
1	B	7	ASP	CA-CB-CG	5.38	117.98	112.60
1	B	125	ASN	OD1-CG-ND2	-5.37	117.23	122.60
1	B	238	ALA	CA-C-N	5.31	125.60	119.92
1	B	238	ALA	C-N-CA	5.31	125.60	119.92
1	A	83	ASN	CA-CB-CG	-5.26	107.34	112.60
1	B	253	HIS	CE1-NE2-CD2	5.25	114.25	109.00
1	A	20	GLN	OE1-CD-NE2	-5.17	117.43	122.60
1	A	101	ASN	OD1-CG-ND2	-5.15	117.45	122.60
1	A	49	ARG	CA-CB-CG	5.12	124.35	114.10
1	B	27	THR	N-CA-C	5.10	121.67	110.80
1	A	70	VAL	N-CA-C	-5.04	98.86	109.34
1	A	149	GLU	CA-CB-CG	5.03	124.16	114.10
1	B	200	VAL	N-CA-C	5.03	115.58	109.30
1	A	277	LEU	N-CA-C	5.01	116.55	111.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2424	538	2482	21	0
1	B	2435	539	2491	26	0
2	A	20	5	10	1	0
2	B	20	5	10	0	0
3	A	20	5	10	1	0
3	B	20	5	10	0	0
4	A	56	112	0	0	0
4	B	54	108	0	0	0
All	All	5049	1317	5013	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:78:ASN:OD1	1:B:96:THR:HG21	1.86	0.76
1:A:125:ASN:HB3	1:A:130:VAL:HB	1.69	0.74
1:B:70:VAL:HG22	1:B:124:SER:HA	1.76	0.68
1:B:125:ASN:HB3	1:B:130:VAL:HB	1.75	0.67
1:A:78:ASN:OD1	1:A:96:THR:HG21	1.96	0.66
1:A:96:THR:HG22	1:A:98:GLU:H	1.61	0.64
1:A:210:SER:HB3	1:A:243:ARG:O	1.97	0.63
1:B:195:LEU:HD21	1:B:198:ARG:HG3	1.82	0.61
1:A:122:GLY:HA2	2:A:336[A]:AFP:O1P	2.00	0.61
1:B:114:VAL:HB	1:B:139:TYR:HB2	1.83	0.59
3:A:337[B]:FBP:O4P	1:B:243:ARG:NH2	2.41	0.54
1:B:226:TYR:CZ	1:B:230:LYS:HE2	2.44	0.52
1:A:317:ILE:HG21	1:A:327:LEU:HD23	1.92	0.52
1:A:126:ILE:HG12	1:A:132:ILE:HD13	1.92	0.51
1:A:166:LEU:HD23	1:B:129:LEU:HD12	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:166:LEU:HD13	1:A:249:VAL:HG12	1.95	0.49
1:B:96:THR:HG23	1:B:117:PHE:CE1	2.47	0.49
1:B:181:VAL:HG21	1:B:291:ALA:HB2	1.94	0.48
1:B:40:ALA:HB2	1:B:84:VAL:HG21	1.95	0.48
1:A:9:ASN:HD21	1:A:15:ARG:NH2	2.12	0.48
1:B:43:ALA:HB3	1:B:80:LEU:HD21	1.96	0.48
1:A:229:ARG:HH11	1:A:330:ILE:HD11	1.79	0.47
1:A:296:THR:HG21	1:A:328:LEU:HD21	1.97	0.46
1:B:248:MET:HE3	1:B:275:LEU:HD13	1.96	0.46
1:A:233:PRO:HA	1:A:234:PRO:HD2	1.93	0.44
1:B:242:ALA:O	1:B:243:ARG:HD3	2.17	0.44
1:A:40:ALA:O	1:A:44:ILE:HG13	2.18	0.44
1:A:252:VAL:HG11	1:A:284:MET:SD	2.58	0.44
1:A:297:THR:HB	1:A:315:PRO:HG2	1.99	0.43
1:B:100:LYS:O	1:B:310:ILE:HD11	2.18	0.43
1:A:73:LEU:HD23	1:A:126:ILE:HD13	2.01	0.43
1:B:268:LYS:HZ2	1:B:269:LYS:H	1.66	0.43
1:B:166:LEU:HD13	1:B:249:VAL:HG12	2.00	0.42
1:B:226:TYR:CE1	1:B:230:LYS:HE2	2.54	0.42
1:B:263:MET:HB3	1:B:265:PRO:HD3	2.02	0.42
1:A:71:LYS:NZ	1:A:71:LYS:HB2	2.34	0.42
1:A:169:SER:HB2	1:B:48:VAL:HG12	2.02	0.42
1:B:229:ARG:CG	1:B:234:PRO:HD3	2.50	0.41
1:B:68:ASP:N	1:B:71:LYS:O	2.54	0.41
1:B:96:THR:HG22	1:B:98:GLU:H	1.85	0.41
1:B:71:LYS:HA	1:B:71:LYS:HD2	1.87	0.41
1:A:190:ILE:HD11	1:A:194:ILE:HD11	2.02	0.41
1:A:77:SER:O	1:A:81:VAL:HG23	2.20	0.41
1:B:229:ARG:HG3	1:B:234:PRO:HD3	2.02	0.41
1:B:204:LYS:O	1:B:322:GLU:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/335 (93%)	294 (94%)	15 (5%)	4 (1%)	9	18
1	B	314/335 (94%)	293 (93%)	18 (6%)	3 (1%)	12	24
All	All	627/670 (94%)	587 (94%)	33 (5%)	7 (1%)	11	22

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	LYS
1	A	268	LYS
1	B	27	THR
1	B	72	LYS
1	A	54	ALA
1	A	333	LYS
1	B	178	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/278 (95%)	250 (94%)	15 (6%)	18	39
1	B	266/278 (96%)	249 (94%)	17 (6%)	16	33
All	All	531/556 (96%)	499 (94%)	32 (6%)	17	36

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	ILE
1	A	20	GLN
1	A	35	ASN
1	A	71	LYS
1	A	98	GLU
1	A	123	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	145	ASP
1	A	210	SER
1	A	224	THR
1	A	230	LYS
1	A	237	SER
1	A	265	PRO
1	A	272	LYS
1	A	300	GLU
1	A	317	ILE
1	B	10	ILE
1	B	20	GLN
1	B	23	LYS
1	B	97	GLU
1	B	108	GLU
1	B	123	SER
1	B	124	SER
1	B	143	SER
1	B	179	ASN
1	B	198	ARG
1	B	204	LYS
1	B	207	SER
1	B	243	ARG
1	B	268	LYS
1	B	270	SER
1	B	276	ARG
1	B	322	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	ASN
1	A	32	GLN
1	A	83	ASN
1	A	101	ASN
1	A	154	GLN
1	A	158	ASN
1	A	179	ASN
1	A	182	ASN
1	A	199	ASN
1	A	228	GLN
1	B	142	ASN
1	B	154	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	182	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	AFP	B	336[A]	-	18,20,20	0.75	0	21,32,32	0.96	1 (4%)
3	FBP	B	337[B]	-	18,20,20	0.66	0	21,32,32	0.96	1 (4%)
2	AFP	A	336[A]	-	18,20,20	0.73	0	21,32,32	0.99	1 (4%)
3	FBP	A	337[B]	-	18,20,20	0.66	0	21,32,32	0.96	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AFP	B	336[A]	-	-	12/13/32/32	0/1/1/1
3	FBP	B	337[B]	-	-	9/13/32/32	0/1/1/1
2	AFP	A	336[A]	-	-	3/13/32/32	0/1/1/1
3	FBP	A	337[B]	-	-	3/13/32/32	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	336[A]	AFP	O2P-P1-O1P	2.16	119.25	110.83
3	A	337[B]	FBP	O2P-P1-O1P	2.12	119.11	110.83
3	B	337[B]	FBP	O2P-P1-O1P	2.10	119.03	110.83
2	B	336[A]	AFP	O2P-P1-O1P	2.10	119.02	110.83

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	336[A]	AFP	C1-O1-P1-O2P
2	A	336[A]	AFP	C1-O1-P1-O3P
2	B	336[A]	AFP	C1-O1-P1-O2P
2	B	336[A]	AFP	C1-O1-P1-O3P
2	B	336[A]	AFP	O1-C1-C2-O2
2	B	336[A]	AFP	O1-C1-C2-C3
2	B	336[A]	AFP	C6-O6-P6-O4P
2	B	336[A]	AFP	C6-O6-P6-O5P
2	B	336[A]	AFP	C6-O6-P6-O6P
3	A	337[B]	FBP	O1-C1-C2-O2
3	A	337[B]	FBP	O1-C1-C2-C3
3	A	337[B]	FBP	O1-C1-C2-O5
3	B	337[B]	FBP	O1-C1-C2-O2
3	B	337[B]	FBP	O1-C1-C2-C3
3	B	337[B]	FBP	O1-C1-C2-O5
3	B	337[B]	FBP	C6-O6-P2-O4P
3	B	337[B]	FBP	C6-O6-P2-O5P
3	B	337[B]	FBP	C6-O6-P2-O6P
3	B	337[B]	FBP	C4-C5-C6-O6
2	B	336[A]	AFP	C4-C5-C6-O6
2	A	336[A]	AFP	C1-O1-P1-O1P
3	B	337[B]	FBP	O5-C5-C6-O6
2	B	336[A]	AFP	O5-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	336[A]	AFP	O1-C1-C2-O5
2	B	336[A]	AFP	C1-O1-P1-O1P
3	B	337[B]	FBP	C1-O1-P1-O2P
2	B	336[A]	AFP	C2-C1-O1-P1

There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	336[A]	AFP	1	0
3	A	337[B]	FBP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand FBP B 337 (B)



Ligand AFP A 336 (A)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.