



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:02 AM UTC

PDB ID : 3FB6 / pdb_00003fb6
Title : KcsA Potassium channel in the partially open state with 16 Å opening at T112
Authors : Cuello, L.G.; Jogini, V.; Cortes, D.M.; Perozo, E.
Deposited on : 2008-11-18
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	16% 77% 21% .
2	B	212	11% 72% 27% .
3	C	104	6% 64% 19% . 13%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3934 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called antibody fab fragment heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	219	Total	C	N	O	S	0	0	0
			1648	1042	275	325	6			

- Molecule 2 is a protein called antibody fab fragment light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	212	Total	C	N	O	S	0	0	0
			1649	1023	283	338	5			

- Molecule 3 is a protein called Voltage-gated potassium channel.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	90	Total	C	N	O	S	0	0	0
			632	415	103	112	2			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	25	GLN	HIS	engineered mutation	UNP P0A334
C	90	CYS	LEU	engineered mutation	UNP P0A334
C	117	GLN	ARG	engineered mutation	UNP P0A334
C	120	GLN	GLU	engineered mutation	UNP P0A334
C	121	GLN	ARG	engineered mutation	UNP P0A334
C	122	GLN	ARG	engineered mutation	UNP P0A334
C	124	GLN	HIS	engineered mutation	UNP P0A334


- Molecule 4 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	5	Total	K	0	0
			5	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: antibody fab fragment heavy chain



- Molecule 2: antibody fab fragment light chain



- Molecule 3: Voltage-gated potassium channel



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	155.71 Å 155.71 Å 74.07 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 3.00 40.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-3.00) 95.4 (40.00-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 3.00 Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.248 , 0.255 0.248 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	88.7	Xtriage
Anisotropy	0.249	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 60.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.029 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3934	wwPDB-VP
Average B, all atoms (Å ²)	103.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.39	0/1692	0.91	5/2312 (0.2%)
2	B	0.36	0/1686	0.87	3/2287 (0.1%)
3	C	0.50	0/645	1.16	6/891 (0.7%)
All	All	0.40	0/4023	0.94	14/5490 (0.3%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	60	ILE	N-CA-C	8.56	120.95	111.88
1	A	103	GLY	N-CA-C	7.95	124.54	115.08
3	C	29	ALA	N-CA-C	-7.27	104.21	113.23
2	B	137	ASN	N-CA-C	6.74	120.81	109.76
3	C	30	GLY	N-CA-C	-6.57	107.17	115.31
3	C	51	GLU	N-CA-C	6.15	118.06	111.36
1	A	96	CYS	N-CA-C	-5.92	100.25	109.72
3	C	28	ALA	N-CA-C	-5.87	105.96	113.23
1	A	139	SER	N-CA-C	-5.29	106.78	113.18
3	C	31	ALA	N-CA-C	-5.28	106.35	112.89
2	B	99	GLY	N-CA-C	-5.14	104.58	112.85
1	A	68	ALA	N-CA-C	5.09	118.05	109.95
1	A	8	GLY	N-CA-C	5.04	118.77	112.73
2	B	138	ASN	N-CA-C	5.03	118.19	111.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1648	0	1620	41	3
2	B	1649	0	1580	60	3
3	C	632	0	635	18	0
4	C	5	0	0	0	0
All	All	3934	0	3835	114	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (114) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:134:CYS:HG	2:B:194:CYS:HG	0.94	0.87
2:B:167:ASP:OD2	2:B:169:LYS:HB2	1.79	0.83
1:A:65:GLN:NE2	1:A:65:GLN:H	1.79	0.80
1:A:120:LYS:H	1:A:120:LYS:HE3	1.49	0.78
1:A:22:CYS:HG	1:A:96:CYS:HG	0.88	0.77
2:B:74:SER:C	2:B:75:ILE:HD12	2.12	0.74
2:B:21:PHE:HE1	2:B:75:ILE:HD13	1.56	0.70
2:B:183:LYS:O	2:B:187:GLU:HG2	1.91	0.70
1:A:140:MET:HE3	1:A:187:THR:HG22	1.74	0.70
3:C:86:LEU:HD12	3:C:86:LEU:O	1.93	0.69
1:A:124:PRO:HB3	1:A:150:TYR:HB3	1.74	0.68
2:B:2:ILE:HD11	2:B:93:ARG:HB2	1.75	0.67
1:A:51:ILE:HD13	1:A:52:ILE:C	2.21	0.66
3:C:91:VAL:O	3:C:95:VAL:HG23	1.96	0.66
2:B:183:LYS:HB3	2:B:183:LYS:NZ	2.11	0.65
2:B:58:ILE:HD12	2:B:58:ILE:N	2.11	0.65
1:A:31:SER:HB2	3:C:62:TYR:CE1	2.31	0.65
2:B:61:ARG:HG3	2:B:76:ASN:O	1.96	0.64
1:A:148:LYS:HB3	1:A:181:THR:HG23	1.79	0.64
2:B:37:GLN:HB2	2:B:47:LEU:HD11	1.79	0.63
1:A:120:LYS:HE3	1:A:120:LYS:N	2.14	0.62
2:B:21:PHE:HE1	2:B:75:ILE:CD1	2.13	0.61
2:B:92:ASN:HD21	3:C:58:GLN:NE2	1.98	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:120:LYS:H	1:A:120:LYS:CE	2.13	0.60
3:C:89:ARG:O	3:C:93:VAL:HG23	2.01	0.60
2:B:2:ILE:CD1	2:B:93:ARG:HB2	2.32	0.59
2:B:13:VAL:CG2	2:B:17:GLU:HB3	2.31	0.59
3:C:44:SER:OG	3:C:66:LEU:HA	2.03	0.58
2:B:75:ILE:HD12	2:B:75:ILE:N	2.18	0.58
1:A:182:LEU:C	1:A:182:LEU:HD12	2.29	0.58
2:B:48:ILE:HD12	2:B:73:LEU:CD1	2.33	0.58
2:B:55:ILE:HB	2:B:58:ILE:HD13	1.86	0.57
1:A:34:ILE:HD13	1:A:79:ALA:HB2	1.86	0.57
1:A:6:GLN:HE21	1:A:109:GLY:HA3	1.69	0.57
2:B:124:GLN:HG2	2:B:129:GLY:O	2.05	0.57
1:A:23:LYS:HD3	1:A:78:THR:CG2	2.35	0.57
3:C:29:ALA:C	3:C:31:ALA:H	2.11	0.57
2:B:2:ILE:HD12	2:B:2:ILE:H	1.70	0.56
2:B:2:ILE:HD12	2:B:2:ILE:N	2.21	0.56
1:A:64:ILE:HD12	1:A:65:GLN:O	2.06	0.56
1:A:215:ILE:HD12	1:A:215:ILE:N	2.21	0.55
2:B:85:ASN:ND2	2:B:103:LYS:HD3	2.22	0.55
2:B:21:PHE:CE1	2:B:75:ILE:HD13	2.41	0.54
1:A:101:GLY:HA3	3:C:62:TYR:CE1	2.43	0.53
2:B:19:VAL:HG12	2:B:75:ILE:HB	1.90	0.53
2:B:2:ILE:HD11	2:B:93:ARG:HD2	1.90	0.53
1:A:51:ILE:HD13	1:A:52:ILE:N	2.24	0.52
1:A:83:LEU:HB3	1:A:86:LEU:HD21	1.92	0.52
2:B:18:ARG:NH1	2:B:76:ASN:OD1	2.43	0.51
2:B:93:ARG:HG2	3:C:58:GLN:HG2	1.91	0.51
3:C:86:LEU:HD12	3:C:86:LEU:C	2.36	0.51
2:B:58:ILE:N	2:B:58:ILE:CD1	2.72	0.50
1:A:23:LYS:HD3	1:A:78:THR:HG23	1.94	0.50
1:A:65:GLN:H	1:A:65:GLN:HE21	1.58	0.50
3:C:51:GLU:HG3	3:C:59:LEU:HB3	1.94	0.50
1:A:17:SER:CB	1:A:84:SER:HA	2.42	0.50
2:B:79:GLU:HG3	2:B:81:GLU:HG2	1.95	0.49
2:B:115:VAL:HG22	2:B:136:LEU:CD1	2.43	0.49
2:B:134:CYS:HG	2:B:194:CYS:CB	2.26	0.48
3:C:29:ALA:C	3:C:31:ALA:N	2.69	0.48
2:B:142:LYS:HB3	2:B:173:TYR:CG	2.49	0.48
2:B:136:LEU:HD22	2:B:136:LEU:N	2.29	0.48
1:A:148:LYS:HB2	1:A:148:LYS:NZ	2.29	0.47
1:A:1:GLN:OE1	1:A:1:GLN:N	2.42	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:150:ILE:HD12	2:B:155:ARG:HH11	1.78	0.47
2:B:13:VAL:HG22	2:B:14:SER:N	2.29	0.47
1:A:1:GLN:N	1:A:1:GLN:CD	2.73	0.47
1:A:12:VAL:O	1:A:116:VAL:HA	2.15	0.46
2:B:142:LYS:HD3	2:B:173:TYR:CE2	2.50	0.46
2:B:92:ASN:ND2	3:C:58:GLN:NE2	2.64	0.46
1:A:34:ILE:CD1	1:A:79:ALA:HB2	2.46	0.45
2:B:13:VAL:HG23	2:B:17:GLU:CD	2.40	0.45
1:A:198:VAL:HG12	1:A:215:ILE:HD13	1.97	0.45
2:B:83:ILE:HD11	2:B:106:ILE:HD12	1.99	0.45
2:B:55:ILE:HB	2:B:58:ILE:CD1	2.45	0.45
2:B:209:PHE:CD1	2:B:209:PHE:C	2.96	0.44
2:B:48:ILE:HD12	2:B:73:LEU:HD13	1.98	0.44
1:A:120:LYS:H	1:A:120:LYS:CD	2.29	0.44
1:A:98:ARG:HD3	1:A:100:ARG:CZ	2.48	0.44
2:B:40:THR:HG22	2:B:41:ASN:ND2	2.33	0.44
1:A:3:GLN:O	1:A:24:ALA:HA	2.18	0.44
3:C:62:TYR:HB2	3:C:63:PRO:HD3	2.00	0.44
2:B:84:ALA:O	2:B:103:LYS:HD2	2.17	0.43
2:B:183:LYS:HB3	2:B:183:LYS:HZ2	1.82	0.43
1:A:3:GLN:O	1:A:4:LEU:HD12	2.18	0.43
1:A:52:ILE:O	1:A:52:ILE:HG23	2.18	0.43
2:B:193:THR:HG22	2:B:208:SER:CB	2.49	0.43
2:B:193:THR:HG22	2:B:208:SER:HB3	1.99	0.43
1:A:124:PRO:CB	1:A:150:TYR:HB3	2.46	0.43
2:B:139:PHE:HE1	2:B:142:LYS:HA	1.83	0.43
1:A:35:HIS:CG	1:A:105:PHE:HE2	2.37	0.42
2:B:18:ARG:NH1	2:B:76:ASN:CG	2.77	0.42
2:B:75:ILE:CD1	2:B:75:ILE:N	2.81	0.42
2:B:118:PHE:HA	2:B:119:PRO:HD3	1.87	0.42
2:B:61:ARG:NH1	2:B:82:ASP:OD1	2.53	0.42
2:B:83:ILE:HD13	2:B:166:GLN:HG2	2.01	0.42
1:A:51:ILE:HD13	1:A:51:ILE:C	2.45	0.42
1:A:140:MET:CE	1:A:187:THR:HG22	2.48	0.42
2:B:21:PHE:CE1	2:B:75:ILE:CD1	2.99	0.42
2:B:55:ILE:O	2:B:58:ILE:HD13	2.19	0.42
3:C:96:MET:O	3:C:100:ILE:HG13	2.19	0.42
1:A:128:PRO:HG3	1:A:213:LYS:HD2	2.01	0.42
1:A:23:LYS:HD3	1:A:78:THR:OG1	2.20	0.41
2:B:55:ILE:CB	2:B:58:ILE:HD13	2.50	0.41
3:C:31:ALA:O	3:C:34:VAL:HB	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:13:VAL:HG22	2:B:17:GLU:HB3	1.99	0.41
2:B:136:LEU:CD1	2:B:196:ALA:HB2	2.51	0.41
2:B:135:PHE:C	2:B:136:LEU:HD22	2.46	0.41
3:C:62:TYR:O	3:C:63:PRO:C	2.61	0.41
1:A:40:ARG:NH2	1:A:89:GLU:HB3	2.36	0.40
1:A:91:SER:O	1:A:92:ALA:HB2	2.21	0.40
2:B:85:ASN:ND2	2:B:103:LYS:CD	2.84	0.40
3:C:66:LEU:O	3:C:69:SER:OG	2.38	0.40
2:B:16:GLY:C	2:B:77:SER:HA	2.46	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:65:GLN:OE1	2:B:18:ARG:NE[3_755]	1.37	0.83
1:A:65:GLN:OE1	2:B:18:ARG:CD[3_755]	1.70	0.50
1:A:65:GLN:OE1	2:B:18:ARG:CZ[3_755]	1.81	0.39

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/219 (99%)	212 (98%)	5 (2%)	0	100	100
2	B	210/212 (99%)	203 (97%)	7 (3%)	0	100	100
3	C	88/104 (85%)	84 (96%)	2 (2%)	2 (2%)	5	25
All	All	515/535 (96%)	499 (97%)	14 (3%)	2 (0%)	30	65

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	77	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	113	TRP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/185 (100%)	178 (96%)	7 (4%)	29	63
2	B	190/190 (100%)	184 (97%)	6 (3%)	34	67
3	C	58/75 (77%)	58 (100%)	0	100	100
All	All	433/450 (96%)	420 (97%)	13 (3%)	36	69

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	ILE
1	A	65	GLN
1	A	120	LYS
1	A	148	LYS
1	A	152	PRO
1	A	154	PRO
1	A	194	PRO
2	B	4	LEU
2	B	58	ILE
2	B	75	ILE
2	B	95	PRO
2	B	165	ASP
2	B	183	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	5	GLN
1	A	6	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	39	GLN
1	A	59	ASN
1	A	65	GLN
2	B	37	GLN
2	B	38	GLN
2	B	41	ASN
2	B	85	ASN
2	B	92	ASN
2	B	190	ASN
3	C	58	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	219/219 (100%)	1.08	35 (15%) 5 3	76, 111, 136, 149	0
2	B	212/212 (100%)	0.98	23 (10%) 11 6	67, 108, 143, 158	0
3	C	90/104 (86%)	0.53	6 (6%) 24 12	56, 74, 120, 138	0
All	All	521/535 (97%)	0.95	64 (12%) 8 5	56, 107, 141, 158	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	112	THR	4.2
2	B	145	ASN	4.1
1	A	136	GLN	3.3
3	C	114	PHE	3.3
1	A	28	THR	3.3
2	B	118	PHE	3.3
2	B	209	PHE	3.3
1	A	55	TYR	3.2
1	A	2	VAL	3.2
3	C	113	TRP	3.1
2	B	157	ASN	2.9
1	A	165	SER	2.9
1	A	129	LEU	2.8
2	B	194	CYS	2.8
1	A	3	GLN	2.8
1	A	124	PRO	2.7
1	A	56	GLY	2.7
1	A	27	TYR	2.7
2	B	183	LYS	2.7
1	A	131	PRO	2.7
1	A	183	SER	2.7
2	B	160	LEU	2.6
3	C	110	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	165	ASP	2.6
1	A	1	GLN	2.6
2	B	176	SER	2.6
1	A	215	ILE	2.6
1	A	196	GLU	2.6
1	A	128	PRO	2.6
3	C	26	TRP	2.6
2	B	75	ILE	2.5
1	A	29	PHE	2.5
1	A	24	ALA	2.5
2	B	128	GLY	2.5
1	A	30	THR	2.4
2	B	143	ASP	2.4
1	A	133	SER	2.4
2	B	103	LYS	2.3
1	A	4	LEU	2.3
1	A	197	THR	2.3
2	B	105	GLU	2.3
1	A	25	SER	2.3
1	A	163	SER	2.3
1	A	65	GLN	2.3
1	A	132	GLY	2.2
1	A	135	ALA	2.2
2	B	192	TYR	2.2
1	A	145	CYS	2.2
2	B	23	CYS	2.2
2	B	55	ILE	2.2
2	B	83	ILE	2.2
2	B	178	THR	2.1
2	B	1	ASP	2.1
1	A	31	SER	2.1
1	A	89	GLU	2.1
2	B	120	PRO	2.1
2	B	117	ILE	2.1
1	A	13	LYS	2.1
1	A	63	LYS	2.0
2	B	182	THR	2.0
1	A	166	SER	2.0
1	A	26	GLY	2.0
3	C	77	GLY	2.0
1	A	146	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	K	C	3	1/1	0.77	0.24	86,86,86,86	1
4	K	C	1	1/1	0.83	0.29	100,100,100,100	1
4	K	C	2	1/1	0.91	0.15	80,80,80,80	1
4	K	C	4	1/1	0.91	0.26	64,64,64,64	1
4	K	C	5	1/1	0.94	0.35	129,129,129,129	1

6.5 Other polymers [i](#)

There are no such residues in this entry.