



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:14 PM UTC

PDB ID : 2FCC / pdb_00002fcc
Title : Crystal Structure of T4 Pyrimidine Dimer Glycosylase (T4-Pdg) Covalently Complexed with a DNA Substrate Containing Abasic Site
Authors : Golan, G.; Zharkov, D.O.; Fernandes, A.S.; Dodson, M.L.; McCullough, A.K.; Grollman, A.P.; Lloyd, R.S.; Shoham, G.
Deposited on : 2005-12-12
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	13	
1	E	13	
2	D	13	
2	F	13	
3	A	137	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	B	137	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment on the left labeled '33%', a green segment in the middle labeled '76%', and a yellow segment on the right labeled '23%'. A small black dot is located at the far right end of the bar.</p>

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 3511 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*CP*CP*AP*GP*GP*AP*(PED)P*GP*AP*AP*GP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	C	13	Total 256	C 121	N 52	O 71	P 12	0	0	0
1	E	13	Total 256	C 121	N 52	O 71	P 12	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*GP*CP*(BRU)P*(BRU)P*CP*AP*(BRU)P*CP*CP*(BRU)P*GP*G)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	Br	C	N	O	P			
2	D	13	Total 262	Br 4	C 122	N 45	O 79	P 12	0	0	0
2	F	13	Total 262	Br 4	C 122	N 45	O 79	P 12	0	0	0

- Molecule 3 is a protein called Endonuclease V.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	137	Total 1137	C 730	N 203	O 202	S 2	0	1	0
3	B	137	Total 1133	C 728	N 203	O 200	S 2	0	1	0

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	C	14	Total O 14 14	0	0
6	D	12	Total O 12 12	0	0
6	E	3	Total O 3 3	0	0
6	F	2	Total O 2 2	0	0
6	A	98	Total O 98 98	0	0
6	B	54	Total O 54 54	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (5'-D(*CP*CP*AP*GP*GP*AP*(PED)P*GP*AP*AP*GP*CP*C)-3')




- Molecule 1: DNA (5'-D(*CP*CP*AP*GP*GP*AP*(PED)P*GP*AP*AP*GP*CP*C)-3')




- Molecule 2: DNA (5'-D(*GP*GP*CP*(BRU)P*(BRU)P*CP*AP*(BRU)P*CP*CP*(BRU)P*GP*G)-3')



- Molecule 2: DNA (5'-D(*GP*GP*CP*(BRU)P*(BRU)P*CP*AP*(BRU)P*CP*CP*(BRU)P*GP*G)-3')





- Molecule 3: Endonuclease V





- Molecule 3: Endonuclease V



4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α , β , γ	184.06Å 184.06Å 100.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.85 – 2.30 39.85 – 2.30	Depositor EDS
% Data completeness (in resolution range)	96.1 (39.85-2.30) 95.9 (39.85-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.19 (at 2.31Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.249 , 0.274 (Not available) , 0.268	Depositor DCC
R_{free} test set	4184 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	32.3	Xtrriage
Anisotropy	0.159	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3511	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PED, BRU, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	C	0.28	0/275	0.76	1/420 (0.2%)
1	E	0.28	0/275	0.73	1/420 (0.2%)
2	D	0.31	0/201	0.70	0/301
2	F	0.27	0/201	0.61	0/301
3	A	0.50	1/1162 (0.1%)	0.86	2/1565 (0.1%)
3	B	0.48	2/1158 (0.2%)	0.83	1/1560 (0.1%)
All	All	0.44	3/3272 (0.1%)	0.81	5/4567 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	E	0	2
All	All	0	3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	71	GLN	CD-OE1	5.61	1.34	1.23
3	B	115	GLN	CD-OE1	5.14	1.33	1.23
3	A	71	GLN	CD-OE1	5.12	1.33	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	206	DA	N9-C1'-C2'	-6.11	104.34	113.50
3	A	101	ARG	N-CA-C	5.79	117.95	110.65

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	52	LEU	N-CA-C	-5.64	101.85	110.14
3	B	36	ALA	N-CA-C	-5.19	107.24	113.21
1	E	206	DA	N9-C1'-C2'	-5.17	105.75	113.50

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	206	DA	Sidechain
1	E	205	DG	Sidechain
1	E	206	DA	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	256	0	143	9	0
1	E	256	0	143	12	0
2	D	262	0	137	9	0
2	F	262	0	137	9	1
3	A	1137	0	1146	17	0
3	B	1133	0	1142	25	0
4	A	10	0	0	0	0
5	A	6	0	8	2	0
5	B	6	0	8	2	0
6	A	98	0	0	1	0
6	B	54	0	0	1	0
6	C	14	0	0	1	0
6	D	12	0	0	1	0
6	E	3	0	0	0	0
6	F	2	0	0	0	0
All	All	3511	0	2864	74	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:218:BRU:H2''	2:F:219:DC:H5''	1.14	1.09
2:F:218:BRU:C2'	2:F:219:DC:H5''	1.93	0.97
2:D:218:BRU:H2''	2:D:219:DC:H5'	1.48	0.95
2:F:220:DA:H2''	2:F:221:BRU:H5''	1.49	0.92
2:F:220:DA:H2''	2:F:221:BRU:C5'	2.04	0.87
3:B:125:ARG:HH11	3:B:125:ARG:HG3	1.40	0.84
2:F:218:BRU:H2''	2:F:219:DC:C5'	2.06	0.80
3:B:104:TYR:CE2	3:B:106:PRO:HG3	2.22	0.74
1:C:211:DG:H2''	1:C:212:DC:H5'	1.68	0.73
2:D:215:DG:H2''	2:D:216:DC:C6	2.24	0.72
1:E:207:PED:H1'2	3:B:23:GLU:OE1	1.95	0.67
2:D:220:DA:H2''	2:D:221:BRU:H5''	1.80	0.64
3:B:125:ARG:HG3	3:B:125:ARG:NH1	2.12	0.63
2:D:220:DA:H2''	2:D:221:BRU:C5'	2.30	0.62
1:C:206:DA:H2'	6:C:1121:HOH:O	1.99	0.61
1:E:212:DC:H2''	1:E:213:DC:C6	2.37	0.60
3:B:20:GLU:HG2	3:B:74:LEU:HD11	1.85	0.59
3:A:104:TYR:CE2	3:A:106:PRO:HG3	2.39	0.58
2:D:218:BRU:C2'	2:D:219:DC:H5'	2.28	0.58
2:F:220:DA:H2''	2:F:221:BRU:H5'	1.85	0.58
3:A:51:ILE:H	3:A:51:ILE:HD13	1.68	0.57
3:A:70:ARG:HH11	5:A:602:GOL:H12	1.69	0.56
1:E:209:DA:H5''	3:B:53:GLY:O	2.05	0.56
3:B:98:GLN:HG2	3:B:101:ARG:NH2	2.20	0.56
1:E:211:DG:H5''	3:B:34:HIS:CE1	2.41	0.56
1:C:207:PED:O4'	3:A:23:GLU:OE2	2.23	0.55
3:A:24:LEU:C	3:A:24:LEU:HD23	2.32	0.54
1:C:213:DC:H5'	1:C:213:DC:H6	1.73	0.54
3:B:24:LEU:HB3	3:B:25:PRO:HD3	1.89	0.54
2:F:218:BRU:C3'	2:F:219:DC:H5''	2.37	0.54
1:E:207:PED:O4'	3:B:23:GLU:OE2	2.25	0.53
3:A:51:ILE:HD13	3:A:51:ILE:N	2.23	0.53
3:B:35:VAL:C	3:B:37:ASN:H	2.18	0.52
3:B:72:ILE:HG13	3:B:90:VAL:HG21	1.92	0.50
3:B:34:HIS:HB3	3:B:39:LYS:HG3	1.94	0.50
3:B:41:VAL:HG22	3:B:100:PHE:CD1	2.47	0.49
3:B:40:ARG:H	3:B:43:ASP:HB2	1.77	0.49
1:E:212:DC:H2''	1:E:213:DC:C5	2.47	0.48
3:A:109:ALA:O	3:A:113:ILE:HG13	2.13	0.48
1:C:211:DG:H2''	1:C:212:DC:C5'	2.40	0.47
3:B:125:ARG:NH1	3:B:125:ARG:CG	2.75	0.47
1:C:212:DC:H2''	1:C:213:DC:H5''	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:211:DG:H5''	3:B:34:HIS:NE2	2.31	0.46
3:B:41:VAL:HG23	3:B:99:GLU:CD	2.39	0.46
3:B:106:PRO:HA	5:B:601:GOL:H31	1.97	0.46
2:D:215:DG:H2''	2:D:216:DC:C5	2.51	0.46
3:B:24:LEU:HD23	3:B:24:LEU:C	2.40	0.46
2:D:225:DG:H2''	6:D:1169:HOH:O	2.15	0.45
1:E:207:PED:H1'2	1:E:207:PED:O4'	2.15	0.45
3:A:118:LEU:HD13	6:A:1107:HOH:O	2.16	0.45
1:E:212:DC:O2	2:F:215:DG:N2	2.48	0.44
3:A:70:ARG:HH11	5:A:602:GOL:C1	2.30	0.43
3:B:51:ILE:H	3:B:51:ILE:HD13	1.82	0.43
3:B:107:HIS:H	5:B:601:GOL:HO3	1.66	0.43
1:E:201:DC:H6	1:E:201:DC:H2'	1.59	0.43
2:D:215:DG:C2'	2:D:216:DC:C5	3.02	0.43
3:A:8:LEU:HB3	3:A:11:GLU:HG2	2.00	0.42
3:A:24:LEU:HB3	3:A:25:PRO:HD3	2.02	0.42
1:C:212:DC:H2''	1:C:213:DC:C5'	2.50	0.42
3:B:113:ILE:HG13	6:B:1171:HOH:O	2.19	0.42
3:B:30:ALA:O	3:B:33:LYS:HB3	2.20	0.42
3:A:120:GLU:O	3:A:124:GLN:HG3	2.20	0.41
1:C:212:DC:C2'	1:C:213:DC:H5''	2.50	0.41
3:A:41:VAL:HG23	3:A:99:GLU:HG3	2.01	0.41
2:D:215:DG:H4'	2:D:215:DG:OP1	2.21	0.41
3:B:122:ILE:HA	3:B:129:TYR:CE2	2.55	0.41
1:E:201:DC:H2''	1:E:202:DC:H5'	2.02	0.41
3:A:33:LYS:O	3:A:36:ALA:HB3	2.20	0.41
1:E:207:PED:O4'	1:E:207:PED:C1'	2.68	0.41
1:C:210:DA:H1'	1:C:211:DG:C8	2.56	0.41
3:A:62:ASP:OD1	3:A:63:LYS:HG3	2.21	0.41
3:A:15:GLN:HG2	3:A:128:TRP:CH2	2.57	0.40
3:A:45:LYS:O	3:A:46:ILE:HB	2.21	0.40
2:F:215:DG:H2''	2:F:216:DC:C5	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:214:DG:N2	2:F:214:DG:N2[12_565]	2.04	0.16

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	136/137 (99%)	133 (98%)	1 (1%)	2 (2%)	8	8
3	B	136/137 (99%)	128 (94%)	7 (5%)	1 (1%)	18	23
All	All	272/274 (99%)	261 (96%)	8 (3%)	3 (1%)	11	13

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	B	38	GLY
3	A	45	LYS
3	A	46	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	119/118 (101%)	116 (98%)	3 (2%)	42	60
3	B	118/118 (100%)	116 (98%)	2 (2%)	53	72
All	All	237/236 (100%)	232 (98%)	5 (2%)	47	66

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	8	LEU
3	A	51	ILE
3	A	118	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	B	43	ASP
3	B	51	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
3	B	37	ASN
3	B	107	HIS
3	B	124	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BRU	D	217	1,2	18,21,22	0.40	0	25,30,33	0.54	0
2	BRU	F	221	1,2	18,21,22	0.42	0	25,30,33	0.64	0
2	BRU	F	224	1,2	18,21,22	0.40	0	25,30,33	0.56	0
2	BRU	F	218	1,2	18,21,22	0.35	0	25,30,33	0.56	1 (4%)
2	BRU	F	217	2	18,21,22	0.36	0	25,30,33	0.58	0
2	BRU	D	218	1,2	18,21,22	0.46	0	25,30,33	0.57	0
2	BRU	D	221	1,2	18,21,22	0.48	0	25,30,33	0.71	0
2	BRU	D	224	1,2	18,21,22	0.42	0	25,30,33	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BRU	D	217	1,2	-	0/7/21/22	0/2/2/2
2	BRU	F	221	1,2	-	4/7/21/22	0/2/2/2
2	BRU	F	224	1,2	-	0/7/21/22	0/2/2/2
2	BRU	F	218	1,2	-	4/7/21/22	0/2/2/2
2	BRU	F	217	2	-	0/7/21/22	0/2/2/2
2	BRU	D	218	1,2	-	0/7/21/22	0/2/2/2
2	BRU	D	221	1,2	-	4/7/21/22	0/2/2/2
2	BRU	D	224	1,2	-	4/7/21/22	0/2/2/2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	218	BRU	BR-C5-C4	2.02	120.35	118.02

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	221	BRU	C2'-C1'-N1-C6
2	F	221	BRU	O4'-C1'-N1-C6
2	F	221	BRU	C2'-C1'-N1-C2
2	F	218	BRU	C2'-C1'-N1-C6
2	F	218	BRU	C2'-C1'-N1-C2
2	D	221	BRU	C2'-C1'-N1-C6
2	F	221	BRU	O4'-C1'-N1-C2
2	F	218	BRU	O4'-C1'-N1-C6
2	D	221	BRU	O4'-C1'-N1-C6
2	D	224	BRU	C2'-C1'-N1-C6
2	F	218	BRU	O4'-C1'-N1-C2
2	D	224	BRU	O4'-C1'-N1-C6
2	D	221	BRU	O4'-C1'-N1-C2
2	D	221	BRU	C2'-C1'-N1-C2
2	D	224	BRU	C2'-C1'-N1-C2
2	D	224	BRU	O4'-C1'-N1-C2

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	221	BRU	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	218	BRU	4	0
2	D	218	BRU	2	0
2	D	221	BRU	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	501	-	4,4,4	0.39	0	6,6,6	0.10	0
5	GOL	A	602	-	5,5,5	0.80	0	5,5,5	0.49	0
4	SO4	A	502	-	4,4,4	0.38	0	6,6,6	0.06	0
5	GOL	B	601	-	5,5,5	0.72	0	5,5,5	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	602	-	-	2/4/4/4	-
5	GOL	B	601	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	602	GOL	O1-C1-C2-C3
5	B	601	GOL	O1-C1-C2-C3
5	B	601	GOL	C1-C2-C3-O3
5	B	601	GOL	O2-C2-C3-O3
5	A	602	GOL	O1-C1-C2-O2
5	B	601	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	602	GOL	2	0
5	B	601	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	12/13 (92%)	0.62	1 (8%) 17 19	24, 45, 55, 58	0
1	E	12/13 (92%)	2.62	7 (58%) 0 0	29, 55, 97, 106	0
2	D	9/13 (69%)	1.76	3 (33%) 1 1	27, 46, 60, 60	0
2	F	9/13 (69%)	3.66	7 (77%) 0 0	35, 60, 113, 115	0
3	A	137/137 (100%)	0.09	5 (3%) 46 48	15, 26, 48, 55	1 (0%)
3	B	137/137 (100%)	1.21	45 (32%) 1 1	16, 38, 76, 85	1 (0%)
All	All	316/326 (96%)	0.84	68 (21%) 2 3	15, 32, 74, 115	2 (0%)

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	214	DG	7.7
1	E	213	DC	6.7
3	B	41	VAL	5.9
3	B	43	ASP	5.6
2	F	215	DG	5.5
1	E	212	DC	5.5
2	F	220	DA	5.1
3	B	35	VAL	4.9
3	B	40	ARG	4.7
3	B	42	ARG	4.6
2	F	216	DC	4.6
3	B	45	LYS	4.3
2	D	214	DG	4.1
3	B	90	VAL	4.0
3	B	34	HIS	4.0
2	F	219	DC	3.7
3	B	44	PHE	3.6
3	B	36	ALA	3.6
3	B	109	ALA	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	B	33	LYS	3.4
3	B	138	ALA	3.4
2	D	215	DG	3.3
3	B	104	TYR	3.3
2	F	226	DG	3.3
3	B	2	THR	3.3
3	B	98	GLN	3.2
3	B	31	VAL	3.2
3	B	54	ALA	3.2
1	E	202	DC	3.1
3	B	46	ILE	3.1
1	E	211	DG	3.1
3	B	39	LYS	3.0
2	D	220	DA	2.9
3	B	30	ALA	2.9
3	B	53	GLY	2.8
1	E	201	DC	2.8
1	E	203	DA	2.7
3	A	37	ASN	2.7
3	B	48	PRO	2.7
1	C	201	DC	2.6
1	E	210	DA	2.6
3	B	55	GLY	2.6
3	B	84[A]	ASN	2.6
3	B	38	GLY	2.6
3	B	51	ILE	2.6
3	B	97	PRO	2.6
3	B	61	TYR	2.6
3	B	113	ILE	2.5
3	B	100	PHE	2.4
3	B	91	GLN	2.4
3	B	49	THR	2.4
2	F	225	DG	2.4
3	B	71	GLN	2.4
3	A	108	GLU	2.4
3	B	32	ARG	2.3
3	B	108	GLU	2.3
3	B	52	LEU	2.3
3	A	40	ARG	2.2
3	B	107	HIS	2.2
3	B	93	ILE	2.1
3	A	45	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	98	GLN	2.1
3	B	47	SER	2.1
3	B	58	THR	2.1
3	B	89	THR	2.1
3	B	99	GLU	2.1
3	B	117	ARG	2.1
3	B	125	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BRU	F	217	20/21	0.59	0.32	104,107,112,119	0
2	BRU	F	218	20/21	0.67	0.28	95,101,104,112	0
2	BRU	F	221	20/21	0.75	0.22	50,60,69,89	0
2	BRU	D	217	20/21	0.78	0.20	55,60,65,85	0
2	BRU	D	221	20/21	0.86	0.15	33,41,52,77	0
2	BRU	D	218	20/21	0.86	0.16	55,58,63,80	0
2	BRU	F	224	20/21	0.88	0.16	43,52,64,82	0
2	BRU	D	224	20/21	0.90	0.16	31,44,59,78	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	B	601	6/6	0.82	0.23	31,33,36,40	0
5	GOL	A	602	6/6	0.86	0.20	55,57,57,59	0
4	SO4	A	502	5/5	0.94	0.14	71,71,72,73	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	A	501	5/5	0.97	0.09	39,40,41,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.