



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:54 PM UTC

PDB ID : 4FCB / pdb\_00004fcb  
Title : Potent and Selective Phosphodiesterase 10A Inhibitors  
Authors : Parris, K.D.  
Deposited on : 2012-05-24  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	345	 4% 79% 12% • 7%
1	B	345	 6% 76% 14% 10%

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5290 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	322	2623	1676	445	477	25	0	2	0
1	B	310	2521	1606	430	460	25	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	435	GLY	-	expression tag	UNP Q9Y233
A	436	SER	-	expression tag	UNP Q9Y233
A	437	HIS	-	expression tag	UNP Q9Y233
A	438	MET	-	expression tag	UNP Q9Y233
B	435	GLY	-	expression tag	UNP Q9Y233
B	436	SER	-	expression tag	UNP Q9Y233
B	437	HIS	-	expression tag	UNP Q9Y233
B	438	MET	-	expression tag	UNP Q9Y233

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0

- Molecule 4 is 3,4-dimethyl-1-propyl-7-(quinolin-2-ylmethoxy)imidazo[1,5-a]quinoxaline (CCD ID: 0T7) (formula: C<sub>25</sub>H<sub>24</sub>N<sub>4</sub>O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 30 25 4 1	0	0
4	B	1	Total C N O 30 25 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	57	Total O 57 57	0	0
5	B	25	Total O 25 25	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.15Å 81.17Å 157.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.17 – 2.10 41.17 – 2.10	Depositor EDS
% Data completeness (in resolution range)	88.8 (41.17-2.10) 89.0 (41.17-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.27 (at 2.10Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.244 , 0.284 0.239 , 0.280	Depositor DCC
$R_{free}$ test set	1720 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.6	Xtrriage
Anisotropy	0.384	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 32.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5290	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0T7, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.51	2/2687 (0.1%)	0.89	4/3638 (0.1%)
1	B	0.49	0/2581	0.86	1/3492 (0.0%)
All	All	0.50	2/5268 (0.0%)	0.87	5/7130 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	725	ILE	CA-CB	6.80	1.57	1.54
1	A	712	VAL	CA-CB	5.56	1.56	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	580	THR	N-CA-C	-9.87	89.77	110.80
1	A	579	SER	N-CA-C	7.79	122.42	111.52
1	A	570	HIS	CA-C-N	6.94	126.55	119.19
1	A	570	HIS	C-N-CA	6.94	126.55	119.19
1	B	736	PRO	N-CA-C	5.54	117.46	110.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	579	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2623	0	2586	27	0
1	B	2521	0	2482	26	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	30	0	24	2	0
4	B	30	0	24	2	0
5	A	57	0	0	2	0
5	B	25	0	0	0	0
All	All	5290	0	5116	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:803:OT7:H5	4:B:803:OT7:H8	1.75	0.66
1:A:648:ASN:HD22	1:A:648:ASN:C	2.04	0.66
1:B:540:ASP:O	1:B:544:LYS:HG2	1.98	0.63
1:B:450:MET:HE3	1:B:495:LEU:HD23	1.81	0.62
1:A:645:ASN:H	1:A:651:HIS:HD2	1.47	0.60
1:A:672:TRP:CZ2	1:A:676:LYS:HD3	2.37	0.59
1:A:560:PHE:HA	1:A:689:GLU:OE1	2.04	0.57
1:A:546:LEU:HD21	1:A:656:ILE:HG23	1.84	0.57
1:B:691:ASP:O	1:B:695:LYS:HG2	2.07	0.54
1:B:693:MET:HB3	1:B:698:ILE:HB	1.89	0.53
1:A:620:ILE:O	1:A:623:THR:OG1	2.24	0.52
1:A:665:LEU:O	1:A:668:VAL:HG22	2.10	0.52
1:B:631:ASN:HD22	1:B:654:ARG:HH11	1.57	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:590:VAL:HG12	1:B:594:GLN:HE21	1.74	0.52
1:A:496[B]:GLU:H	1:A:496[B]:GLU:CD	2.18	0.51
1:A:706:ARG:O	1:A:709:LYS:HG3	2.11	0.51
1:B:687:TRP:HB3	1:B:706:ARG:HB3	1.93	0.50
1:B:542:GLU:CD	1:B:652:ARG:HH21	2.19	0.50
1:B:617:ARG:O	1:B:621:ILE:HG12	2.13	0.49
1:B:667:SER:OG	1:B:678:THR:HG21	2.12	0.49
1:B:586:PHE:O	1:B:590:VAL:HG23	2.13	0.49
1:A:441:CYS:HA	1:A:507:LYS:HD2	1.95	0.49
1:A:450:MET:HE3	1:A:495:LEU:HD23	1.95	0.48
1:B:579:SER:HB2	1:B:582:GLU:HB2	1.96	0.47
1:A:608:GLU:OE1	5:A:950:HOH:O	2.20	0.47
1:A:473:GLU:HA	1:A:476:TRP:CE2	2.49	0.47
1:B:673:PRO:HA	1:B:676:LYS:HE2	1.95	0.47
1:A:494:GLU:HG2	1:A:496[B]:GLU:HG2	1.97	0.46
1:B:538:PHE:O	1:B:543:ARG:NH2	2.49	0.46
1:A:569:ASP:HA	1:A:570:HIS:O	2.14	0.46
1:A:449:LEU:HD13	1:A:477:PRO:HB2	1.97	0.46
1:A:546:LEU:HD21	1:A:656:ILE:CG2	2.45	0.46
4:A:803:OT7:H8	4:A:803:OT7:H7	1.98	0.46
1:B:683:TYR:CE2	4:B:803:OT7:H21	2.51	0.46
1:B:665:LEU:O	1:B:668:VAL:HG22	2.16	0.45
1:A:531:LEU:HD22	1:A:543:ARG:HG2	1.99	0.45
1:A:631:ASN:HD22	1:A:654:ARG:HH11	1.64	0.44
1:A:582:GLU:HA	1:A:585:HIS:HD2	1.82	0.44
1:B:690:GLY:HA3	1:B:704:MET:O	2.17	0.43
1:B:706:ARG:O	1:B:709:LYS:HG3	2.17	0.43
1:B:449:LEU:HD13	1:B:477:PRO:HB2	2.00	0.43
1:B:687:TRP:CH2	1:B:709:LYS:HG2	2.52	0.43
1:A:645:ASN:H	1:A:651:HIS:CD2	2.31	0.43
1:B:601:PHE:HB3	1:B:604:LEU:HD22	2.00	0.43
1:A:648:ASN:C	1:A:648:ASN:ND2	2.76	0.42
1:A:725:ILE:HB	1:A:726:PRO:HD3	2.02	0.42
1:A:575:LEU:HD22	1:A:576:TYR:CZ	2.55	0.42
4:A:803:OT7:H5	5:A:939:HOH:O	2.20	0.42
1:B:438:MET:HE1	1:B:588:GLN:OE1	2.20	0.42
1:A:492:CYS:SG	1:A:544:LYS:HE2	2.59	0.42
1:A:582:GLU:HA	1:A:585:HIS:CD2	2.54	0.41
1:B:645:ASN:ND2	1:B:647:ASN:H	2.18	0.41
1:B:683:TYR:HB3	1:B:687:TRP:CZ2	2.56	0.41
1:A:479:ILE:O	1:A:483:MET:HG3	2.20	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:631:ASN:HD22	1:B:654:ARG:NH1	2.19	0.40
1:B:720:TYR:HA	1:B:724:ALA:HB3	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	322/345 (93%)	308 (96%)	11 (3%)	3 (1%)	14	10
1	B	307/345 (89%)	294 (96%)	13 (4%)	0	100	100
All	All	629/690 (91%)	602 (96%)	24 (4%)	3 (0%)	24	22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	570	HIS
1	A	492	CYS
1	A	569	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/308 (95%)	285 (98%)	7 (2%)	43	49

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	282/308 (92%)	275 (98%)	7 (2%)	42	48
All	All	574/616 (93%)	560 (98%)	14 (2%)	44	49

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	570	HIS
1	A	575	LEU
1	A	580	THR
1	A	648	ASN
1	A	699	GLN
1	A	742	LEU
1	A	743	LYS
1	B	453[A]	THR
1	B	453[B]	THR
1	B	494	GLU
1	B	579	SER
1	B	633	LYS
1	B	711	GLU
1	B	750	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	535	HIS
1	A	584	HIS
1	A	631	ASN
1	A	645	ASN
1	A	648	ASN
1	A	651	HIS
1	A	680	ASN
1	A	714	GLN
1	B	594	GLN
1	B	599	ASN
1	B	611	GLN
1	B	631	ASN
1	B	634	GLN
1	B	645	ASN
1	B	680	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	OT7	B	803	-	33,34,34	0.99	1 (3%)	41,49,49	1.20	3 (7%)
4	OT7	A	803	-	33,34,34	0.90	1 (3%)	41,49,49	1.21	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OT7	B	803	-	-	4/8/8/8	0/5/5/5
4	OT7	A	803	-	-	1/8/8/8	0/5/5/5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	803	OT7	C16-N4	2.64	1.36	1.32

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	803	0T7	C16-N4	2.50	1.36	1.32

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	803	0T7	C14-C13-N3	-4.08	115.57	119.62
4	B	803	0T7	C3-C4-N1	-3.45	119.44	125.36
4	B	803	0T7	C14-C13-N3	-3.08	116.56	119.62
4	A	803	0T7	C3-C4-N1	-2.81	120.54	125.36
4	B	803	0T7	N2-C4-N1	-2.09	110.57	113.23

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	803	0T7	C1-C2-C3-C4
4	B	803	0T7	C9-C10-O1-C15
4	B	803	0T7	C11-C10-O1-C15
4	B	803	0T7	C2-C3-C4-N2
4	B	803	0T7	C2-C3-C4-N1

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	803	0T7	2	0
4	A	803	0T7	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	322/345 (93%)	0.55	15 (4%) 36 38	24, 48, 71, 76	2 (0%)
1	B	310/345 (89%)	0.72	21 (6%) 23 25	29, 53, 72, 84	1 (0%)
All	All	632/690 (91%)	0.63	36 (5%) 29 31	24, 51, 71, 84	3 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	698	ILE	5.5
1	B	564	TYR	3.7
1	A	571	PRO	3.5
1	B	578	THR	3.5
1	B	562	ASN	3.3
1	A	492	CYS	3.2
1	A	570	HIS	3.2
1	B	600	ILE	3.1
1	A	568	PHE	3.0
1	B	560	PHE	2.9
1	A	495	LEU	2.8
1	A	493	PHE	2.8
1	B	703	MET	2.7
1	A	580	THR	2.7
1	B	700	PRO	2.6
1	B	446	TRP	2.6
1	B	437	HIS	2.6
1	B	697	GLY	2.5
1	A	579	SER	2.4
1	A	698	ILE	2.4
1	B	704	MET	2.3
1	B	442	THR	2.3
1	A	448	GLY	2.3
1	A	446	TRP	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	492	CYS	2.2
1	A	462	ILE	2.2
1	B	594	GLN	2.2
1	A	488	CYS	2.2
1	B	495	LEU	2.2
1	B	604	LEU	2.2
1	A	452	PHE	2.1
1	B	701	ILE	2.1
1	B	558	ARG	2.1
1	B	440	ILE	2.1
1	A	494	GLU	2.1
1	B	601	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	OT7	B	803	30/30	0.80	0.16	61,63,65,65	0
4	OT7	A	803	30/30	0.88	0.11	49,51,54,54	0
3	MG	B	802	1/1	0.97	0.04	40,40,40,40	0
2	ZN	A	801	1/1	0.99	0.03	42,42,42,42	0
2	ZN	B	801	1/1	0.99	0.05	46,46,46,46	0
3	MG	A	802	1/1	0.99	0.06	33,33,33,33	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 0T7 B 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 0T7 A 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.