



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 08:26 AM UTC

PDB ID : 5FCS / pdb_00005fcs
Title : Diabody
Authors : Mosyak, L.; Root, A.
Deposited on : 2015-12-15
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	247	 6% 83% 10% 6%
2	L	256	 4% 86% 8% 6%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4119 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Diabody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	233	1724	1079	294	342	9	0	0	0

- Molecule 2 is a protein called Diabody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	240	1805	1133	308	357	7	0	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	L	1	5	4	1	0	0
3	L	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	L	1	Total	O	S	0	0
			5	4	1		
3	L	1	Total	O	S	0	0
			5	4	1		
3	L	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	259	Total	O	0	0
			259	259		
4	L	306	Total	O	0	0
			306	306		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Diabody



- Molecule 2: Diabody



4 Data and refinement statistics i

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	142.81Å 142.81Å 62.69Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.43 – 2.01 29.43 – 2.01	Depositor EDS
% Data completeness (in resolution range)	96.5 (29.43-2.01) 96.8 (29.43-2.01)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 2.01Å)	Xtrriage
Refinement program	BUSTER 2.11.1	Depositor
R, R_{free}	0.176 , 0.205 0.178 , 0.203	Depositor DCC
R_{free} test set	2374 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	26.8	Xtrriage
Anisotropy	0.181	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.033 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4119	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.85	0/1761	1.13	7/2399 (0.3%)
2	L	0.83	0/1847	1.04	1/2509 (0.0%)
All	All	0.84	0/3608	1.08	8/4908 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	219	THR	N-CA-C	-7.64	102.92	111.71
1	H	217	ILE	CB-CA-C	-6.02	101.66	110.81
1	H	224	ASP	N-CA-C	5.99	119.06	111.69
1	H	150	TYR	N-CA-C	5.89	119.65	110.17
2	L	188	PHE	CA-CB-CG	5.78	119.58	113.80
1	H	82	ALA	CA-C-N	5.58	128.31	120.28
1	H	82	ALA	C-N-CA	5.58	128.31	120.28
1	H	222	ALA	N-CA-C	-5.01	101.70	109.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1724	0	1674	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	L	1805	0	1730	12	0
3	L	25	0	0	0	0
4	H	259	0	0	2	0
4	L	306	0	0	1	1
All	All	4119	0	3404	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:120:VAL:HG22	4:H:411:HOH:O	1.83	0.77
1:H:225:ILE:HD12	4:H:411:HOH:O	2.00	0.60
1:H:124:GLN:HE21	1:H:227:GLY:HA3	1.68	0.58
2:L:6:GLN:HE21	2:L:102:GLY:HA3	1.71	0.56
1:H:218:ASP:HB3	1:H:220:ALA:H	1.71	0.56
1:H:22:CYS:HB3	1:H:73:ALA:HB3	1.92	0.52
1:H:239:CYS:HG	2:L:246:CYS:HG	1.54	0.51
2:L:80:GLN:HE22	2:L:119:GLU:H	1.56	0.51
1:H:218:ASP:HB2	1:H:222:ALA:H	1.75	0.50
1:H:124:GLN:HE22	1:H:213:TYR:HA	1.76	0.50
1:H:6:GLN:HE21	1:H:101:GLY:HA3	1.77	0.49
1:H:30:THR:H	1:H:33:ASN:HD22	1.61	0.48
2:L:203:MET:HE2	2:L:206:LEU:HD21	1.95	0.48
2:L:40:LEU:HG	2:L:235:GLN:HG3	1.97	0.47
2:L:80:GLN:NE2	2:L:119:GLU:H	2.14	0.46
2:L:62:ARG:HB2	2:L:77:THR:O	2.14	0.46
1:H:41:LYS:HB2	1:H:44:GLN:HG3	1.99	0.44
2:L:106:LYS:HE3	4:L:566:HOH:O	2.19	0.42
2:L:38:GLN:HB2	2:L:48:LEU:HD11	2.01	0.42
2:L:170:ARG:HG3	2:L:177:ALA:HB3	2.02	0.42
1:H:185:ARG:NH2	1:H:208:ASP:OD2	2.44	0.42
1:H:154:TRP:CE2	1:H:199:MET:HB2	2.55	0.41
1:H:217:ILE:HG22	1:H:218:ASP:N	2.33	0.41
2:L:152:MET:HB3	2:L:199:LEU:HD22	2.03	0.40
1:H:127:ALA:O	2:L:244:GLY:HA2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:L:518:HOH:O	4:L:518:HOH:O[5_555]	1.62	0.58

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	229/247 (93%)	225 (98%)	4 (2%)	0	100	100
2	L	236/256 (92%)	232 (98%)	4 (2%)	0	100	100
All	All	465/503 (92%)	457 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	180/187 (96%)	180 (100%)	0	100	100
2	L	194/204 (95%)	194 (100%)	0	100	100
All	All	374/391 (96%)	374 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	6	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	33	ASN
1	H	36	ASN
1	H	40	GLN
1	H	121	GLN
1	H	124	GLN
1	H	157	GLN
1	H	161	GLN
1	H	183	GLN
2	L	6	GLN
2	L	39	GLN
2	L	53	ASN
2	L	80	GLN
2	L	157	GLN
2	L	202	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	L	304	-	4,4,4	0.53	0	6,6,6	0.25	0
3	SO4	L	302	-	4,4,4	0.47	0	6,6,6	0.16	0
3	SO4	L	301	-	4,4,4	0.39	0	6,6,6	0.36	0
3	SO4	L	305	-	4,4,4	0.49	0	6,6,6	0.20	0
3	SO4	L	303	-	4,4,4	0.60	0	6,6,6	0.35	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	233/247 (94%)	-0.00	14 (6%) 27 26	17, 28, 56, 78	0
2	L	240/256 (93%)	-0.22	9 (3%) 44 43	16, 26, 47, 61	0
All	All	473/503 (94%)	-0.11	23 (4%) 35 34	16, 27, 52, 78	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	110	GLY	5.3
1	H	172	TYR	4.1
2	L	116	GLY	3.8
1	H	150	TYR	3.6
1	H	147	PHE	3.6
1	H	239	CYS	3.4
1	H	148	THR	3.1
2	L	246	CYS	3.1
1	H	219	THR	3.0
2	L	96	LEU	2.9
1	H	218	ASP	2.8
1	H	146	THR	2.8
1	H	171	ALA	2.8
2	L	118	GLY	2.7
2	L	110	LEU	2.7
1	H	221	SER	2.5
1	H	145	TYR	2.3
2	L	117	GLY	2.2
2	L	161	LYS	2.2
1	H	236	SER	2.2
1	H	173	ASN	2.2
2	L	42	GLY	2.1
2	L	97	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	L	304	5/5	0.59	0.22	79,82,83,83	0
3	SO4	L	305	5/5	0.83	0.12	54,57,59,63	0
3	SO4	L	302	5/5	0.88	0.15	50,51,54,58	0
3	SO4	L	303	5/5	0.90	0.15	39,39,42,45	0
3	SO4	L	301	5/5	0.95	0.14	51,52,54,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.