



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 05:02 AM UTC

PDB ID : 7FCL / pdb_00007fcl
Title : Zebrafish SIGIRR TIR domain
Authors : Wang, X.; Zhou, J.
Deposited on : 2021-07-15
Resolution : 3.04 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3685 (3.08-3.00)
Clashscore	190562	4007 (3.08-3.00)
Ramachandran outliers	187476	3834 (3.08-3.00)
RSRZ outliers	180081	3684 (3.08-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	155	 65% 29% 6%
1	B	155	 72% 23% 6%
1	C	155	 73% 21% 6%

2 Entry composition i

There is only 1 type of molecule in this entry. The entry contains 3621 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SIGIRR protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	146	1207	773	211	215	8	0	0	0
1	B	146	1207	773	211	215	8	0	0	0
1	C	146	1207	773	211	215	8	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	179	MET	-	initiating methionine	UNP K9K3G6
A	218	ASN	ASP	engineered mutation	UNP K9K3G6
A	328	HIS	-	expression tag	UNP K9K3G6
A	329	HIS	-	expression tag	UNP K9K3G6
A	330	HIS	-	expression tag	UNP K9K3G6
A	331	HIS	-	expression tag	UNP K9K3G6
A	332	HIS	-	expression tag	UNP K9K3G6
A	333	HIS	-	expression tag	UNP K9K3G6
B	179	MET	-	initiating methionine	UNP K9K3G6
B	218	ASN	ASP	engineered mutation	UNP K9K3G6
B	328	HIS	-	expression tag	UNP K9K3G6
B	329	HIS	-	expression tag	UNP K9K3G6
B	330	HIS	-	expression tag	UNP K9K3G6
B	331	HIS	-	expression tag	UNP K9K3G6
B	332	HIS	-	expression tag	UNP K9K3G6
B	333	HIS	-	expression tag	UNP K9K3G6
C	179	MET	-	initiating methionine	UNP K9K3G6
C	218	ASN	ASP	engineered mutation	UNP K9K3G6
C	328	HIS	-	expression tag	UNP K9K3G6
C	329	HIS	-	expression tag	UNP K9K3G6
C	330	HIS	-	expression tag	UNP K9K3G6
C	331	HIS	-	expression tag	UNP K9K3G6
C	332	HIS	-	expression tag	UNP K9K3G6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	333	HIS	-	expression tag	UNP K9K3G6

4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	84.00Å 145.36Å 75.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.56 – 3.04 33.56 – 3.04	Depositor EDS
% Data completeness (in resolution range)	99.2 (33.56-3.04) 97.4 (33.56-3.04)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.28 (at 3.06Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.239 , 0.273 0.246 , 0.275	Depositor DCC
R_{free} test set	478 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å ²)	75.3	Xtrriage
Anisotropy	0.360	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 80.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.438 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.416 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3621	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/1237	0.86	0/1675
1	B	0.55	0/1237	0.82	0/1675
1	C	0.56	0/1237	0.88	1/1675 (0.1%)
All	All	0.56	0/3711	0.85	1/5025 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	237	ARG	CB-CA-C	-5.60	101.95	111.02

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1207	0	1202	31	0
1	B	1207	0	1202	23	0
1	C	1207	0	1203	22	1
All	All	3621	0	3607	73	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 73 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:291:GLN:O	1:B:295:GLN:OE1	1.61	1.17
1:B:291:GLN:C	1:B:295:GLN:OE1	2.29	0.74
1:B:190:ASN:O	1:B:190:ASN:ND2	2.24	0.68
1:A:225:ALA:O	1:A:261:GLN:NE2	2.27	0.67
1:B:225:ALA:O	1:B:261:GLN:NE2	2.30	0.64

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:299:CYS:CB	1:C:299:CYS:SG[3_556]	1.81	0.39

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/155 (93%)	138 (96%)	5 (4%)	1 (1%)	18	49
1	B	144/155 (93%)	138 (96%)	4 (3%)	2 (1%)	9	33
1	C	144/155 (93%)	138 (96%)	4 (3%)	2 (1%)	9	33
All	All	432/465 (93%)	414 (96%)	13 (3%)	5 (1%)	10	37

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	310	MET
1	B	310	MET
1	A	310	MET
1	B	271	ARG
1	C	271	ARG

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	146/155 (94%)	-1.27	0 100 100	44, 76, 94, 101	0
1	B	146/155 (94%)	-1.25	0 100 100	44, 75, 96, 102	0
1	C	146/155 (94%)	-1.23	0 100 100	43, 75, 94, 102	0
All	All	438/465 (94%)	-1.25	0 100 100	43, 75, 96, 102	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.