



wwPDB EM Validation Summary Report ⓘ

Mar 24, 2026 – 05:37 AM UTC

PDB ID : 8FCU / pdb_00008fcu
EMDB ID : EMD-28993
Title : Cryo-EM structure of Cascade-DNA-TniQ-TnsC complex in type I-B CAST system
Authors : Chang, L.; Wang, S.
Deposited on : 2022-12-01
Resolution : 3.19 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	14455 (2.69 - 3.69)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	212	68% (green), 30% (yellow), 2% (orange), 0% (red), 1% (grey)
2	B	220	60% (green), 36% (yellow), 2% (orange), 2% (red), 2% (grey)
3	C	323	60% (green), 28% (yellow), 12% (grey)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	323	 59% 29% 12%
3	E	323	 60% 27% 11%
3	F	323	 60% 28% 12%
3	G	323	 61% 28% 12%
3	H	323	 59% 29% 12%
4	I	534	 55% 37% 7%
5	J	138	 44% 35% 21%
5	K	138	 46% 33% 21%
5	L	138	 56% 23% 21%
6	M	71	 15% 56% 15% 11%
7	N	85	 22% 52% 26%
8	O	85	 6% 14% 21% 65%
9	P	329	 67% 31% 2% 2%
10	Q	383	 8% 59% 21% 19%

2 Entry composition i

There are 12 unique types of molecules in this entry. The entry contains 32692 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type I-B CRISPR-associated protein Cas5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	212	1745	1116	313	305	11	0	0

- Molecule 2 is a protein called Type I-B CRISPR-associated protein Cas6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	213	1702	1084	297	316	5	0	0

- Molecule 3 is a protein called Type I-B CRISPR-associated protein Cas7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	285	2323	1490	393	430	10	0	0
3	D	285	2323	1490	393	430	10	0	0
3	E	286	2330	1494	394	432	10	0	0
3	F	285	2323	1490	393	430	10	0	0
3	G	285	2323	1490	393	430	10	0	0
3	H	285	2323	1490	393	430	10	0	0

- Molecule 4 is a protein called Type I-MYXAN CRISPR-associated Cas8a1/Cmx1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	I	494	4147	2689	698	748	12	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	-7	MET	-	initiating methionine	UNP A0A235IGR9
I	-6	HIS	-	expression tag	UNP A0A235IGR9
I	-5	HIS	-	expression tag	UNP A0A235IGR9
I	-4	HIS	-	expression tag	UNP A0A235IGR9
I	-3	HIS	-	expression tag	UNP A0A235IGR9
I	-2	HIS	-	expression tag	UNP A0A235IGR9
I	-1	HIS	-	expression tag	UNP A0A235IGR9
I	0	HIS	-	expression tag	UNP A0A235IGR9
I	1	HIS	-	expression tag	UNP A0A235IGR9
I	2	ILE	-	expression tag	UNP A0A235IGR9

- Molecule 5 is a protein called Type I-B CRISPR-associated protein Cas11.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	J	109	Total	C	N	O	S	0	0
			917	598	150	167	2		
5	K	109	Total	C	N	O	S	0	0
			917	598	150	167	2		
5	L	109	Total	C	N	O	S	0	0
			917	598	150	167	2		

- Molecule 6 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	M	63	Total	C	N	O	P	0	0
			1342	600	236	443	63		

- Molecule 7 is a DNA chain called Target DNA strand.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	N	63	Total	C	N	O	P	0	0
			1282	616	221	382	63		

- Molecule 8 is a DNA chain called Non-target DNA strand.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	O	30	Total	C	N	O	P	0	0
			620	296	115	179	30		


- Molecule 9 is a protein called TniQ.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	P	327	2618	1688	438	473	19	0	0

- Molecule 10 is a protein called TnsC.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	Q	310	2508	1605	435	460	8	0	0

- Molecule 11 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
11	Q	1	31	10	5	13	3	0


- Molecule 12 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
			Total	Mg	
12	Q	1	1	1	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Type I-B CRISPR-associated protein Cas5



- Molecule 2: Type I-B CRISPR-associated protein Cas6





- Molecule 3: Type I-B CRISPR-associated protein Cas7








• Molecule 3: Type I-B CRISPR-associated protein Cas7




• Molecule 3: Type I-B CRISPR-associated protein Cas7




• Molecule 3: Type I-B CRISPR-associated protein Cas7



• Molecule 3: Type I-B CRISPR-associated protein Cas7







• Molecule 8: Non-target DNA strand



• Molecule 9: TniQ



• Molecule 10: TnsC



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	96534	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	54	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	35.722	Depositor
Minimum map value	-15.332	Depositor
Average map value	0.069	Depositor
Map value standard deviation	0.876	Depositor
Recommended contour level	2.6	Depositor
Map size (\AA)	404.73602, 404.73602, 404.73602	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.054, 1.054, 1.054	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.18	0/1788	0.40	0/2422
2	B	0.17	0/1735	0.47	0/2341
3	C	0.13	0/2384	0.36	0/3220
3	D	0.17	0/2384	0.39	0/3220
3	E	0.12	0/2391	0.34	0/3231
3	F	0.15	0/2384	0.39	0/3220
3	G	0.12	0/2384	0.33	0/3220
3	H	0.14	0/2384	0.36	0/3220
4	I	0.19	0/4249	0.41	0/5741
5	J	0.17	0/938	0.45	0/1258
5	K	0.16	0/938	0.43	0/1258
5	L	0.13	0/938	0.32	0/1258
6	M	0.17	0/1500	0.41	1/2335 (0.0%)
7	N	0.22	0/1434	0.46	0/2209
8	O	0.18	0/695	0.39	0/1069
9	P	0.13	0/2673	0.35	0/3607
10	Q	0.12	0/2562	0.31	0/3468
All	All	0.16	0/33761	0.38	1/46297 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	M	3	G	C4'-C3'-O3'	5.21	117.22	109.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1745	0	1772	56	0
2	B	1702	0	1735	57	0
3	C	2323	0	2249	69	0
3	D	2323	0	2249	84	0
3	E	2330	0	2255	75	0
3	F	2323	0	2249	84	0
3	G	2323	0	2249	78	0
3	H	2323	0	2249	82	0
4	I	4147	0	4143	201	0
5	J	917	0	925	39	0
5	K	917	0	925	43	0
5	L	917	0	925	30	0
6	M	1342	0	675	64	0
7	N	1282	0	716	60	0
8	O	620	0	341	20	0
9	P	2618	0	2687	81	0
10	Q	2508	0	2514	63	0
11	Q	31	0	12	1	0
12	Q	1	0	0	0	0
All	All	32692	0	30870	1025	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 1025 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:I:366:LEU:HD13	4:I:379:LEU:HD12	1.35	1.08
4:I:366:LEU:HD22	4:I:379:LEU:HB2	1.48	0.96
1:A:165:ARG:HA	1:A:190:TYR:HA	1.46	0.93
5:J:49:ARG:HH12	5:K:71:ARG:HG3	1.34	0.93
4:I:137:VAL:HG13	4:I:335:LYS:HA	1.51	0.90

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/212 (99%)	203 (97%)	7 (3%)	0	100	100
2	B	211/220 (96%)	201 (95%)	9 (4%)	1 (0%)	24	59
3	C	281/323 (87%)	249 (89%)	30 (11%)	2 (1%)	18	52
3	D	281/323 (87%)	246 (88%)	32 (11%)	3 (1%)	11	43
3	E	282/323 (87%)	258 (92%)	22 (8%)	2 (1%)	18	52
3	F	281/323 (87%)	243 (86%)	36 (13%)	2 (1%)	18	52
3	G	281/323 (87%)	263 (94%)	16 (6%)	2 (1%)	18	52
3	H	281/323 (87%)	257 (92%)	22 (8%)	2 (1%)	18	52
4	I	492/534 (92%)	478 (97%)	12 (2%)	2 (0%)	30	62
5	J	107/138 (78%)	97 (91%)	10 (9%)	0	100	100
5	K	107/138 (78%)	98 (92%)	8 (8%)	1 (1%)	14	47
5	L	107/138 (78%)	105 (98%)	2 (2%)	0	100	100
9	P	325/329 (99%)	312 (96%)	13 (4%)	0	100	100
10	Q	306/383 (80%)	291 (95%)	15 (5%)	0	100	100
All	All	3552/4030 (88%)	3301 (93%)	234 (7%)	17 (0%)	26	59

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	D	136	ARG
3	E	247	PRO
3	F	247	PRO
4	I	334	TYR
3	G	267	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/193 (100%)	189 (98%)	4 (2%)	47	71
2	B	187/193 (97%)	184 (98%)	3 (2%)	55	75
3	C	250/287 (87%)	250 (100%)	0	100	100
3	D	250/287 (87%)	249 (100%)	1 (0%)	84	86
3	E	251/287 (88%)	247 (98%)	4 (2%)	55	75
3	F	250/287 (87%)	249 (100%)	1 (0%)	84	86
3	G	250/287 (87%)	250 (100%)	0	100	100
3	H	250/287 (87%)	248 (99%)	2 (1%)	73	82
4	I	449/489 (92%)	445 (99%)	4 (1%)	70	81
5	J	98/126 (78%)	98 (100%)	0	100	100
5	K	98/126 (78%)	98 (100%)	0	100	100
5	L	98/126 (78%)	98 (100%)	0	100	100
9	P	292/294 (99%)	290 (99%)	2 (1%)	76	83
10	Q	269/332 (81%)	267 (99%)	2 (1%)	76	83
All	All	3185/3601 (88%)	3162 (99%)	23 (1%)	73	83

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	H	155	VAL
4	I	331	ILE
4	I	116	LEU
4	I	358	TRP
2	B	171	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	42	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
10	Q	328	GLN
3	H	84	HIS
5	L	53	ASN
3	H	42	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
6	M	62/71 (87%)	24 (38%)	3 (4%)

5 of 24 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
6	M	2	U
6	M	3	G
6	M	4	C
6	M	8	A
6	M	9	G

All (3) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
6	M	3	G
6	M	14	G
6	M	26	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	ATP	Q	600	12	32,33,33	0.27	0	48,52,52	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	ATP	Q	600	12	-	5/22/38/38	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	Q	600	ATP	C4'-C5'-O5'-PA
11	Q	600	ATP	C5'-O5'-PA-O1A
11	Q	600	ATP	PB-O3B-PG-O1G
11	Q	600	ATP	PG-O3B-PB-O2B
11	Q	600	ATP	PG-O3B-PB-O1B


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	Q	600	ATP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-28993. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192



Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 201




Y Index: 196




Z Index: 141


6.3.2 Raw map



X Index: 202



Y Index: 196




Z Index: 141


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z

6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 2.6. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 569 nm³; this corresponds to an approximate mass of 514 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.313 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.313 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.19	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	6.25	10.72	7.23

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.25 differs from the reported value 3.19 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-28993 and PDB model 8FCU. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 2.6 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (2.6).





































9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 95% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (2.6) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9530	 0.3930
A	 0.9830	 0.4680
B	 0.9850	 0.3870
C	 0.9910	 0.4610
D	 0.9900	 0.4810
E	 0.9900	 0.4820
F	 0.9850	 0.4810
G	 0.9900	 0.4860
H	 0.9880	 0.4750
I	 0.9270	 0.3370
J	 0.9770	 0.4090
K	 0.9900	 0.4340
L	 0.9830	 0.4390
M	 0.9930	 0.4010
N	 0.9160	 0.3380
O	 0.7790	 0.1670
P	 0.9380	 0.3200
Q	 0.7770	 0.0930

