



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 06:10 AM EDT

PDB ID : 2FDP / pdb_00002fdp
Title : Crystal structure of beta-secretase complexed with an amino-ethylene inhibitor
Authors : Yang, W.; Lu, W.; Lu, Y.; Zhong, M.; Sun, J.; Thomas, A.E.; Wilkinson, J.M.; Fucini, R.V.; Lam, M.; Randal, M.; Shi, X.P.; Jacobs, J.W.; McDowell, R.S.; Gordon, E.M.; Ballinger, M.D.
Deposited on : 2005-12-14
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	 4% 81% 15% •
1	B	388	 4% 83% 12% • 5%
1	C	388	 4% 82% 14% •

2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 9008 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	373	Total	C	N	O	S	0	0	0
			2940	1882	489	555	14			
1	B	369	Total	C	N	O	S	0	0	0
			2908	1865	484	545	14			
1	C	374	Total	C	N	O	S	0	0	0
			2937	1881	489	553	14			

- Molecule 2 is N1-((2S,3S,5R)-3-AMINO-6-(4-FLUOROPHENYLAMINO)-5-METHYL-6-OXO-1-PHENYLHEXAN-2-YL)-N3,N3-DIPROPYLISOPHTHALAMIDE (CCD ID: FRP) (formula: C₃₃H₄₁FN₄O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
2	A	1	Total	C	F	N	O	0	0
			41	33	1	4	3		
2	B	1	Total	C	F	N	O	0	0
			41	33	1	4	3		


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
2	C	1	41	33	1	4	3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	36	Total	O	0	0
			36	36		
3	B	35	Total	O	0	0
			35	35		
3	C	29	Total	O	0	0
			29	29		



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.19Å 102.19Å 99.53Å 90.00° 103.21° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 20.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.9 (20.00-2.50) 98.7 (20.00-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.55 (at 2.50Å)	Xtrriage
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.225 , 0.254 0.244 , 0.267	Depositor DCC
R_{free} test set	2794 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	42.7	Xtrriage
Anisotropy	0.095	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 17.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9008	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FRP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/3014	0.76	1/4092 (0.0%)
1	B	0.36	0/2982	0.75	0/4049
1	C	0.35	0/3011	0.75	0/4089
All	All	0.35	0/9007	0.75	1/12230 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	178	GLY	N-CA-C	5.23	117.25	110.45

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2940	0	2849	15	0
1	B	2908	0	2826	11	0
1	C	2937	0	2841	15	0
2	A	41	0	41	1	0
2	B	41	0	41	1	0
2	C	41	0	41	2	0
3	A	36	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	35	0	0	0	0
3	C	29	0	0	0	0
All	All	9008	0	8639	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:MET:HE3	1:A:176:ILE:HD11	1.85	0.57
1:A:257:PHE:HD2	1:A:268:VAL:HG21	1.71	0.56
1:C:110:ILE:HB	1:C:113:SER:HB3	1.89	0.55
1:B:235:ARG:HB2	1:B:332:VAL:HB	1.88	0.55
1:B:197:TRP:CG	1:B:198:TYR:H	2.25	0.54
1:A:197:TRP:CG	1:A:198:TYR:H	2.25	0.54
2:C:386:FRP:O3	2:C:386:FRP:H27	2.08	0.54
1:C:197:TRP:CG	1:C:198:TYR:H	2.26	0.54
1:A:235:ARG:HB2	1:A:332:VAL:HB	1.91	0.53
1:C:235:ARG:HB2	1:C:332:VAL:HB	1.91	0.52
1:C:174:MET:HE3	1:C:176:ILE:HD11	1.91	0.52
1:C:2:MET:HG2	1:C:90:GLY:HA2	1.92	0.52
1:B:301:LEU:HB3	1:B:302:PRO:HD2	1.93	0.51
1:A:194:ARG:HD2	1:A:202:ILE:HD11	1.95	0.49
1:C:276:PRO:HB2	1:C:279:ILE:HG12	1.95	0.48
1:B:2:MET:HG2	1:B:90:GLY:HA2	1.96	0.48
1:A:208:ILE:HG12	1:A:283:ILE:HG12	1.95	0.48
1:C:245:VAL:O	1:C:249:LYS:HB2	2.14	0.48
1:B:95:VAL:HG11	1:B:140:LEU:HA	1.97	0.47
1:A:95:VAL:HG11	1:A:140:LEU:HA	1.97	0.47
2:B:386:FRP:O3	2:B:386:FRP:H27	2.15	0.47
1:A:19:THR:HB	1:A:24:PRO:HB3	1.97	0.46
1:A:298:ILE:HG22	1:A:370:VAL:HG22	1.97	0.45
1:B:276:PRO:HB2	1:B:279:ILE:HG12	1.99	0.45
1:C:235:ARG:HB3	1:C:327:SER:HB2	1.96	0.45
1:B:50:ARG:O	1:B:116:GLU:HG2	2.16	0.45
2:A:386:FRP:H27	2:A:386:FRP:O3	2.17	0.44
1:A:302:PRO:HA	1:A:305:TYR:CE2	2.54	0.43
1:C:126:ILE:HG23	1:C:197:TRP:HB2	2.00	0.43
1:B:179:ILE:HG23	1:B:342:TYR:HE2	1.83	0.42
1:B:228:ASP:O	1:B:334:GLY:HA2	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:TRP:CD1	1:A:198:TYR:H	2.37	0.42
1:B:3:VAL:HG13	1:B:183:LEU:HD21	2.02	0.42
1:C:134:GLU:HA	1:C:135:PRO:HD3	1.94	0.42
1:A:199:TYR:HB3	1:A:352:ILE:HD11	2.02	0.42
1:A:307:ARG:HA	1:A:308:PRO:HD3	1.86	0.42
1:A:205:ARG:HB3	1:A:286:TYR:HB2	2.02	0.42
1:C:278:ASN:H	1:C:278:ASN:HD22	1.69	0.41
1:B:42:ALA:HB2	1:B:101:ALA:HB1	2.03	0.41
1:C:205:ARG:HB3	1:C:286:TYR:HB2	2.03	0.41
1:C:298:ILE:HD12	1:C:300:ILE:HD11	2.03	0.41
1:A:146:VAL:HA	1:A:147:PRO:HD3	1.96	0.40
1:C:71:TYR:HB3	2:C:386:FRP:H15	2.04	0.40
1:C:179:ILE:HG23	1:C:342:TYR:HE2	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	367/388 (95%)	356 (97%)	10 (3%)	1 (0%)	36	55
1	B	363/388 (94%)	353 (97%)	10 (3%)	0	100	100
1	C	368/388 (95%)	358 (97%)	9 (2%)	1 (0%)	36	55
All	All	1098/1164 (94%)	1067 (97%)	29 (3%)	2 (0%)	43	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	223	ASP
1	A	223	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/331 (96%)	292 (92%)	27 (8%)	10	22
1	B	315/331 (95%)	287 (91%)	28 (9%)	9	20
1	C	317/331 (96%)	287 (90%)	30 (10%)	8	17
All	All	951/993 (96%)	866 (91%)	85 (9%)	9	20

All (85) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48(P)	VAL
1	A	9	LYS
1	A	48	LEU
1	A	75	LYS
1	A	125	GLU
1	A	132	SER
1	A	142	LYS
1	A	169	SER
1	A	213	LEU
1	A	239	LYS
1	A	242	GLU
1	A	247	SER
1	A	256	LYS
1	A	259	ASP
1	A	266	GLN
1	A	267	LEU
1	A	271	GLN
1	A	274	THR
1	A	288	MET
1	A	292	THR
1	A	325	SER
1	A	361	VAL
1	A	362	HIS
1	A	364	GLU
1	A	375	VAL
1	A	379	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	380	GLU
1	B	46(P)	SER
1	B	48(P)	VAL
1	B	4	ASP
1	B	46	PRO
1	B	48	LEU
1	B	65	LYS
1	B	110	ILE
1	B	125	GLU
1	B	142	LYS
1	B	169	SER
1	B	195	ARG
1	B	207	GLU
1	B	208	ILE
1	B	218	LYS
1	B	239	LYS
1	B	242	GLU
1	B	247	SER
1	B	249	LYS
1	B	254	THR
1	B	266	GLN
1	B	267	LEU
1	B	279	ILE
1	B	301	LEU
1	B	304	GLN
1	B	321	LYS
1	B	339	GLU
1	B	363	ASP
1	B	367	THR
1	C	48(P)	VAL
1	C	48	LEU
1	C	59	THR
1	C	64	ARG
1	C	65	LYS
1	C	68	TYR
1	C	86	SER
1	C	142	LYS
1	C	195	ARG
1	C	200	GLU
1	C	213	LEU
1	C	214	LYS
1	C	218	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	239	LYS
1	C	246	LYS
1	C	247	SER
1	C	249	LYS
1	C	254	THR
1	C	256	LYS
1	C	265	GLU
1	C	267	LEU
1	C	284	SER
1	C	291	VAL
1	C	326	GLN
1	C	336	VAL
1	C	339	GLU
1	C	350	LYS
1	C	364	GLU
1	C	367	THR
1	C	375	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	HIS
1	A	98	ASN
1	A	233	ASN
1	A	385	ASN
1	B	12	GLN
1	B	89	HIS
1	B	266	GLN
1	B	278	ASN
1	C	278	ASN
1	C	304	GLN
1	C	326	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FRP	B	386	-	43,43,43	0.66	1 (2%)	52,57,57	0.97	3 (5%)
2	FRP	A	386	-	43,43,43	0.63	1 (2%)	52,57,57	0.97	3 (5%)
2	FRP	C	386	-	43,43,43	0.64	1 (2%)	52,57,57	0.94	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FRP	B	386	-	-	2/42/42/42	0/3/3/3
2	FRP	A	386	-	-	5/42/42/42	0/3/3/3
2	FRP	C	386	-	-	5/42/42/42	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	386	FRP	C26-N3	-2.51	1.36	1.41
2	B	386	FRP	C26-N3	-2.49	1.36	1.41
2	A	386	FRP	C26-N3	-2.31	1.37	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	386	FRP	C17-C15-N2	-2.95	105.89	110.08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	386	FRP	C17-C15-N2	-2.80	106.10	110.08
2	B	386	FRP	C17-C15-N2	-2.80	106.11	110.08
2	A	386	FRP	C3-C7-N1	2.56	121.86	118.66
2	C	386	FRP	C3-C7-N1	2.36	121.61	118.66
2	B	386	FRP	C22-C23-C24	-2.26	119.84	122.80
2	A	386	FRP	C22-C23-C24	-2.23	119.88	122.80
2	B	386	FRP	C3-C7-N1	2.11	121.29	118.66

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	386	FRP	N4-C16-C18-C19
2	C	386	FRP	N4-C16-C18-C19
2	C	386	FRP	C16-C18-C19-C21
2	A	386	FRP	C14-C10-C9-N1
2	C	386	FRP	C13-C11-C8-N1
2	A	386	FRP	C15-C17-C32-C33
2	A	386	FRP	C15-C17-C32-C31
2	A	386	FRP	C13-C11-C8-N1
2	A	386	FRP	N4-C16-C18-C19
2	B	386	FRP	C14-C10-C9-N1
2	C	386	FRP	C15-C16-C18-C19
2	C	386	FRP	C16-C18-C19-C20



There are no ring outliers.


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	386	FRP	1	0
2	A	386	FRP	1	0
2	C	386	FRP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	373/388 (96%)	0.05	16 (4%) 40 35	6, 19, 39, 53	0
1	B	369/388 (95%)	0.15	15 (4%) 41 37	5, 20, 34, 51	0
1	C	374/388 (96%)	0.07	15 (4%) 42 38	5, 20, 40, 58	0
All	All	1116/1164 (95%)	0.09	46 (4%) 41 37	5, 19, 38, 58	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	259	ASP	4.9
1	B	365	PHE	4.7
1	C	68	TYR	4.3
1	C	311	ASP	4.1
1	C	316	GLN	3.9
1	C	365	PHE	3.9
1	A	272	ALA	3.7
1	C	310	GLU	3.7
1	B	364	GLU	3.5
1	B	363	ASP	3.4
1	A	46(P)	SER	3.4
1	A	259	ASP	3.3
1	B	49	HIS	3.1
1	B	270	TRP	3.0
1	B	319	CYS	3.0
1	A	316	GLN	3.0
1	A	254	THR	2.9
1	B	258	PRO	2.7
1	C	157	ALA	2.7
1	B	265	GLU	2.6
1	C	258	PRO	2.6
1	C	361	VAL	2.6
1	C	315	SER	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	254	THR	2.5
1	B	256	LYS	2.4
1	B	267	LEU	2.4
1	A	270	TRP	2.4
1	A	55	GLN	2.4
1	B	278	ASN	2.4
1	C	223	ASP	2.3
1	A	380	GLU	2.3
1	C	7	ARG	2.3
1	B	309	VAL	2.3
1	C	49	HIS	2.2
1	A	315	SER	2.2
1	B	43	ALA	2.2
1	A	362	HIS	2.2
1	A	269	CYS	2.2
1	C	46(P)	SER	2.2
1	B	271	GLN	2.1
1	A	157	ALA	2.1
1	C	385	ASN	2.1
1	A	365	PHE	2.0
1	A	260	GLY	2.0
1	A	278	ASN	2.0
1	A	222	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
-----	------	-------	-----	-------	------	-----	----------------------------	-------




2	FRP	B	386	41/41	0.95	0.08	12,14,20,20	0
2	FRP	C	386	41/41	0.95	0.07	11,15,19,21	0
2	FRP	A	386	41/41	0.96	0.07	10,13,19,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around FRP C 386:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FRP A 386:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.