



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 04:30 AM UTC

PDB ID : 3FDY / pdb_00003fdy
Title : Pyranose 2-oxidase thermostable triple mutant, T169G/E542K/V546C
Authors : Tan, T.C.; Divne, C.
Deposited on : 2008-11-26
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	623	6% 84% 8% 8%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pyranose oxidase (Pyranose 2-oxidase).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	576	4538	2865	778	869	26	0	0	0

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	169	GLY	THR	engineered mutation	UNP Q7ZA32
A	542	LYS	GLU	engineered mutation	UNP Q7ZA32
A	546	CYS	VAL	engineered mutation	UNP Q7ZA32

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	470	Total	O	0	0
			470	470		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyranose oxidase (Pyranose 2-oxidase)



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	101.58Å 101.58Å 126.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.80 – 1.55 53.80 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.8 (53.80-1.55) 99.8 (53.80-1.55)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.16	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.186 , 0.219 (Not available) , 0.208	Depositor DCC
R_{free} test set	2899 reflections (3.01%)	wwPDB-VP
Wilson B-factor (Å ²)	9.6	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5073	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.25	4/4654 (0.1%)	1.14	10/6325 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	139	ARG	CD-NE	-6.03	1.37	1.46
1	A	450	HIS	C-O	-5.29	1.18	1.24
1	A	603	ILE	C-O	-5.25	1.18	1.24
1	A	224	ASN	N-CA	-5.03	1.40	1.46

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	139	ARG	NE-CZ-NH2	-10.85	109.43	119.20
1	A	139	ARG	CD-NE-CZ	7.00	134.19	124.40
1	A	139	ARG	CG-CD-NE	-6.30	98.15	112.00
1	A	139	ARG	CA-CB-CG	6.18	126.47	114.10
1	A	139	ARG	NE-CZ-NH1	6.09	127.59	121.50
1	A	180	ARG	CA-C-N	-5.68	114.12	120.14
1	A	180	ARG	C-N-CA	-5.68	114.12	120.14
1	A	616	SER	CA-C-N	-5.40	114.42	120.14
1	A	616	SER	C-N-CA	-5.40	114.42	120.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	GLN	N-CA-C	5.00	116.90	109.25

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	436	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4538	0	4387	26	0
2	A	53	0	29	0	0
3	A	12	0	12	0	0
4	A	470	0	0	4	0
All	All	5073	0	4428	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:201:LYS:HE2	1:A:205:TYR:OH	1.77	0.83
1:A:110:GLN:HE21	1:A:167:HIS:HD1	1.29	0.81
1:A:97:GLN:HB2	4:A:1401:HOH:O	1.97	0.64
1:A:432:GLU:H	1:A:432:GLU:CD	2.04	0.63
1:A:97:GLN:HG3	1:A:250:PHE:CE2	2.36	0.61
1:A:421:GLU:OE2	4:A:1389:HOH:O	2.17	0.58
1:A:100:ILE:HD13	1:A:100:ILE:O	2.04	0.57
1:A:457:GLY:O	1:A:461:GLN:HG3	2.06	0.55
1:A:47:TYR:O	1:A:313:ALA:HA	2.07	0.54
1:A:404:HIS:HE1	4:A:1254:HOH:O	1.94	0.51
1:A:81:ASP:OD1	1:A:81:ASP:C	2.54	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:402:ASN:ND2	4:A:1279:HOH:O	2.41	0.47
1:A:201:LYS:HE2	1:A:205:TYR:HH	1.79	0.47
1:A:449:ILE:HG12	1:A:471:TRP:CE3	2.50	0.46
1:A:618:PHE:C	1:A:618:PHE:CD1	2.93	0.46
1:A:100:ILE:HD13	1:A:453:ALA:HB2	1.99	0.45
1:A:618:PHE:C	1:A:618:PHE:HD1	2.24	0.45
1:A:358:GLU:HG2	1:A:544:GLY:HA2	1.98	0.44
1:A:165:SER:HA	1:A:168:TRP:CD1	2.53	0.44
1:A:167:HIS:CD2	1:A:167:HIS:C	2.97	0.42
1:A:346:PRO:HG2	1:A:350:PRO:HA	2.03	0.41
1:A:339:ARG:O	1:A:340:PRO:C	2.62	0.41
1:A:532:PHE:CE1	1:A:538:PRO:HG3	2.55	0.41
1:A:218:ARG:HG3	1:A:430:ASP:OD2	2.21	0.41
1:A:439:PHE:C	1:A:439:PHE:CD1	2.98	0.41
1:A:426:ILE:HA	1:A:427:PRO:HD3	1.98	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	517/623 (83%)	506 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	503/541 (93%)	491 (98%)	12 (2%)	43 15

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	ILE
1	A	100	ILE
1	A	206	PHE
1	A	312	LYS
1	A	347	GLU
1	A	385	THR
1	A	389	LEU
1	A	413	LYS
1	A	432	GLU
1	A	455	SER
1	A	496	ASN
1	A	542	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	ASN
1	A	257	ASN
1	A	263	GLN
1	A	276	ASN
1	A	341	ASN
1	A	419	HIS
1	A	460	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FAD	A	801	1	58,58,58	1.24	6 (10%)	85,89,89	2.31	32 (37%)
3	MES	A	901	-	12,12,12	1.94	1 (8%)	15,16,16	2.69	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	801	1	-	3/34/50/50	0/6/6/6
3	MES	A	901	-	-	2/6/14/14	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	901	MES	C8-S	-6.59	1.68	1.77
2	A	801	FAD	C8A-N9A	-3.59	1.31	1.37
2	A	801	FAD	O4B-C4B	-3.18	1.37	1.45
2	A	801	FAD	O3B-C3B	-2.20	1.37	1.43
2	A	801	FAD	C6A-N6A	-2.16	1.28	1.34
2	A	801	FAD	C2-N3	-2.09	1.34	1.39
2	A	801	FAD	C2B-C1B	-2.09	1.46	1.53

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	FAD	N3A-C2A-N1A	-6.79	118.30	128.58
2	A	801	FAD	C4A-N9A-C8A	6.60	112.67	105.74
3	A	901	MES	C5-N4-C3	6.37	122.56	108.84
2	A	801	FAD	O4'-C4'-C3'	5.10	121.18	109.25
2	A	801	FAD	N9A-C8A-N7A	-5.06	106.76	113.94
3	A	901	MES	O3S-S-C8	-5.00	96.22	106.00
2	A	801	FAD	O4B-C1B-N9A	4.86	117.42	108.09
2	A	801	FAD	C2B-C1B-N9A	4.66	124.89	113.30
2	A	801	FAD	O2B-C2B-C3B	4.37	125.83	111.82
2	A	801	FAD	C5'-C4'-C3'	-3.95	104.76	112.22
3	A	901	MES	C7-N4-C5	3.95	121.76	111.24
2	A	801	FAD	O3B-C3B-C4B	3.54	121.25	111.08
2	A	801	FAD	C5A-N7A-C8A	3.49	108.94	103.45
2	A	801	FAD	C4-C4X-C10	-3.35	111.19	116.93
2	A	801	FAD	N3A-C4A-N9A	3.26	132.71	127.17
2	A	801	FAD	C2A-N1A-C6A	3.24	124.05	118.73
2	A	801	FAD	O4-C4-C4X	-3.23	118.00	126.53
2	A	801	FAD	C2A-N3A-C4A	3.11	119.44	111.83
2	A	801	FAD	C4'-C3'-C2'	-2.91	108.72	113.57
2	A	801	FAD	C4-C4X-N5	2.87	122.17	118.21
2	A	801	FAD	O2B-C2B-C1B	2.73	119.52	110.10
2	A	801	FAD	O4B-C4B-C5B	2.73	118.07	109.33
3	A	901	MES	O1S-S-C8	-2.66	102.71	106.73
2	A	801	FAD	O4B-C4B-C3B	2.62	110.35	105.15
2	A	801	FAD	C7M-C7-C6	-2.62	114.96	119.57
2	A	801	FAD	C8M-C8-C9	-2.61	114.97	119.57
2	A	801	FAD	C5A-C4A-N9A	-2.60	102.98	105.81
3	A	901	MES	O3S-S-O1S	2.48	117.60	111.40
2	A	801	FAD	C3B-C2B-C1B	2.42	106.04	101.46
2	A	801	FAD	C4X-C4-N3	2.40	119.37	113.25
2	A	801	FAD	C1B-N9A-C8A	-2.22	122.18	127.09
2	A	801	FAD	C4-N3-C2	-2.21	121.71	125.64
2	A	801	FAD	O2A-PA-O1A	2.15	122.43	112.44
2	A	801	FAD	C5A-C4A-N3A	-2.07	123.86	126.72
2	A	801	FAD	C7M-C7-C8	2.06	124.96	120.76
2	A	801	FAD	C4A-C5A-N7A	-2.05	108.24	110.58
2	A	801	FAD	C9A-C5X-N5	-2.02	120.32	122.45

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	901	MES	C8-C7-N4-C5

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	901	MES	N4-C7-C8-S
2	A	801	FAD	PA-O3P-P-O5'
2	A	801	FAD	P-O3P-PA-O2A
2	A	801	FAD	O4B-C4B-C5B-O5B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	576/623 (92%)	-0.02	39 (6%) 23 27	4, 10, 23, 37	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	387	GLY	5.4
1	A	399	ALA	4.1
1	A	232	GLY	4.1
1	A	81	ASP	3.9
1	A	618	PHE	3.9
1	A	459	VAL	3.8
1	A	384	GLY	3.7
1	A	389	LEU	3.7
1	A	386	PRO	3.7
1	A	47	TYR	3.7
1	A	458	ALA	3.7
1	A	407	TRP	3.7
1	A	344	ASN	3.5
1	A	45	ILE	3.5
1	A	456	TYR	3.5
1	A	393	VAL	3.4
1	A	48	ASP	3.4
1	A	398	GLY	3.3
1	A	391	TYR	3.3
1	A	400	SER	3.2
1	A	385	THR	3.2
1	A	390	THR	3.1
1	A	342	PRO	2.8
1	A	453	ALA	2.7
1	A	397	PRO	2.7
1	A	309	PHE	2.6
1	A	343	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	401	THR	2.5
1	A	300	ILE	2.4
1	A	452	ASP	2.4
1	A	345	PRO	2.3
1	A	43	MET	2.2
1	A	82	SER	2.1
1	A	44	ASP	2.1
1	A	561	GLU	2.1
1	A	405	PRO	2.1
1	A	299	HIS	2.1
1	A	460	GLN	2.1
1	A	396	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MES	A	901	12/12	0.97	0.09	14,16,17,18	0
2	FAD	A	801	53/53	0.99	0.05	4,7,10,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.