



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:29 PM UTC

PDB ID : 5FD2 / pdb\_00005fd2  
Title : B-Raf wild-type kinase domain in complex with a purinylpyridinylamino-based inhibitor  
Authors : Whittington, D.A.; Epstein, L.F.  
Deposited on : 2015-12-15  
Resolution : 2.89 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	 12% 69% 18% • 12%
1	B	300	 12% 60% 17% • 22%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4072 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Serine/threonine-protein kinase B-raf.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	264	2115	1355	370	377	13	0	0	0
1	B	234	1863	1195	321	334	13	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	427	HIS	-	expression tag	UNP P15056
A	428	HIS	-	expression tag	UNP P15056
A	429	HIS	-	expression tag	UNP P15056
A	430	HIS	-	expression tag	UNP P15056
A	431	HIS	-	expression tag	UNP P15056
A	432	HIS	-	expression tag	UNP P15056
B	427	HIS	-	expression tag	UNP P15056
B	428	HIS	-	expression tag	UNP P15056
B	429	HIS	-	expression tag	UNP P15056
B	430	HIS	-	expression tag	UNP P15056
B	431	HIS	-	expression tag	UNP P15056
B	432	HIS	-	expression tag	UNP P15056

- Molecule 2 is 6-[2-[[3-(dimethylsulfamoylamino)-2,6-bis(fluoranyl)phenyl]amino]pyridin-3-yl]-7 {H}-purine (CCD ID: 5XJ) (formula: C<sub>18</sub>H<sub>16</sub>F<sub>2</sub>N<sub>8</sub>O<sub>2</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	N	O			S
2	A	1	31	18	2	8	2	1	0	0
2	B	1	31	18	2	8	2	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	30	30	30	0	0
3	B	2	2	2	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase B-raf



- Molecule 1: Serine/threonine-protein kinase B-raf



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.79Å 80.09Å 245.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.89 50.00 – 2.89	Depositor EDS
% Data completeness (in resolution range)	97.6 (50.00-2.89) 97.7 (50.00-2.89)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.31 (at 2.91Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.213 , 0.278 0.213 , 0.275	Depositor DCC
$R_{free}$ test set	952 reflections (5.68%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	50.4	Xtrriage
Anisotropy	0.076	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 48.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4072	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5XJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.78	0/2161	0.88	0/2916
1	B	0.67	0/1902	0.77	0/2568
All	All	0.73	0/4063	0.83	0/5484

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2115	0	2150	33	0
1	B	1863	0	1853	37	0
2	A	31	0	0	3	0
2	B	31	0	0	2	0
3	A	30	0	0	0	0
3	B	2	0	0	0	0
All	All	4072	0	4003	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:484:MET:HE3	1:A:524:GLN:OE1	1.63	0.98
1:B:498:PHE:CE2	1:B:525:LEU:HD11	2.14	0.83
1:B:518:GLY:O	1:B:528:VAL:HG12	1.85	0.77
1:A:650:MET:SD	1:A:689:MET:HE3	2.30	0.72
1:A:575:ARG:HD2	1:A:598:ALA:HB1	1.72	0.70
1:B:639:VAL:HG13	1:B:710:ILE:HD11	1.75	0.69
1:A:548:PHE:HB2	1:A:553:LEU:HD21	1.76	0.68
1:B:480:VAL:HG12	1:B:517:MET:HE1	1.74	0.67
1:B:668:MET:HE3	1:B:674:LEU:HB2	1.82	0.61
1:A:712:ALA:HB3	1:B:716:LEU:HD21	1.82	0.61
1:B:650:MET:HE3	1:B:685:CYS:HB3	1.82	0.60
1:B:514:LEU:HD23	2:B:801:5XJ:C29	2.32	0.59
1:A:712:ALA:HB3	1:B:716:LEU:CD2	2.34	0.58
1:B:517:MET:HE2	1:B:529:THR:C	2.29	0.58
1:A:576:ASP:O	1:A:581:ASN:ND2	2.37	0.57
1:B:504:VAL:O	1:B:508:THR:HG23	2.03	0.57
1:B:626:ARG:O	1:B:627:MET:C	2.47	0.57
1:A:717:LEU:C	1:A:717:LEU:HD12	2.29	0.57
1:A:626:ARG:NH2	1:A:670:GLY:O	2.37	0.56
1:B:642:PHE:CZ	1:B:646:LEU:HD21	2.41	0.56
1:A:505:LEU:HD21	1:A:596:GLY:CA	2.36	0.55
1:B:540:HIS:CD2	1:B:584:LEU:HD12	2.40	0.55
1:B:650:MET:HE3	1:B:685:CYS:CB	2.37	0.55
1:B:620:MET:HB2	1:B:625:ILE:HD12	1.89	0.54
1:B:582:ILE:HG23	1:B:590:VAL:HG13	1.90	0.54
1:A:505:LEU:HD21	1:A:596:GLY:HA2	1.90	0.54
1:B:454:ASP:N	1:B:454:ASP:OD1	2.41	0.53
1:A:548:PHE:HB2	1:A:553:LEU:CD2	2.37	0.53
1:A:586:GLU:HA	1:A:586:GLU:OE1	2.10	0.52
2:A:801:5XJ:N8	2:A:801:5XJ:N16	2.57	0.52
1:B:482:VAL:HG22	1:B:528:VAL:HG23	1.91	0.52
1:A:650:MET:SD	1:A:689:MET:HG2	2.50	0.51
1:A:600:VAL:HG13	1:A:604:TRP:CE3	2.48	0.49
1:B:471:VAL:HG22	1:B:483:LYS:HA	1.95	0.48
1:B:557:ALA:HB2	1:B:649:LEU:CD1	2.43	0.48
1:A:677:ASP:C	1:A:677:ASP:OD1	2.57	0.47
1:B:622:PRO:O	1:B:625:ILE:HG22	2.14	0.47
1:A:463:ILE:HD11	1:A:473:LYS:HB2	1.96	0.47
1:A:669:VAL:HG23	1:A:674:LEU:HD23	1.96	0.46
1:A:650:MET:HE1	1:A:689:MET:HG2	1.97	0.46
1:B:498:PHE:CG	1:B:525:LEU:HD21	2.51	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:717:LEU:HD12	1:B:717:LEU:C	2.41	0.45
1:B:656:TYR:CD1	1:B:674:LEU:HD11	2.52	0.45
1:B:717:LEU:HD12	1:B:717:LEU:O	2.17	0.44
1:A:587:ASP:OD1	1:A:589:THR:N	2.50	0.44
1:B:686:PRO:O	1:B:687:LYS:C	2.60	0.44
1:B:578:LYS:HG2	1:B:581:ASN:ND2	2.32	0.43
1:A:582:ILE:HG23	1:A:590:VAL:CG1	2.48	0.43
1:A:584:LEU:CD2	1:A:590:VAL:HG22	2.49	0.43
1:B:498:PHE:CD2	1:B:525:LEU:HD21	2.53	0.43
1:A:647:TYR:CZ	1:A:651:THR:HG21	2.54	0.43
1:B:457:ILE:O	1:B:458:THR:CB	2.66	0.43
1:A:505:LEU:HD13	2:A:801:5XJ:C28	2.49	0.43
1:A:582:ILE:HG23	1:A:590:VAL:HG13	2.00	0.43
1:A:452:ILE:HD12	1:A:528:VAL:HG21	2.01	0.42
1:A:471:VAL:HG21	2:A:801:5XJ:C13	2.49	0.42
1:B:665:ILE:O	1:B:669:VAL:HG23	2.19	0.42
1:A:650:MET:HE3	1:A:685:CYS:HB3	2.00	0.42
1:B:521:THR:O	1:B:521:THR:HG23	2.20	0.42
1:A:626:ARG:O	1:A:628:GLN:N	2.53	0.42
2:B:801:5XJ:N8	2:B:801:5XJ:N16	2.65	0.42
1:A:552:LYS:O	1:A:556:ILE:HG12	2.19	0.42
1:B:564:MET:HE1	1:B:574:HIS:HB2	2.01	0.42
1:B:686:PRO:HG2	1:B:717:LEU:HD13	2.02	0.41
1:A:578:LYS:HD2	1:A:580:ASN:HB2	2.03	0.41
1:B:650:MET:HE3	1:B:685:CYS:SG	2.61	0.41
1:B:696:CYS:O	1:B:704:ARG:HD2	2.21	0.41
1:A:635:PHE:O	1:A:639:VAL:HG23	2.21	0.40
1:B:639:VAL:HG13	1:B:710:ILE:CD1	2.47	0.40
1:A:555:ASP:OD1	1:A:558:ARG:NH1	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/300 (87%)	246 (95%)	12 (5%)	2 (1%)	16	44
1	B	226/300 (75%)	207 (92%)	15 (7%)	4 (2%)	6	25
All	All	486/600 (81%)	453 (93%)	27 (6%)	6 (1%)	10	34

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	523	PRO
1	A	627	MET
1	B	627	MET
1	A	588	LEU
1	B	594	ASP
1	B	545	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/266 (87%)	219 (94%)	13 (6%)	19	50
1	B	201/266 (76%)	194 (96%)	7 (4%)	32	66
All	All	433/532 (81%)	413 (95%)	20 (5%)	24	57

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	451	GLU
1	A	506	ARG
1	A	521	THR
1	A	551	ILE
1	A	587	ASP
1	A	588	LEU
1	A	600	VAL
1	A	614	SER
1	A	623	GLU
1	A	625	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	657	SER
1	A	664	GLN
1	A	720	SER
1	B	454	ASP
1	B	498	PHE
1	B	500	ASN
1	B	623	GLU
1	B	625	ILE
1	B	657	SER
1	B	717	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	512	ASN
1	A	539	HIS
1	A	581	ASN
1	A	631	ASN
1	B	539	HIS
1	B	581	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	5XJ	B	801	-	33,34,34	2.57	6 (18%)	47,50,50	2.58	21 (44%)
2	5XJ	A	801	-	33,34,34	2.55	6 (18%)	47,50,50	2.71	22 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5XJ	B	801	-	-	8/19/19/19	0/4/4/4
2	5XJ	A	801	-	-	8/19/19/19	0/4/4/4

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	5XJ	C10-C15	-7.93	1.41	1.49
2	A	801	5XJ	C10-C15	-7.76	1.41	1.49
2	B	801	5XJ	O27-S24	6.39	1.50	1.43
2	A	801	5XJ	C3-N7	-6.28	1.32	1.42
2	A	801	5XJ	O25-S24	6.21	1.50	1.43
2	B	801	5XJ	O25-S24	6.12	1.49	1.43
2	B	801	5XJ	C19-N21	-5.55	1.33	1.38
2	A	801	5XJ	O27-S24	5.53	1.49	1.43
2	B	801	5XJ	C3-N7	-5.33	1.34	1.42
2	A	801	5XJ	C19-N21	-5.15	1.34	1.38
2	B	801	5XJ	C20-N23	-2.29	1.34	1.37
2	A	801	5XJ	C1-N8	-2.11	1.34	1.41

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	5XJ	C19-N21-C22	7.38	108.53	103.84
2	B	801	5XJ	O27-S24-O25	-7.35	106.67	119.80
2	A	801	5XJ	O27-S24-O25	-6.17	108.79	119.80
2	B	801	5XJ	C19-N21-C22	6.00	107.66	103.84

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	5XJ	C20-C19-N18	-5.33	119.58	124.21
2	B	801	5XJ	C20-C19-N18	-5.28	119.63	124.21
2	A	801	5XJ	F30-C6-C1	4.59	122.40	117.64
2	A	801	5XJ	N18-C17-N16	-4.40	121.92	128.58
2	A	801	5XJ	N18-C19-N21	4.31	132.92	125.78
2	B	801	5XJ	C10-C9-N8	4.02	124.18	119.97
2	B	801	5XJ	N18-C19-N21	3.98	132.38	125.78
2	B	801	5XJ	N18-C17-N16	-3.85	122.76	128.58
2	A	801	5XJ	C17-N18-C19	3.82	120.29	113.19
2	A	801	5XJ	C20-C19-N21	-3.73	107.50	110.46
2	A	801	5XJ	N23-C22-N21	-3.57	108.44	113.87
2	B	801	5XJ	C13-N14-C9	3.45	123.45	116.74
2	B	801	5XJ	C17-N18-C19	3.45	119.61	113.19
2	A	801	5XJ	C29-N26-C28	-3.38	108.05	114.75
2	A	801	5XJ	C20-N23-C22	3.37	109.09	106.27
2	B	801	5XJ	C20-C19-N21	-3.10	108.00	110.46
2	B	801	5XJ	C29-N26-C28	-3.08	108.65	114.75
2	A	801	5XJ	C13-N14-C9	3.05	122.67	116.74
2	B	801	5XJ	C29-N26-S24	-2.99	107.70	117.71
2	B	801	5XJ	C20-C15-N16	-2.95	117.09	120.35
2	B	801	5XJ	N23-C22-N21	-2.85	109.53	113.87
2	B	801	5XJ	C20-N23-C22	2.84	108.64	106.27
2	A	801	5XJ	C20-C15-N16	-2.71	117.35	120.35
2	A	801	5XJ	O27-S24-N26	2.70	111.43	107.13
2	B	801	5XJ	C6-C1-C2	2.69	118.58	115.64
2	A	801	5XJ	C6-C1-C2	2.65	118.55	115.64
2	A	801	5XJ	C17-N16-C15	2.60	121.89	117.87
2	A	801	5XJ	C10-C15-N16	2.58	119.77	115.17
2	A	801	5XJ	C15-C20-N23	2.52	134.59	130.36
2	B	801	5XJ	C10-C15-N16	2.50	119.63	115.17
2	B	801	5XJ	C15-C20-N23	2.44	134.46	130.36
2	B	801	5XJ	C17-N16-C15	2.38	121.55	117.87
2	B	801	5XJ	C28-N26-S24	-2.31	109.97	117.71
2	B	801	5XJ	C10-C9-N14	-2.28	119.01	121.88
2	A	801	5XJ	C28-N26-S24	-2.26	110.14	117.71
2	B	801	5XJ	O25-S24-N26	2.18	110.60	107.13
2	A	801	5XJ	C12-C13-N14	-2.14	120.03	123.42
2	A	801	5XJ	C2-C3-N7	-2.10	118.30	120.73
2	A	801	5XJ	C29-N26-S24	-2.05	110.83	117.71

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	801	5XJ	C28-N26-S24-O27
2	A	801	5XJ	C28-N26-S24-N7
2	A	801	5XJ	C3-N7-S24-O25
2	B	801	5XJ	C28-N26-S24-O25
2	B	801	5XJ	C28-N26-S24-N7
2	B	801	5XJ	C29-N26-S24-O27
2	B	801	5XJ	C3-N7-S24-O25
2	A	801	5XJ	C11-C10-C15-N16
2	A	801	5XJ	C11-C10-C15-C20
2	B	801	5XJ	C11-C10-C15-C20
2	B	801	5XJ	C11-C10-C15-N16
2	A	801	5XJ	C2-C1-N8-C9
2	A	801	5XJ	C6-C1-N8-C9
2	A	801	5XJ	C9-C10-C15-C20
2	B	801	5XJ	C2-C1-N8-C9
2	B	801	5XJ	C9-C10-C15-C20

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	5XJ	2	0
2	A	801	5XJ	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	264/300 (88%)	-0.13	4 (1%) 72 64	18, 37, 72, 111	0
1	B	234/300 (78%)	0.86	35 (14%) 5 4	41, 72, 164, 187	0
All	All	498/600 (83%)	0.34	39 (7%) 19 16	18, 53, 147, 187	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	497	ALA	6.4
1	B	587	ASP	4.4
1	B	453	PRO	3.6
1	B	458	THR	3.4
1	B	482	VAL	3.3
1	B	470	THR	3.2
1	B	449	ASP	3.2
1	B	485	LEU	3.1
1	B	525	LEU	3.1
1	B	685	CYS	3.1
1	B	498	PHE	3.0
1	B	499	LYS	2.9
1	B	532	CYS	2.9
1	B	479	ASP	2.8
1	B	455	GLY	2.8
1	B	472	TYR	2.8
1	B	456	GLN	2.7
1	B	507	LYS	2.7
1	B	596	GLY	2.7
1	B	450	TRP	2.6
1	B	527	ILE	2.6
1	B	554	ILE	2.4
1	B	504	VAL	2.4
1	B	530	GLN	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	614	SER	2.3
1	B	627	MET	2.3
1	B	452	ILE	2.3
1	A	587	ASP	2.3
1	A	588	LEU	2.3
1	B	614	SER	2.3
1	B	517	MET	2.3
1	B	628	GLN	2.2
1	B	471	VAL	2.2
1	B	514	LEU	2.1
1	B	720	SER	2.1
1	B	511	VAL	2.1
1	A	604	TRP	2.1
1	B	512	ASN	2.0
1	B	709	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	5XJ	B	801	31/31	0.82	0.14	59,74,85,90	0
2	5XJ	A	801	31/31	0.97	0.07	24,27,33,34	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 5XJ B 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 5XJ A 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.