



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:19 AM UTC

PDB ID : 6FDU / pdb_00006fd
Title : Structure of Chlamydia trachomatis effector protein Cdu1 bound to Compound 3
Authors : Ramirez, Y.; Kisker, C.; Altmann, E.
Deposited on : 2017-12-26
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	266	
1	B	266	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4087 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deubiquitinase and deneddylase Dub1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	240	Total	C	N	O	S	0	3	0
			1948	1267	318	349	14			
1	B	239	Total	C	N	O	S	0	5	0
			1959	1274	322	350	13			

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	136	MET	-	initiating methionine	UNP B0B9A0
A	137	LYS	-	expression tag	UNP B0B9A0
A	138	HIS	-	expression tag	UNP B0B9A0
A	139	HIS	-	expression tag	UNP B0B9A0
A	140	HIS	-	expression tag	UNP B0B9A0
A	141	HIS	-	expression tag	UNP B0B9A0
A	142	HIS	-	expression tag	UNP B0B9A0
A	143	HIS	-	expression tag	UNP B0B9A0
A	144	SER	-	expression tag	UNP B0B9A0
A	145	ALA	-	expression tag	UNP B0B9A0
A	146	GLY	-	expression tag	UNP B0B9A0
A	147	LEU	-	expression tag	UNP B0B9A0
A	148	GLU	-	expression tag	UNP B0B9A0
A	149	VAL	-	expression tag	UNP B0B9A0
A	150	LEU	-	expression tag	UNP B0B9A0
A	151	PHE	-	expression tag	UNP B0B9A0
A	152	GLN	-	expression tag	UNP B0B9A0
A	153	GLY	-	expression tag	UNP B0B9A0
A	154	PRO	-	expression tag	UNP B0B9A0
B	136	MET	-	initiating methionine	UNP B0B9A0
B	137	LYS	-	expression tag	UNP B0B9A0
B	138	HIS	-	expression tag	UNP B0B9A0
B	139	HIS	-	expression tag	UNP B0B9A0
B	140	HIS	-	expression tag	UNP B0B9A0
B	141	HIS	-	expression tag	UNP B0B9A0

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	142	HIS	-	expression tag	UNP B0B9A0
B	143	HIS	-	expression tag	UNP B0B9A0
B	144	SER	-	expression tag	UNP B0B9A0
B	145	ALA	-	expression tag	UNP B0B9A0
B	146	GLY	-	expression tag	UNP B0B9A0
B	147	LEU	-	expression tag	UNP B0B9A0
B	148	GLU	-	expression tag	UNP B0B9A0
B	149	VAL	-	expression tag	UNP B0B9A0
B	150	LEU	-	expression tag	UNP B0B9A0
B	151	PHE	-	expression tag	UNP B0B9A0
B	152	GLN	-	expression tag	UNP B0B9A0
B	153	GLY	-	expression tag	UNP B0B9A0
B	154	PRO	-	expression tag	UNP B0B9A0

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	B	1	Total Cl 1 1	0	0

- Molecule 3 is (2 {S},3 {S})-2-[[[(2 {S})-2-[3,5-bis(chloranyl)phenyl]-2-(dimethylamino)ethanoyl]amino]- {N}]-[[2-(iminomethyl)pyrimidin-4-yl]methyl]-3-methyl-pentanamide (CCD ID: D5W) (formula: C₂₂H₂₈Cl₂N₆O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	Cl	N	O	0	0
			32	22	2	6	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	52	Total	O	0	0
			52	52		
4	B	94	Total	O	0	0
			94	94		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Deubiquitinase and deneddylase Dub1



- Molecule 1: Deubiquitinase and deneddylase Dub1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.08Å 57.09Å 116.38Å 90.00° 96.00° 90.00°	Depositor
Resolution (Å)	43.84 – 2.30 43.84 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.84-2.30) 99.7 (43.84-2.30)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.200 , 0.240 0.205 , 0.242	Depositor DCC
R_{free} test set	1523 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å ²)	35.0	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 25.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4087	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, D5W

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.16	7/2013 (0.3%)	1.20	2/2737 (0.1%)
1	B	1.19	10/2028 (0.5%)	1.21	7/2758 (0.3%)
All	All	1.18	17/4041 (0.4%)	1.21	9/5495 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	324	SER	N-CA	7.98	1.57	1.46
1	B	271	VAL	N-CA	7.69	1.53	1.46
1	A	280	ILE	N-CA	-7.66	1.37	1.46
1	B	284	GLN	C-O	-7.65	1.15	1.24
1	A	316	ASP	C-O	-6.81	1.16	1.24
1	B	280	ILE	N-CA	-6.67	1.38	1.46
1	B	180	THR	C-O	-6.19	1.16	1.24
1	B	337	ILE	C-O	-6.08	1.17	1.23
1	A	296	ASN	C-O	-5.86	1.17	1.23
1	B	316	ASP	C-O	-5.82	1.17	1.24
1	B	317	GLN	C-O	-5.68	1.17	1.24
1	A	216	GLY	C-O	-5.60	1.16	1.24
1	A	284	GLN	C-O	-5.60	1.17	1.24
1	B	236	LEU	C-O	-5.50	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	250	PRO	C-O	-5.43	1.17	1.24
1	B	237	GLN	C-O	-5.32	1.18	1.24
1	B	216	GLY	C-O	-5.02	1.16	1.24

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	260	GLN	CB-CA-C	5.71	120.27	110.79
1	B	186	ILE	CA-C-O	5.56	122.43	119.15
1	B	400	GLU	CB-CG-CD	5.39	121.76	112.60
1	B	284	GLN	CB-CG-CD	5.34	121.68	112.60
1	A	400	GLU	CB-CG-CD	5.33	121.67	112.60
1	B	400	GLU	CB-CA-C	5.22	118.94	110.22
1	B	215	ARG	NE-CZ-NH1	-5.17	116.33	121.50
1	B	303	ASN	CB-CA-C	-5.03	102.29	110.85
1	B	384	GLN	CB-CG-CD	-5.03	104.05	112.60

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	163	VAL	Peptide
1	A	339	ARG	Sidechain
1	B	215	ARG	Sidechain
1	B	238	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1948	0	1955	23	0
1	B	1959	0	1962	21	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	B	32	0	0	1	0
4	A	52	0	0	0	0
4	B	94	0	0	3	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4087	0	3917	43	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:162:LEU:O	1:A:163:VAL:HG23	1.82	0.79
1:A:253:LYS:HA	1:A:253:LYS:HE3	1.71	0.71
1:B:220:LYS:O	1:B:221:GLY:O	2.08	0.71
1:B:302[A]:GLU:H	1:B:302[A]:GLU:CD	2.02	0.67
1:A:220:LYS:O	1:A:221:GLY:O	2.16	0.64
1:A:229:LYS:HD3	1:A:230:GLU:N	2.13	0.64
1:B:383:VAL:O	1:B:387:GLU:HG3	2.05	0.56
1:B:335:GLU:OE1	1:B:339[A]:ARG:NH1	2.32	0.56
1:A:234:ARG:NH1	1:A:400:GLU:HG3	2.22	0.54
1:B:220:LYS:O	1:B:221:GLY:C	2.51	0.54
1:B:338:GLN:HE22	1:B:345:CYS:HB3	1.74	0.53
1:A:162:LEU:O	1:A:163:VAL:CG2	2.56	0.52
1:A:220:LYS:O	1:A:221:GLY:C	2.54	0.51
1:A:163:VAL:HG22	1:A:380:ALA:HB2	1.93	0.50
1:A:229:LYS:HD3	1:A:230:GLU:HG3	1.93	0.50
1:B:363:ASP:OD1	1:B:366[B]:ASN:OD1	2.30	0.49
1:A:229:LYS:CD	1:A:230:GLU:HG3	2.43	0.48
1:A:334:LYS:O	1:B:271:VAL:CG1	2.61	0.48
1:A:302[A]:GLU:H	1:A:302[A]:GLU:CD	2.22	0.48
1:B:234:ARG:NH1	1:B:400:GLU:HG3	2.31	0.46
1:A:334:LYS:O	1:B:271:VAL:HG13	2.16	0.45
3:B:501:D5W:CAN	3:B:501:D5W:CAV	2.94	0.45
1:B:317:GLN:HG2	4:B:686:HOH:O	2.17	0.44
1:A:253:LYS:HA	1:A:253:LYS:CE	2.43	0.44
1:B:235:HIS:CD2	1:B:401:ALA:HB2	2.54	0.43
1:A:286:ARG:O	1:A:327:PHE:HA	2.19	0.43
1:B:238:ARG:HH11	1:B:238:ARG:HG3	1.84	0.43
1:A:185:LEU:C	1:A:185:LEU:HD23	2.44	0.43
1:A:215:ARG:HD2	1:A:217:GLU:CD	2.44	0.42
1:B:220:LYS:C	1:B:221:GLY:O	2.61	0.42
1:A:241:LEU:C	1:A:241:LEU:HD23	2.44	0.42
1:B:200:LYS:HD2	1:B:311:PHE:CD1	2.55	0.42
1:A:198:ASN:HA	1:A:201:GLN:OE1	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:286:ARG:O	1:B:327:PHE:HA	2.19	0.42
1:B:211:LYS:NZ	1:B:320:PRO:HD2	2.35	0.41
1:B:273:ALA:O	1:B:297:TYR:OH	2.33	0.41
1:A:223:LEU:HD21	1:A:229:LYS:HA	2.02	0.41
1:B:215:ARG:HD2	1:B:217:GLU:CD	2.46	0.41
1:B:324:SER:HA	4:B:680:HOH:O	2.20	0.40
1:A:162:LEU:C	1:A:163:VAL:HG23	2.46	0.40
1:B:272:GLY:HA3	4:B:614:HOH:O	2.20	0.40
1:A:220:LYS:C	1:A:221:GLY:O	2.63	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:650:HOH:O	4:B:667:HOH:O[1_455]	2.11	0.09
1:B:169:ASP:OD2	1:B:220:LYS:NZ[2_448]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/266 (91%)	232 (96%)	8 (3%)	1 (0%)	30	38
1	B	242/266 (91%)	235 (97%)	6 (2%)	1 (0%)	30	38
All	All	483/532 (91%)	467 (97%)	14 (3%)	2 (0%)	30	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	221	GLY
1	B	221	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/237 (92%)	202 (93%)	16 (7%)	13	18
1	B	219/237 (92%)	210 (96%)	9 (4%)	27	41
All	All	437/474 (92%)	412 (94%)	25 (6%)	20	27

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	162	LEU
1	A	170	ILE
1	A	202	LYS
1	A	229	LYS
1	A	230	GLU
1	A	253	LYS
1	A	257	ASP
1	A	263	LYS
1	A	267	ILE
1	A	271	VAL
1	A	277	LEU
1	A	302[A]	GLU
1	A	302[B]	GLU
1	A	306	LYS
1	A	334	LYS
1	A	339	ARG
1	B	202	LYS
1	B	220	LYS
1	B	229	LYS
1	B	263	LYS
1	B	271	VAL
1	B	302[A]	GLU
1	B	302[B]	GLU
1	B	335	GLU
1	B	393	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	264	GLN
1	A	284	GLN
1	B	284	GLN
1	B	338	GLN
1	B	354	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	D5W	B	501	1	31,33,33	1.32	6 (19%)	42,45,45	2.22	15 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	D5W	B	501	1	-	5/31/33/33	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	501	D5W	CAN-CAI	2.95	1.43	1.39
3	B	501	D5W	CAM-CLB	2.66	1.80	1.74
3	B	501	D5W	CAK-CLA	2.55	1.80	1.74
3	B	501	D5W	CAJ-CAK	2.34	1.42	1.38
3	B	501	D5W	CAI-CAH	2.06	1.55	1.52
3	B	501	D5W	CAJ-CAI	2.03	1.42	1.39

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	501	D5W	CAH-CAG-N	4.55	120.04	114.59
3	B	501	D5W	CAN-CAM-CLB	4.39	124.67	119.17
3	B	501	D5W	C6-C5-C4	4.18	120.91	116.68
3	B	501	D5W	CAJ-CAK-CLA	4.08	124.28	119.17
3	B	501	D5W	CAL-CAK-CLA	-3.85	114.33	119.17
3	B	501	D5W	CAI-CAH-CAG	3.72	119.38	111.80
3	B	501	D5W	C5-C4-N3	-3.58	117.88	122.40
3	B	501	D5W	C5-C6-N1	-3.47	119.73	123.97
3	B	501	D5W	CAM-CAN-CAI	3.29	123.35	119.43
3	B	501	D5W	CAJ-CAI-CAH	3.00	124.41	119.81
3	B	501	D5W	C-CA-N	2.84	118.04	110.32
3	B	501	D5W	CB-CA-N	-2.51	106.19	111.35
3	B	501	D5W	CAL-CAM-CLB	-2.47	116.07	119.17
3	B	501	D5W	CAN-CAI-CAJ	-2.17	115.14	118.09
3	B	501	D5W	CAM-CAL-CAK	2.06	119.91	117.43

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	501	D5W	N-CAG-CAH-NAU
3	B	501	D5W	OAQ-CAG-CAH-NAU
3	B	501	D5W	CG2-CB-CG1-CD1
3	B	501	D5W	N3-C4-CAX-NAF
3	B	501	D5W	C5-C4-CAX-NAF

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	501	D5W	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/266 (90%)	0.11	9 (3%) 44 46	22, 42, 75, 102	3 (1%)
1	B	239/266 (89%)	0.06	9 (3%) 44 46	19, 38, 72, 123	5 (2%)
All	All	479/532 (90%)	0.08	18 (3%) 44 46	19, 40, 75, 123	8 (1%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	163	VAL	6.2
1	A	162	LEU	4.4
1	B	267	ILE	4.3
1	B	270	SER	4.2
1	B	164	PRO	3.4
1	B	271	VAL	3.4
1	B	265	LEU	3.0
1	A	163	VAL	2.9
1	A	271	VAL	2.9
1	A	191	PHE	2.7
1	B	269	GLY	2.6
1	A	221	GLY	2.6
1	A	222	PRO	2.5
1	A	267	ILE	2.5
1	B	221	GLY	2.1
1	A	265	LEU	2.1
1	B	263	LYS	2.0
1	A	268	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	D5W	B	501	32/32	0.79	0.19	35,89,119,126	0
2	CL	B	502	1/1	0.93	0.22	72,72,72,72	0
2	CL	A	501	1/1	0.95	0.23	67,67,67,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.