



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 12:15 PM UTC

PDB ID : 3FER / pdb\_00003fer  
Title : Crystal structure of n-terminal actin-binding domain from human filamin b (tandem ch-domains). northeast structural genomics consortium target hr5571a.  
Authors : Kuzin, A.P.; Abashidze, M.; Seetharaman, R.; Shastry, R.; Sahdev, S.; Ciccosanti, C.; Xiao, R.; Everett, J.K.; Huang, Y.; Acton, T.; Rost, B.; Montelione, G.T.; Tong, L.; Hunt, J.F.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2008-11-30  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	 2% 59% 27% 11%
1	B	262	 3% 53% 31% 5% 11%
1	C	262	 5% 61% 22% 14%

*Continued on next page...*

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	D	262	 <p>6% 54% 30% 11%</p>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7688 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Filamin-B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	234	1900	1219	330	342	4	5	0	0	0
1	B	234	1900	1219	330	342	4	5	0	0	0
1	C	225	1827	1172	320	327	4	4	0	0	0
1	D	234	1900	1219	330	342	4	5	0	0	0

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	expression tag	UNP O75369
A	2	GLY	-	expression tag	UNP O75369
A	3	HIS	-	expression tag	UNP O75369
A	4	HIS	-	expression tag	UNP O75369
A	5	HIS	-	expression tag	UNP O75369
A	6	HIS	-	expression tag	UNP O75369
A	7	HIS	-	expression tag	UNP O75369
A	8	HIS	-	expression tag	UNP O75369
A	9	SER	-	expression tag	UNP O75369
A	10	HIS	-	expression tag	UNP O75369
B	1	MSE	-	expression tag	UNP O75369
B	2	GLY	-	expression tag	UNP O75369
B	3	HIS	-	expression tag	UNP O75369
B	4	HIS	-	expression tag	UNP O75369
B	5	HIS	-	expression tag	UNP O75369
B	6	HIS	-	expression tag	UNP O75369
B	7	HIS	-	expression tag	UNP O75369
B	8	HIS	-	expression tag	UNP O75369
B	9	SER	-	expression tag	UNP O75369
B	10	HIS	-	expression tag	UNP O75369
C	1	MSE	-	expression tag	UNP O75369

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	2	GLY	-	expression tag	UNP O75369
C	3	HIS	-	expression tag	UNP O75369
C	4	HIS	-	expression tag	UNP O75369
C	5	HIS	-	expression tag	UNP O75369
C	6	HIS	-	expression tag	UNP O75369
C	7	HIS	-	expression tag	UNP O75369
C	8	HIS	-	expression tag	UNP O75369
C	9	SER	-	expression tag	UNP O75369
C	10	HIS	-	expression tag	UNP O75369
D	1	MSE	-	expression tag	UNP O75369
D	2	GLY	-	expression tag	UNP O75369
D	3	HIS	-	expression tag	UNP O75369
D	4	HIS	-	expression tag	UNP O75369
D	5	HIS	-	expression tag	UNP O75369
D	6	HIS	-	expression tag	UNP O75369
D	7	HIS	-	expression tag	UNP O75369
D	8	HIS	-	expression tag	UNP O75369
D	9	SER	-	expression tag	UNP O75369
D	10	HIS	-	expression tag	UNP O75369

- Molecule 2 is ACETIC ACID (CCD ID: ACY) (formula: C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	54	Total 54	O 54	0	0
3	B	47	Total 47	O 47	0	0
3	C	22	Total 22	O 22	0	0
3	D	30	Total 30	O 30	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Filamin-B




- Molecule 1: Filamin-B




- Molecule 1: Filamin-B





● Molecule 1: Filamin-B



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.85Å 82.41Å 132.25Å 90.00° 94.14° 90.00°	Depositor
Resolution (Å)	19.86 – 2.40 19.86 – 2.40	Depositor EDS
% Data completeness (in resolution range)	72.0 (19.86-2.40) 83.8 (19.86-2.40)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.06 (at 2.39Å)	Xtrriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.215 , 0.259 0.224 , 0.268	Depositor DCC
$R_{free}$ test set	4228 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.8	Xtrriage
Anisotropy	0.601	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 36.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7688	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.53 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3033e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/1943	0.97	7/2630 (0.3%)
1	B	0.47	0/1943	1.00	8/2630 (0.3%)
1	C	0.43	0/1867	0.91	3/2527 (0.1%)
1	D	0.47	0/1943	0.96	5/2630 (0.2%)
All	All	0.45	0/7696	0.96	23/10417 (0.2%)

There are no bond length outliers.

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	234	ASP	N-CA-C	-8.34	93.04	110.80
1	B	53	ASP	N-CA-C	7.89	122.17	112.54
1	B	55	SER	N-CA-C	7.86	120.75	111.71
1	A	135	MSE	N-CA-C	6.87	121.56	112.35
1	B	176	GLN	N-CA-C	6.66	118.54	111.28

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1900	0	1919	67	0
1	B	1900	0	1919	68	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1827	0	1857	59	0
1	D	1900	0	1919	84	0
2	B	4	0	3	1	0
2	C	4	0	3	0	0
3	A	54	0	0	3	0
3	B	47	0	0	4	0
3	C	22	0	0	1	0
3	D	30	0	0	2	0
All	All	7688	0	7620	278	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 278 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:50:LEU:HG	1:B:120:ILE:HD11	1.35	1.09
1:D:171:PHE:HB2	1:D:237:GLU:HG3	1.35	1.03
1:C:37:ASN:HD21	1:C:47:ILE:H	1.05	1.02
1:B:169:THR:H	1:B:174:ASN:HD21	1.12	0.98
1:D:37:ASN:HD21	1:D:47:ILE:H	1.05	0.98

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/262 (88%)	217 (94%)	12 (5%)	1 (0%)	30	43
1	B	230/262 (88%)	215 (94%)	14 (6%)	1 (0%)	30	43
1	C	221/262 (84%)	212 (96%)	7 (3%)	2 (1%)	14	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	D	230/262 (88%)	208 (90%)	14 (6%)	8 (4%)	3 2
All	All	911/1048 (87%)	852 (94%)	47 (5%)	12 (1%)	9 14

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	139	GLU
1	D	140	ASP
1	D	233	PRO
1	C	53	ASP
1	D	234	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	212/229 (93%)	204 (96%)	8 (4%)	29 49
1	B	212/229 (93%)	198 (93%)	14 (7%)	15 26
1	C	204/229 (89%)	194 (95%)	10 (5%)	22 39
1	D	212/229 (93%)	197 (93%)	15 (7%)	13 23
All	All	840/916 (92%)	793 (94%)	47 (6%)	19 33

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	209	ARG
1	D	120	ILE
1	C	212	MSE
1	D	24	TRP
1	D	141	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	223	GLN
1	D	130	HIS
1	C	246	GLN
1	D	37	ASN
1	D	174	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ACY	B	263	-	3,3,3	0.58	0	3,3,3	0.87	0
2	ACY	C	263	-	3,3,3	0.65	0	3,3,3	0.78	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	263	ACY	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	229/262 (87%)	-0.04	6 (2%) 57 53	15, 31, 60, 83	0
1	B	229/262 (87%)	0.00	7 (3%) 51 47	14, 31, 59, 76	0
1	C	221/262 (84%)	0.47	13 (5%) 28 24	21, 42, 77, 103	0
1	D	229/262 (87%)	0.27	16 (6%) 22 19	19, 37, 75, 92	0
All	All	908/1048 (86%)	0.17	42 (4%) 37 33	14, 34, 69, 103	0

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	142	GLY	5.8
1	D	233	PRO	5.4
1	D	23	PRO	4.6
1	C	260	PRO	4.4
1	C	73	TYR	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ACY	C	263	4/4	0.75	0.17	55,57,57,57	0
2	ACY	B	263	4/4	0.86	0.13	55,55,56,56	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.