



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 24, 2026 – 09:19 AM UTC

PDB ID : 4FE8 / pdb_00004fe8
Title : Crystal Structure of Htt36Q3H-EX1-X1-C1(Alpha)
Authors : Kim, M.
Deposited on : 2012-05-29
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 9664 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein,Huntingtin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	402	Total 3146	C 2016	N 520	O 602	S 8	0	2	0
1	B	401	Total 3130	C 2007	N 520	O 595	S 8	0	1	0
1	C	410	Total 3209	C 2051	N 535	O 615	S 8	0	1	0

There are 90 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	359	ALA	-	linker	UNP P0AEX9
A	360	ALA	-	linker	UNP P0AEX9
A	361	LEU	-	linker	UNP P0AEX9
A	362	ALA	-	linker	UNP P0AEX9
A	363	ALA	-	linker	UNP P0AEX9
A	364	ALA	-	linker	UNP P0AEX9
A	365	GLN	-	linker	UNP P0AEX9
A	366	THR	-	linker	UNP P0AEX9
A	367	ASN	-	linker	UNP P0AEX9
A	368	ALA	-	linker	UNP P0AEX9
A	369	ALA	-	linker	UNP P0AEX9
A	370	ALA	-	linker	UNP P0AEX9
A	388	GLN	-	insertion	UNP P42858
A	389	GLN	-	insertion	UNP P42858
A	390	GLN	-	insertion	UNP P42858
A	391	GLN	-	insertion	UNP P42858
A	392	GLN	-	insertion	UNP P42858
A	393	GLN	-	insertion	UNP P42858
A	394	GLN	-	insertion	UNP P42858
A	395	HIS	-	insertion	UNP P42858
A	396	GLN	-	insertion	UNP P42858
A	397	HIS	-	insertion	UNP P42858
A	398	GLN	-	insertion	UNP P42858

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	399	HIS	-	insertion	UNP P42858
A	400	GLN	-	insertion	UNP P42858
A	401	GLN	-	insertion	UNP P42858
A	402	GLN	-	insertion	UNP P42858
A	403	GLN	-	insertion	UNP P42858
A	404	GLN	-	insertion	UNP P42858
A	405	GLN	-	insertion	UNP P42858
B	359	ALA	-	linker	UNP P0AEX9
B	360	ALA	-	linker	UNP P0AEX9
B	361	LEU	-	linker	UNP P0AEX9
B	362	ALA	-	linker	UNP P0AEX9
B	363	ALA	-	linker	UNP P0AEX9
B	364	ALA	-	linker	UNP P0AEX9
B	365	GLN	-	linker	UNP P0AEX9
B	366	THR	-	linker	UNP P0AEX9
B	367	ASN	-	linker	UNP P0AEX9
B	368	ALA	-	linker	UNP P0AEX9
B	369	ALA	-	linker	UNP P0AEX9
B	370	ALA	-	linker	UNP P0AEX9
B	388	GLN	-	insertion	UNP P42858
B	389	GLN	-	insertion	UNP P42858
B	390	GLN	-	insertion	UNP P42858
B	391	GLN	-	insertion	UNP P42858
B	392	GLN	-	insertion	UNP P42858
B	393	GLN	-	insertion	UNP P42858
B	394	GLN	-	insertion	UNP P42858
B	395	HIS	-	insertion	UNP P42858
B	396	GLN	-	insertion	UNP P42858
B	397	HIS	-	insertion	UNP P42858
B	398	GLN	-	insertion	UNP P42858
B	399	HIS	-	insertion	UNP P42858
B	400	GLN	-	insertion	UNP P42858
B	401	GLN	-	insertion	UNP P42858
B	402	GLN	-	insertion	UNP P42858
B	403	GLN	-	insertion	UNP P42858
B	404	GLN	-	insertion	UNP P42858
B	405	GLN	-	insertion	UNP P42858
C	359	ALA	-	linker	UNP P0AEX9
C	360	ALA	-	linker	UNP P0AEX9
C	361	LEU	-	linker	UNP P0AEX9
C	362	ALA	-	linker	UNP P0AEX9
C	363	ALA	-	linker	UNP P0AEX9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	364	ALA	-	linker	UNP P0AEX9
C	365	GLN	-	linker	UNP P0AEX9
C	366	THR	-	linker	UNP P0AEX9
C	367	ASN	-	linker	UNP P0AEX9
C	368	ALA	-	linker	UNP P0AEX9
C	369	ALA	-	linker	UNP P0AEX9
C	370	ALA	-	linker	UNP P0AEX9
C	388	GLN	-	insertion	UNP P42858
C	389	GLN	-	insertion	UNP P42858
C	390	GLN	-	insertion	UNP P42858
C	391	GLN	-	insertion	UNP P42858
C	392	GLN	-	insertion	UNP P42858
C	393	GLN	-	insertion	UNP P42858
C	394	GLN	-	insertion	UNP P42858
C	395	HIS	-	insertion	UNP P42858
C	396	GLN	-	insertion	UNP P42858
C	397	HIS	-	insertion	UNP P42858
C	398	GLN	-	insertion	UNP P42858
C	399	HIS	-	insertion	UNP P42858
C	400	GLN	-	insertion	UNP P42858
C	401	GLN	-	insertion	UNP P42858
C	402	GLN	-	insertion	UNP P42858
C	403	GLN	-	insertion	UNP P42858
C	404	GLN	-	insertion	UNP P42858
C	405	GLN	-	insertion	UNP P42858

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	5	Total Zn 5 5	0	0
2	B	13	Total Zn 13 13	0	0
2	C	13	Total Zn 13 13	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	38	Total O 38 38	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	61	Total	O	0	0
			61	61		
3	C	49	Total	O	0	0
			49	49		

MolProbity failed to run properly - this section is therefore empty.

3 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	155.05Å 177.28Å 78.87Å 90.00° 109.03° 90.00°	Depositor
Resolution (Å)	35.00 – 3.00 35.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	95.9 (35.00-3.00) 95.8 (35.00-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.40 (at 3.00Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.227 , 0.275 0.253 , 0.292	Depositor DCC
R_{free} test set	1942 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	80.8	Xtrriage
Anisotropy	0.056	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 61.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9664	wwPDB-VP
Average B, all atoms (Å ²)	90.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

4.6 Ligand geometry [i](#)

Of 31 ligands modelled in this entry, 31 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	402/452 (88%)	0.23	14 (3%) 47 27	24, 84, 120, 155	15 (3%)
1	B	401/452 (88%)	0.22	13 (3%) 50 29	32, 89, 132, 159	13 (3%)
1	C	410/452 (90%)	0.35	28 (6%) 23 12	31, 92, 141, 190	28 (6%)
All	All	1213/1356 (89%)	0.27	55 (4%) 38 20	24, 89, 132, 190	56 (4%)

The worst 5 of 55 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	402	GLN	7.7
1	A	400	GLN	7.6
1	A	401	GLN	7.3
1	C	171	TYR	6.5
1	C	408	GLN	6.3

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	B	513	1/1	0.71	0.20	60,60,60,60	1
2	ZN	B	512	1/1	0.76	0.12	60,60,60,60	1
2	ZN	C	502	1/1	0.78	0.08	60,60,60,60	1
2	ZN	C	503	1/1	0.80	0.14	60,60,60,60	1
2	ZN	A	505	1/1	0.81	0.11	60,60,60,60	1
2	ZN	B	503	1/1	0.85	0.08	60,60,60,60	1
2	ZN	C	501	1/1	0.88	0.32	60,60,60,60	1
2	ZN	C	511	1/1	0.88	0.12	60,60,60,60	1
2	ZN	C	512	1/1	0.88	0.18	60,60,60,60	1
2	ZN	C	513	1/1	0.88	0.14	60,60,60,60	1
2	ZN	B	509	1/1	0.89	0.07	60,60,60,60	1
2	ZN	B	501	1/1	0.89	0.09	60,60,60,60	1
2	ZN	C	505	1/1	0.89	0.12	60,60,60,60	1
2	ZN	B	502	1/1	0.90	0.06	60,60,60,60	1
2	ZN	C	504	1/1	0.90	0.05	60,60,60,60	1
2	ZN	A	504	1/1	0.91	0.07	60,60,60,60	1
2	ZN	A	502	1/1	0.91	0.08	60,60,60,60	1
2	ZN	B	510	1/1	0.92	0.12	55,55,55,55	1
2	ZN	C	506	1/1	0.93	0.11	60,60,60,60	1
2	ZN	A	501	1/1	0.93	0.10	60,60,60,60	1
2	ZN	A	503	1/1	0.94	0.08	60,60,60,60	1
2	ZN	C	507	1/1	0.94	0.14	60,60,60,60	1
2	ZN	C	510	1/1	0.94	0.06	60,60,60,60	1
2	ZN	B	511	1/1	0.95	0.04	60,60,60,60	1
2	ZN	B	504	1/1	0.95	0.09	60,60,60,60	1
2	ZN	B	505	1/1	0.97	0.17	60,60,60,60	1
2	ZN	B	506	1/1	0.97	0.11	59,59,59,59	1
2	ZN	B	507	1/1	0.98	0.05	60,60,60,60	1
2	ZN	B	508	1/1	0.98	0.04	60,60,60,60	1
2	ZN	C	508	1/1	0.98	0.10	60,60,60,60	1
2	ZN	C	509	1/1	0.98	0.05	60,60,60,60	1

5.5 Other polymers [i](#)

There are no such residues in this entry.