



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:59 AM UTC

PDB ID : 4FEK / pdb_00004fek
Title : Crystal Structure of putative diflavin flavoprotein A 5 (fragment 1-254) from Nostoc sp. PCC 7120, Northeast Structural Genomics Consortium Target NsR435A , Northeast Structural Genomics Consortium (NESG) Target NsR435A
Authors : Kuzin, A.; Abashidze, M.; Seetharaman, J.; Janjua, J.; Xiao, R.; Foote, E.L.; Ciccocanti, C.; Wang, H.; Everett, J.K.; Nair, R.; Acton, T.B.; Rost, B.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2012-05-30
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	: 4-5-2 with Phenix2.0
Mogul	: 2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	: 2.0
EDS	: 3.0
Percentile statistics	: 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	: 9.0.010 (Gargrove)
Density-Fitness	: 1.0.12


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	
1	B	262	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

Ideal geometry (proteins)	: Engh & Huber (2001)
Ideal geometry (DNA, RNA)	: Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	: 2.49

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PEG	A	301	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4209 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative diflavin flavoprotein A 5.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	252	Total	C	N	O	S	Se	0	2	0
			2022	1304	347	364	4	3			
1	B	248	Total	C	N	O	S	Se	0	0	0
			1976	1275	336	358	4	3			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	255	LEU	-	expression tag	UNP Q8Z0C1
A	256	GLU	-	expression tag	UNP Q8Z0C1
A	257	HIS	-	expression tag	UNP Q8Z0C1
A	258	HIS	-	expression tag	UNP Q8Z0C1
A	259	HIS	-	expression tag	UNP Q8Z0C1
A	260	HIS	-	expression tag	UNP Q8Z0C1
A	261	HIS	-	expression tag	UNP Q8Z0C1
A	262	HIS	-	expression tag	UNP Q8Z0C1
B	255	LEU	-	expression tag	UNP Q8Z0C1
B	256	GLU	-	expression tag	UNP Q8Z0C1
B	257	HIS	-	expression tag	UNP Q8Z0C1
B	258	HIS	-	expression tag	UNP Q8Z0C1
B	259	HIS	-	expression tag	UNP Q8Z0C1
B	260	HIS	-	expression tag	UNP Q8Z0C1
B	261	HIS	-	expression tag	UNP Q8Z0C1
B	262	HIS	-	expression tag	UNP Q8Z0C1

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		
2	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	137	Total	O	0	1
			138	138		
5	B	47	Total	O	0	1
			48	48		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Putative diflavin flavoprotein A 5



• Molecule 1: Putative diflavin flavoprotein A 5



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.65Å 57.99Å 106.61Å 90.00° 102.42° 90.00°	Depositor
Resolution (Å)	28.37 – 2.00 28.37 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.1 (28.37-2.00) 99.1 (28.37-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.10 (at 2.00Å)	Xtriage
Refinement program	PHENIX dev_988	Depositor
R, R_{free}	0.209 , 0.240 0.206 , 0.235	Depositor DCC
R_{free} test set	2038 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtriage
Anisotropy	0.454	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 57.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4209	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CL, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/2075	0.86	2/2814 (0.1%)
1	B	0.45	0/2019	0.79	0/2738
All	All	0.50	0/4094	0.83	2/5552 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	193	PHE	N-CA-C	-5.88	103.45	111.56
1	A	192	SER	N-CA-C	5.35	117.65	109.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2022	0	2041	22	0
1	B	1976	0	1995	24	0
2	A	7	0	10	4	0
2	B	7	0	10	0	0
3	A	1	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	138	0	0	1	0
5	B	48	0	0	1	0
All	All	4209	0	4056	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:MSE:HE2	1:A:246:TYR:HE2	1.49	0.76
1:A:205:MSE:HE3	1:A:212:VAL:HG22	1.73	0.70
1:A:205:MSE:HE2	1:A:246:TYR:CE2	2.31	0.66
1:A:7:ARG:NH1	1:A:62:SER:OG	2.30	0.65
1:B:205:MSE:HE3	1:B:212:VAL:HG22	1.80	0.63
1:B:226:LEU:HD11	1:B:234:LEU:HB3	1.81	0.63
1:A:12:LEU:HD22	1:A:13:PRO:HD2	1.84	0.59
1:B:227:TYR:HB2	1:B:235:VAL:HB	1.84	0.59
1:B:24:ARG:NH1	1:B:182:GLY:O	2.35	0.58
1:A:175:ILE:HD12	1:A:205:MSE:HE1	1.87	0.57
2:A:301:PEG:H21	5:A:420:HOH:O	2.05	0.56
1:A:100:LEU:HD13	1:A:106:ILE:HD11	1.86	0.56
1:A:205:MSE:HE3	1:A:212:VAL:CG2	2.38	0.54
1:A:211:HIS:HB3	2:A:301:PEG:H32	1.88	0.53
1:B:55:ILE:HD13	1:B:80:LEU:HD21	1.89	0.53
1:B:149:LEU:HD22	1:B:219:ILE:HG22	1.92	0.52
1:B:205:MSE:HE2	1:B:246:TYR:CE2	2.45	0.52
1:B:205:MSE:HE2	1:B:246:TYR:HE2	1.75	0.52
1:B:168:GLN:HB2	1:B:225:ARG:HG3	1.90	0.51
1:A:5:LYS:HB2	1:A:26:TRP:CD1	2.46	0.51
1:B:100:LEU:HD13	1:B:106:ILE:HD11	1.93	0.51
1:B:46:TYR:CZ	1:B:228:ALA:HB1	2.45	0.51
1:B:23:ALA:HB1	1:B:63:PHE:CD1	2.48	0.48
1:B:148:PHE:HB3	1:B:160:LEU:HD11	1.96	0.47
1:A:46:TYR:CZ	1:A:228:ALA:HB1	2.50	0.47
1:A:124:ASP:OD1	1:A:124:ASP:N	2.42	0.46
1:A:227:TYR:HB2	1:A:235:VAL:HB	1.97	0.46
1:A:5:LYS:HD2	1:A:26:TRP:CE2	2.51	0.45
1:A:211:HIS:HB3	2:A:301:PEG:H22	1.98	0.45
1:B:24:ARG:NH1	1:B:183:ASP:HA	2.33	0.44
1:B:5:LYS:HB2	1:B:26:TRP:CD1	2.53	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:194:LYS:HD2	1:B:197:GLN:OE1	2.18	0.44
1:B:24:ARG:HH12	1:B:183:ASP:HA	1.83	0.43
1:A:47:VAL:HG11	1:A:75:VAL:HG11	1.99	0.43
1:B:12:LEU:HD12	1:B:185:VAL:HG21	2.00	0.43
1:A:78:LYS:HA	1:A:78:LYS:HD2	1.78	0.43
1:A:151:ILE:O	2:A:301:PEG:H31	2.19	0.42
1:B:55:ILE:HG23	1:B:80:LEU:HD11	2.01	0.42
1:A:11:VAL:HG22	1:A:21:LEU:HG	2.02	0.41
1:B:84:ILE:HG12	1:B:109:VAL:HB	2.02	0.41
1:B:175:ILE:HD12	1:B:205:MSE:HE1	2.03	0.41
1:B:240:ILE:HB	5:B:422:HOH:O	2.20	0.41
1:A:193:PHE:CE1	1:A:233:PRO:HG2	2.55	0.41
1:A:84:ILE:HG12	1:A:109:VAL:HB	2.03	0.40
1:B:115:ALA:O	1:B:119:ARG:HG3	2.21	0.40
1:B:174:LYS:HA	1:B:174:LYS:HD3	1.93	0.40
1:A:88:PHE:CZ	1:A:93:ILE:HD11	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/262 (95%)	245 (98%)	5 (2%)	0	100	100
1	B	242/262 (92%)	231 (96%)	10 (4%)	1 (0%)	30	27
All	All	492/524 (94%)	476 (97%)	15 (3%)	1 (0%)	43	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	192	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/224 (98%)	209 (95%)	11 (5%)	22	20
1	B	214/224 (96%)	208 (97%)	6 (3%)	38	41
All	All	434/448 (97%)	417 (96%)	17 (4%)	28	28

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	LEU
1	A	28	ARG
1	A	37	LEU
1	A	75	VAL
1	A	100	LEU
1	A	112	LEU
1	A	125	ASP
1	A	127	LEU
1	A	140	LEU
1	A	146	LEU
1	A	191	GLU
1	B	12	LEU
1	B	37	LEU
1	B	75	VAL
1	B	100	LEU
1	B	128	ASN
1	B	140	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	168	GLN
1	B	17	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PEG	B	301	-	6,6,6	0.57	0	5,5,5	0.76	0
4	PO4	A	303	-	4,4,4	1.39	0	6,6,6	1.21	0
2	PEG	A	301	-	6,6,6	0.57	0	5,5,5	1.34	0
4	PO4	B	302	-	4,4,4	0.91	0	6,6,6	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	B	301	-	-	0/4/4/4	-
2	PEG	A	301	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	PEG	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	249/262 (95%)	0.09	9 (3%) 46 45	9, 27, 53, 85	2 (0%)
1	B	245/262 (93%)	1.63	95 (38%) 1 1	18, 52, 84, 109	0
All	All	494/524 (94%)	0.85	104 (21%) 2 2	9, 35, 76, 109	2 (0%)

All (104) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	192	SER	5.8
1	B	6	PRO	5.6
1	B	115	ALA	5.6
1	B	190	TRP	5.5
1	A	259	HIS	5.5
1	B	121	ALA	5.4
1	B	255	LEU	5.1
1	B	116	GLY	5.1
1	B	241	ALA	4.7
1	B	191	GLU	4.5
1	B	120	ALA	4.5
1	B	114	ALA	4.4
1	B	145	VAL	4.4
1	A	258	HIS	4.2
1	B	249	TRP	4.0
1	B	240	ILE	4.0
1	B	245	ALA	3.8
1	B	244	GLN	3.8
1	B	112	LEU	3.8
1	B	133	ARG	3.6
1	B	36	ALA	3.6
1	A	191	GLU	3.6
1	A	193	PHE	3.5
1	B	127	LEU	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	124	ASP	3.5
1	B	137	THR	3.5
1	B	38	GLU	3.4
1	B	117	ASP	3.3
1	B	100	LEU	3.3
1	B	192	SER	3.2
1	B	135	LYS	3.2
1	B	165	VAL	3.2
1	B	180	ILE	3.1
1	B	128	ASN	3.1
1	B	148	PHE	2.9
1	B	247	ALA	2.9
1	A	123	PRO	2.9
1	B	40	GLY	2.9
1	B	110	CYS	2.8
1	B	130	LEU	2.8
1	B	26	TRP	2.8
1	B	104	PRO	2.8
1	B	31	PHE	2.8
1	B	182	GLY	2.8
1	B	78	LYS	2.8
1	B	134	GLY	2.8
1	B	118	LEU	2.8
1	B	149	LEU	2.8
1	B	194	LYS	2.8
1	B	223	GLN	2.7
1	B	193	PHE	2.7
1	B	90	PRO	2.7
1	B	119	ARG	2.7
1	B	181	CYS	2.7
1	B	208	HIS	2.7
1	B	210	ILE	2.6
1	B	94	PRO	2.6
1	B	27	SER	2.6
1	B	125	ASP	2.6
1	B	166	GLN	2.6
1	B	107	THR	2.6
1	B	214	ALA	2.6
1	B	108	PHE	2.5
1	B	217	GLU	2.5
1	B	252	ALA	2.5
1	B	5	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	106	ILE	2.5
1	B	256	GLU	2.4
1	B	143	GLY	2.4
1	B	189	ASN	2.4
1	B	60	VAL	2.4
1	B	35	TYR	2.3
1	B	63	PHE	2.3
1	B	251	LYS	2.3
1	B	93	ILE	2.3
1	B	157	PRO	2.3
1	B	103	ALA	2.3
1	B	30	ARG	2.3
1	B	141	GLY	2.3
1	B	239	LEU	2.3
1	B	129	ILE	2.3
1	B	98	ALA	2.3
1	B	221	ASP	2.3
1	B	140	LEU	2.3
1	B	33	ILE	2.2
1	B	131	PRO	2.2
1	A	221	ASP	2.2
1	B	139	ASP	2.2
1	B	32	GLU	2.2
1	B	138	LEU	2.2
1	B	196	ASP	2.2
1	B	102	LEU	2.2
1	B	113	PRO	2.1
1	B	154	PRO	2.1
1	B	75	VAL	2.1
1	B	39	ARG	2.1
1	B	29	LEU	2.1
1	B	220	SER	2.1
1	B	184	ASP	2.1
1	A	257	HIS	2.1
1	B	69	GLU	2.0
1	B	198	ARG	2.0
1	B	28	ARG	2.0
1	B	159	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PEG	A	301	7/7	0.75	0.17	33,43,45,50	0
2	PEG	B	301	7/7	0.79	0.23	109,113,117,118	0
4	PO4	B	302	5/5	0.96	0.09	44,50,52,52	0
4	PO4	A	303	5/5	0.98	0.05	21,22,23,24	0
3	CL	A	302	1/1	0.99	0.02	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.