



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:37 PM UTC

PDB ID : 4FEQ / pdb_00004feq
Title : Inhibitor bound structure of the kinase domain of the murine receptor tyrosine kinase TYRO3 (Sky)
Authors : Ohren, J.F.; Powell, N.A.; Kohrt, J.T.; Perrin, L.A.
Deposited on : 2012-05-30
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	25% 59% 10% • 31%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1844 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase receptor TYRO3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	224	1776	1136	303	316	21	0	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	484	MET	-	initiating methionine	UNP P55144
A	801	HIS	-	expression tag	UNP P55144
A	802	HIS	-	expression tag	UNP P55144
A	803	HIS	-	expression tag	UNP P55144
A	804	HIS	-	expression tag	UNP P55144
A	805	HIS	-	expression tag	UNP P55144
A	806	HIS	-	expression tag	UNP P55144

- Molecule 2 is 4-(cyclopentylamino)-N-[3-(2-oxopyrrolidin-1-yl)propyl]-2-{{2-(pyridin-4-yl)ethyl}amino}pyrimidine-5-carboxamide (CCD ID: 0T8) (formula: C₂₄H₃₃N₇O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			33	24	7	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	35	Total	O	0	0
			35	35		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase receptor TYRO3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	47.78Å 57.16Å 60.68Å 90.00° 100.69° 90.00°	Depositor
Resolution (Å)	47.67 – 2.20 47.67 – 2.20	Depositor EDS
% Data completeness (in resolution range)	91.7 (47.67-2.20) 91.9 (47.67-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.65 (at 2.20Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.275 , 0.327 0.269 , 0.317	Depositor DCC
R_{free} test set	842 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	45.1	Xtriage
Anisotropy	0.199	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	1844	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 0T8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/1810	0.84	0/2434

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1776	0	1765	23	0
2	A	33	0	33	0	0
3	A	35	0	0	1	0
All	All	1844	0	1798	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:626:MET:CE	1:A:716:MET:HE3	2.03	0.88

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:626:MET:HE1	1:A:716:MET:HE3	1.64	0.77
1:A:686:VAL:N	3:A:1006:HOH:O	2.18	0.75
1:A:626:MET:CE	1:A:716:MET:CE	2.72	0.67
1:A:626:MET:HE3	1:A:716:MET:CE	2.26	0.64
1:A:687:LYS:HD3	1:A:722:PRO:O	1.98	0.62
1:A:715:ILE:HG22	1:A:716:MET:HE2	1.86	0.57
1:A:760:SER:O	1:A:766:ARG:NH1	2.38	0.57
1:A:643:HIS:HD2	1:A:645:ASP:H	1.54	0.54
1:A:783:LEU:H	1:A:783:LEU:HD23	1.76	0.51
1:A:626:MET:HE3	1:A:716:MET:HE2	1.92	0.51
1:A:626:MET:HE1	1:A:716:MET:CE	2.37	0.51
1:A:596:MET:HE1	1:A:660:CYS:SG	2.52	0.49
1:A:662:ALA:O	1:A:663:ASP:HB3	2.11	0.47
1:A:562:MET:HE2	1:A:641:PHE:HE1	1.82	0.45
1:A:563:LYS:NZ	1:A:574:VAL:O	2.50	0.44
1:A:562:MET:SD	1:A:664:PHE:HB3	2.58	0.44
1:A:567:HIS:CG	1:A:568:PRO:HD2	2.53	0.43
1:A:626:MET:HE1	1:A:716:MET:HG2	2.01	0.42
1:A:514:LEU:HD21	1:A:524:GLU:HB3	2.01	0.41
1:A:758:CYS:O	1:A:766:ARG:HD2	2.20	0.41
1:A:622:LEU:HB3	1:A:716:MET:HE1	2.02	0.41
1:A:624:ARG:HE	1:A:624:ARG:HB2	1.74	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	210/323 (65%)	205 (98%)	4 (2%)	1 (0%)	24 27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	663	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	191/280 (68%)	184 (96%)	7 (4%)	30 41

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	556	LEU
1	A	593	LEU
1	A	608	SER
1	A	697	ASN
1	A	736	ILE
1	A	766	ARG
1	A	773	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	506	GLN
1	A	507	GLN
1	A	526	GLN
1	A	643	HIS
1	A	702	HIS
1	A	778	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	OT8	A	901	-	36,36,36	0.80	3 (8%)	44,47,47	2.29	14 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OT8	A	901	-	-	8/21/38/38	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	OT8	C26-C24	2.62	1.62	1.52
2	A	901	OT8	C26-C21	2.31	1.59	1.51
2	A	901	OT8	C2-C1	-2.12	1.40	1.42

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	OT8	C1-C2-C4	7.74	123.19	120.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	0T8	N3-C5-N1	-7.01	119.64	126.42
2	A	901	0T8	C9-N4-C4	4.31	131.86	122.11
2	A	901	0T8	C3-N3-C5	3.59	120.73	115.81
2	A	901	0T8	C3-C2-C1	3.10	117.79	114.67
2	A	901	0T8	C2-C1-N1	-2.92	119.10	122.65
2	A	901	0T8	C1-N2-C6	-2.86	119.34	124.18
2	A	901	0T8	C18-C22-N7	-2.56	119.23	123.60
2	A	901	0T8	C24-C20-N6	2.55	111.47	108.71
2	A	901	0T8	C2-C3-N3	-2.34	121.02	124.32
2	A	901	0T8	C21-N6-C20	2.30	116.73	113.76
2	A	901	0T8	C23-N7-C22	2.23	121.95	116.86
2	A	901	0T8	C5-N1-C1	2.18	121.71	116.36
2	A	901	0T8	N5-C5-N3	2.05	119.97	117.21

There are no chirality outliers.


All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	901	0T8	C9-C13-C15-N6
2	A	901	0T8	C3-C2-C4-O1
2	A	901	0T8	C13-C15-N6-C21
2	A	901	0T8	C3-C2-C4-N4
2	A	901	0T8	C13-C15-N6-C20
2	A	901	0T8	N3-C5-N5-C10
2	A	901	0T8	N1-C5-N5-C10
2	A	901	0T8	C13-C9-N4-C4

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/323 (69%)	1.55	80 (35%) ⓘ ⓘ	24, 49, 84, 89	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	783	LEU	6.4
1	A	724	ALA	4.8
1	A	749	GLU	4.6
1	A	555	PHE	4.6
1	A	508	PHE	4.6
1	A	725	GLY	4.6
1	A	556	LEU	4.5
1	A	511	GLY	4.5
1	A	534	PHE	4.2
1	A	610	ILE	4.0
1	A	558	GLU	3.9
1	A	541	MET	3.8
1	A	542	LEU	3.7
1	A	525	ALA	3.7
1	A	537	VAL	3.5
1	A	514	LEU	3.5
1	A	526	GLN	3.5
1	A	515	GLY	3.4
1	A	512	ARG	3.4
1	A	576	VAL	3.4
1	A	618	PRO	3.3
1	A	516	LYS	3.3
1	A	562	MET	3.2
1	A	782	HIS	3.2
1	A	520	GLY	3.2
1	A	510	LEU	3.2
1	A	639	ARG	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	597	LYS	3.1
1	A	574	VAL	3.0
1	A	619	LEU	3.0
1	A	577	SER	3.0
1	A	589	PRO	3.0
1	A	533	SER	3.0
1	A	665	GLY	2.9
1	A	765	GLN	2.9
1	A	535	VAL	2.9
1	A	557	ARG	2.9
1	A	776	LEU	2.9
1	A	608	SER	2.7
1	A	524	GLU	2.7
1	A	720	GLN	2.7
1	A	527	LEU	2.7
1	A	592	ILE	2.7
1	A	779	ILE	2.7
1	A	522	VAL	2.6
1	A	523	ARG	2.6
1	A	575	GLY	2.6
1	A	559	ALA	2.6
1	A	560	ALA	2.6
1	A	778	ASN	2.5
1	A	538	ALA	2.5
1	A	751	VAL	2.5
1	A	737	GLY	2.5
1	A	609	ARG	2.5
1	A	780	LEU	2.4
1	A	734	TYR	2.4
1	A	561	CYS	2.4
1	A	745	PRO	2.3
1	A	536	LYS	2.3
1	A	513	MET	2.2
1	A	532	GLY	2.2
1	A	773	ARG	2.2
1	A	528	LYS	2.2
1	A	539	VAL	2.2
1	A	624	ARG	2.2
1	A	564	GLU	2.2
1	A	727	GLU	2.2
1	A	777	GLU	2.2
1	A	521	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	590	MET	2.1
1	A	655	GLU	2.1
1	A	726	ILE	2.1
1	A	730	GLU	2.1
1	A	732	TYR	2.1
1	A	621	THR	2.0
1	A	716	MET	2.0
1	A	505	GLU	2.0
1	A	775	GLU	2.0
1	A	593	LEU	2.0
1	A	739	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	OT8	A	901	33/33	0.87	0.17	47,51,63,65	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OT8 A 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.