



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:37 PM UTC

PDB ID : 9FE8 / pdb_00009fe8
Title : Crystal Structure of reduced NuoEF variant P228R(NuoF) from Aquifex aeolicus
Authors : Wohlwend, D.; Friedrich, T.; Goeppert-Asadollahpour, S.
Deposited on : 2024-05-17
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	
1	C	160	
2	B	434	
2	D	434	

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 9829 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH-quinone oxidoreductase subunit E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	156	Total	C	N	O	S	0	0	0
			1268	821	204	234	9			
1	C	159	Total	C	N	O	S	0	1	0
			1296	837	211	239	9			

- Molecule 2 is a protein called NADH-quinone oxidoreductase subunit F.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	418	Total	C	N	O	S	0	1	0
			3298	2118	551	616	13			
2	D	418	Total	C	N	O	S	0	3	0
			3323	2136	555	619	13			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	228	ARG	PRO	engineered mutation	UNP O66841
B	427	ALA	-	expression tag	UNP O66841
B	428	GLY	-	expression tag	UNP O66841
B	429	HIS	-	expression tag	UNP O66841
B	430	HIS	-	expression tag	UNP O66841
B	431	HIS	-	expression tag	UNP O66841
B	432	HIS	-	expression tag	UNP O66841
B	433	HIS	-	expression tag	UNP O66841
B	434	HIS	-	expression tag	UNP O66841
D	228	ARG	PRO	engineered mutation	UNP O66841
D	427	ALA	-	expression tag	UNP O66841
D	428	GLY	-	expression tag	UNP O66841
D	429	HIS	-	expression tag	UNP O66841
D	430	HIS	-	expression tag	UNP O66841
D	431	HIS	-	expression tag	UNP O66841
D	432	HIS	-	expression tag	UNP O66841

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	433	HIS	-	expression tag	UNP O66841
D	434	HIS	-	expression tag	UNP O66841

- Molecule 3 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Fe	S	0	0
			4	2	2		
3	C	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O_4S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Na	0	0
			3	3		
5	B	3	Total	Na	0	0
			3	3		
5	C	1	Total	Na	0	0
			1	1		
5	D	1	Total	Na	0	0
			1	1		

- Molecule 6 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	Fe	S	0	0
			8	4	4		
6	D	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 7 is 1-DEOXY-1-(7,8-DIMETHYL-2,4-DIOXO-3,4-DIHYDRO-2H-BENZO[G]PT ERIDIN-1-ID-10(5H)-YL)-5-O-PHOSPHONATO-D-RIBITOL (CCD ID: FNR) (formula: C₁₇H₂₃N₄O₉P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	N	O	P	0
			31	17	4	9	1	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	D	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			6	3	3		
8	B	1	Total	C	O	0	0
			6	3	3		
8	B	1	Total	C	O	0	0
			6	3	3		
8	D	1	Total	C	O	0	0
			6	3	3		
8	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is 3[N-MORPHOLINO]PROPANE SULFONIC ACID (CCD ID: MPO) (formula: $C_7H_{15}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	D	1	Total	C	N	O	S	0	0
			13	7	1	4	1		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	52	Total	O	0	1
			53	53		
10	B	198	Total	O	0	0
			198	198		
10	C	55	Total	O	0	0
			55	55		
10	D	171	Total	O	0	0
			171	171		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NADH-quinone oxidoreductase subunit E

Chain A: 




- Molecule 1: NADH-quinone oxidoreductase subunit E

Chain C: 




- Molecule 2: NADH-quinone oxidoreductase subunit F

Chain B: 



- Molecule 2: NADH-quinone oxidoreductase subunit F

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.08Å 116.22Å 190.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.64 – 2.34 49.64 – 2.34	Depositor EDS
% Data completeness (in resolution range)	99.4 (49.64-2.34) 99.4 (49.64-2.34)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.184 , 0.206 (Not available) , 0.194	Depositor DCC
R_{free} test set	2968 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	38.2	Xtriage
Anisotropy	0.290	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 32.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9829	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.85 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6611e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FNR, SF4, SO4, MPO, GOL, FES, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/1297	0.99	0/1752
1	C	0.54	0/1325	1.02	0/1790
2	B	0.55	0/3378	1.00	1/4571 (0.0%)
2	D	0.55	0/3404	0.98	5/4603 (0.1%)
All	All	0.54	0/9404	0.99	6/12716 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	D	0	1
All	All	0	2

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	376	ASP	CA-CB-CG	5.83	118.43	112.60
2	D	110	ASP	CA-CB-CG	5.42	118.02	112.60
2	D	383	VAL	N-CA-CB	5.32	118.57	110.58
2	D	245	ASP	CA-CB-CG	5.14	117.74	112.60
2	B	298	ASP	CA-CB-CG	5.12	117.72	112.60
2	D	129	GLU	CB-CA-C	5.08	117.96	110.14

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	401	ARG	Sidechain
2	D	87	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1268	0	1269	19	0
1	C	1296	0	1292	5	0
2	B	3298	0	3259	19	0
2	D	3323	0	3286	20	0
3	A	4	0	0	0	0
3	C	4	0	0	0	0
4	A	10	0	0	0	0
4	B	5	0	0	0	0
4	C	10	0	0	0	0
4	D	5	0	0	0	0
5	A	3	0	0	0	0
5	B	3	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	B	8	0	0	0	0
6	D	8	0	0	1	0
7	B	31	0	22	0	0
7	D	31	0	22	0	0
8	B	18	0	24	2	0
8	D	12	0	16	0	0
9	D	13	0	15	0	0
10	A	53	0	0	1	0
10	B	198	0	0	1	0
10	C	55	0	0	0	0
10	D	171	0	0	1	0
All	All	9829	0	9205	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:384:ASN:HD22	2:B:404:ARG:HH21	1.06	1.00
1:A:38:GLN:HE22	1:A:74:ASP:H	1.31	0.77
2:B:68:GLY:HA3	8:B:504:GOL:H31	1.75	0.68
2:B:384:ASN:ND2	2:B:404:ARG:HH21	1.87	0.67
2:B:233:MET:HE1	2:B:241:ILE:HD11	1.78	0.66
2:B:384:ASN:HD22	2:B:404:ARG:NH2	1.88	0.64
2:B:208:GLN:HB2	10:B:772:HOH:O	2.02	0.59
2:D:94:ASP:O	2:D:95:GLU:C	2.47	0.57
2:B:106:ILE:HD11	2:B:251:LEU:HD11	1.88	0.55
1:A:62:HIS:HE1	2:B:194:GLY:O	1.89	0.55
2:B:25:LYS:HG2	8:B:503:GOL:H2	1.90	0.54
1:A:133:GLU:HG3	2:B:8:PRO:HB2	1.89	0.53
1:A:38:GLN:HE21	1:A:44:ILE:HG12	1.74	0.53
2:D:233:MET:HE1	2:D:241:ILE:HD11	1.88	0.53
2:B:94:ASP:O	2:B:95:GLU:C	2.52	0.51
1:C:25:LYS:HD2	2:D:212:TRP:CZ2	2.46	0.50
2:D:360:GLN:NE2	2:D:384:ASN:HD22	2.09	0.50
1:A:133:GLU:HB3	1:A:145:LYS:HB2	1.94	0.49
2:D:66:ARG:HB2	2:D:220:ASN:HD22	1.77	0.49
1:A:45:PRO:HB2	1:A:48:SER:HB3	1.93	0.49
2:D:237:GLU:HG2	10:D:731:HOH:O	2.13	0.48
1:A:49:LEU:HG	1:A:63:VAL:HG12	1.96	0.48
2:D:253:PRO:HG2	2:D:321:THR:HA	1.94	0.48
1:A:50:LYS:HB3	1:A:51:PRO:CD	2.44	0.48
1:A:110:LYS:HB3	1:A:111:PRO:HD2	1.97	0.47
1:A:62:HIS:HD2	10:A:347:HOH:O	1.99	0.46
2:D:2:ARG:NE	2:D:147:ASP:OD2	2.48	0.46
1:C:62:HIS:HE1	2:D:194:GLY:O	1.99	0.46
2:D:391:SER:HB2	6:D:501:SF4:S2	2.55	0.46
1:A:50:LYS:HB3	1:A:51:PRO:HD3	1.97	0.46
1:A:23:PHE:HE2	1:A:32:LEU:HD12	1.81	0.45
1:A:16:LEU:HD13	1:A:55:MET:HE1	1.98	0.45
1:A:91:CYS:HA	1:A:134:ALA:HB1	1.99	0.45
1:A:110:LYS:HB3	1:A:111:PRO:CD	2.47	0.45
2:D:211:LEU:HG	2:D:212:TRP:CD1	2.53	0.44
2:D:351:THR:HB	2:D:352:PRO:HD3	1.99	0.44
2:B:129:GLU:HG3	2:B:172:TYR:HE2	1.82	0.44
2:D:50:SER:OG	2:D:53:GLU:HG3	2.17	0.44
2:B:388[B]:GLN:N	2:B:389:PRO:CD	2.82	0.43
1:A:8:PHE:CE2	1:A:55:MET:HE2	2.53	0.43
1:A:38:GLN:NE2	1:A:44:ILE:HG12	2.33	0.43
2:D:154:LYS:HD2	2:D:155:LYS:HG2	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:52:GLU:H	2:B:52:GLU:CD	2.25	0.43
2:B:388[A]:GLN:N	2:B:389:PRO:CD	2.82	0.43
2:D:202:LYS:HA	2:D:203:PRO:C	2.44	0.43
2:D:205:TYR:O	2:D:208:GLN:HG2	2.19	0.43
2:D:252:PHE:CE1	2:D:309:MET:HB3	2.54	0.43
2:B:89:PHE:O	2:B:130:ALA:HA	2.18	0.42
1:C:84:ARG:O	1:C:137:PHE:HA	2.20	0.42
2:D:348:GLY:HA2	2:D:354[B]:ARG:HB2	2.02	0.42
2:B:348:GLY:HA2	2:B:354:ARG:HB2	2.02	0.41
2:B:17:ASN:OD1	2:B:21:LYS:HA	2.20	0.41
1:C:50:LYS:HB3	1:C:51:PRO:HD3	2.02	0.41
1:A:15:LYS:HD2	1:A:15:LYS:HA	1.94	0.41
1:C:41:TYR:O	1:C:78:LYS:HE2	2.20	0.41
2:D:402:LEU:C	2:D:402:LEU:HD23	2.45	0.41
2:D:64:ARG:HE	2:D:311[A]:TYR:HD2	1.68	0.41
1:A:90:VAL:HG11	2:B:255:SER:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	154/160 (96%)	149 (97%)	5 (3%)	0	100	100
1	C	158/160 (99%)	154 (98%)	4 (2%)	0	100	100
2	B	417/434 (96%)	405 (97%)	11 (3%)	1 (0%)	43	50
2	D	419/434 (96%)	406 (97%)	12 (3%)	1 (0%)	43	50
All	All	1148/1188 (97%)	1114 (97%)	32 (3%)	2 (0%)	43	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	95	GLU
2	B	95	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/146 (97%)	141 (99%)	1 (1%)	76	84
1	C	144/146 (99%)	144 (100%)	0	100	100
2	B	342/357 (96%)	339 (99%)	3 (1%)	70	80
2	D	344/357 (96%)	343 (100%)	1 (0%)	86	91
All	All	972/1006 (97%)	967 (100%)	5 (0%)	81	88

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	LEU
2	B	52	GLU
2	B	355	VAL
2	B	393	CYS
2	D	393	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	ASN
1	A	38	GLN
1	A	62	HIS
2	B	48	ASN
2	B	83	ASN
2	B	220	ASN
2	B	349	GLN
2	B	360	GLN
2	B	375	GLN
2	B	384	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	62	HIS
2	D	220	ASN
2	D	344	HIS
2	D	349	GLN
2	D	360	GLN
2	D	362	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 8 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	202	5	4,4,4	0.30	0	6,6,6	0.09	0
4	SO4	B	506	-	4,4,4	0.30	0	6,6,6	0.12	0
7	FNR	B	502	-	32,33,33	0.37	0	41,50,50	0.63	0
4	SO4	D	506	-	4,4,4	0.22	0	6,6,6	0.15	0
6	SF4	D	501	2	0,12,12	-	-	-		
8	GOL	B	505	-	5,5,5	0.13	0	5,5,5	0.34	0
4	SO4	C	203	-	4,4,4	0.32	0	6,6,6	0.22	0
7	FNR	D	502	-	32,33,33	0.38	0	41,50,50	0.57	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	GOL	D	505	-	5,5,5	0.10	0	5,5,5	0.28	0
9	MPO	D	503	-	13,13,13	0.73	1 (7%)	17,17,17	0.69	0
4	SO4	C	202	-	4,4,4	0.29	0	6,6,6	0.08	0
8	GOL	B	504	-	5,5,5	0.18	0	5,5,5	0.59	0
4	SO4	A	203	-	4,4,4	0.28	0	6,6,6	0.10	0
6	SF4	B	501	2	0,12,12	-	-	-		
3	FES	C	201	1	0,4,4	-	-	-		
8	GOL	D	504	-	5,5,5	0.10	0	5,5,5	0.35	0
3	FES	A	201	1	0,4,4	-	-	-		
8	GOL	B	503	-	5,5,5	0.18	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	FNR	B	502	-	-	4/18/18/18	0/3/3/3
8	GOL	D	505	-	-	0/4/4/4	-
6	SF4	D	501	2	-	-	0/6/5/5
8	GOL	B	505	-	-	0/4/4/4	-
7	FNR	D	502	-	-	3/18/18/18	0/3/3/3
9	MPO	D	503	-	-	0/7/15/15	0/1/1/1
8	GOL	B	504	-	-	2/4/4/4	-
6	SF4	B	501	2	-	-	0/6/5/5
3	FES	C	201	1	-	-	0/1/1/1
8	GOL	D	504	-	-	0/4/4/4	-
3	FES	A	201	1	-	-	0/1/1/1
8	GOL	B	503	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	D	503	MPO	O3-S1	2.54	1.56	1.47

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	B	504	GOL	C1-C2-C3-O3
8	B	504	GOL	O2-C2-C3-O3
8	B	503	GOL	O1-C1-C2-C3
7	D	502	FNR	C4'-C5'-O5'-P
7	B	502	FNR	C4'-C5'-O5'-P
8	B	503	GOL	O1-C1-C2-O2
7	B	502	FNR	O2'-C2'-C3'-C4'
7	D	502	FNR	O2'-C2'-C3'-C4'
7	B	502	FNR	O2'-C2'-C3'-O3'
7	B	502	FNR	N10-C1'-C2'-O2'
7	D	502	FNR	N10-C1'-C2'-O2'


There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	501	SF4	1	0
8	B	504	GOL	1	0
8	B	503	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand FNR B 502




Ligand SF4 D 501




Ligand FNR D 502




Ligand SF4 B 501



Ligand FES C 201





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	156/160 (97%)	0.24	0 100 100	33, 56, 79, 112	0
1	C	159/160 (99%)	-0.01	2 (1%) 75 79	21, 46, 75, 115	1 (0%)
2	B	418/434 (96%)	-0.11	3 (0%) 84 87	19, 40, 62, 90	1 (0%)
2	D	418/434 (96%)	-0.13	5 (1%) 76 80	18, 41, 59, 95	3 (0%)
All	All	1151/1188 (96%)	-0.06	10 (0%) 81 84	18, 42, 68, 115	5 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	311[A]	TYR	5.7
1	C	2	PHE	4.2
2	B	419	LYS	3.0
2	D	419	LYS	2.9
2	D	2	ARG	2.5
2	B	2	ARG	2.4
1	C	6	PHE	2.4
2	D	416	LYS	2.3
2	D	240	TYR	2.1
2	B	246	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	B	506	5/5	0.62	0.09	94,97,102,106	0
4	SO4	A	203	5/5	0.75	0.10	74,76,79,80	0
4	SO4	A	202	5/5	0.76	0.14	78,87,97,99	0
4	SO4	C	202	5/5	0.77	0.09	75,77,88,90	0
8	GOL	B	503	6/6	0.81	0.13	46,50,53,54	0
8	GOL	B	504	6/6	0.81	0.27	72,81,85,86	0
8	GOL	D	504	6/6	0.83	0.17	55,62,68,76	0
8	GOL	B	505	6/6	0.86	0.18	64,75,79,80	0
9	MPO	D	503	13/13	0.86	0.19	58,78,91,94	0
4	SO4	C	203	5/5	0.87	0.15	59,67,71,74	0
5	NA	A	205	1/1	0.90	0.10	59,59,59,59	0
5	NA	A	204	1/1	0.90	0.18	57,57,57,57	0
5	NA	C	204	1/1	0.91	0.15	53,53,53,53	0
5	NA	B	509	1/1	0.91	0.20	59,59,59,59	0
5	NA	D	507	1/1	0.92	0.21	43,43,43,43	0
4	SO4	D	506	5/5	0.92	0.11	57,62,71,81	0
5	NA	B	508	1/1	0.92	0.24	58,58,58,58	0
8	GOL	D	505	6/6	0.93	0.18	68,69,72,74	0
5	NA	A	206	1/1	0.95	0.15	47,47,47,47	0
5	NA	B	507	1/1	0.96	0.05	45,45,45,45	0
7	FNR	B	502	31/31	0.96	0.06	28,32,34,37	0
7	FNR	D	502	31/31	0.96	0.07	30,33,34,36	0
6	SF4	D	501	8/8	0.99	0.02	31,33,34,34	0
3	FES	A	201	4/4	0.99	0.03	33,33,33,35	0
6	SF4	B	501	8/8	0.99	0.03	33,34,35,36	0
3	FES	C	201	4/4	1.00	0.02	32,32,32,33	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FNR B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FNR D 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SF4 D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FES A 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SF4 B 501:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FES C 201:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.