



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:25 AM UTC

PDB ID : 5FFI / pdb_00005ffi
Title : [2Fe:2S] ferredoxin FeSII from Azotobacter vinelandii
Authors : Schlesier, J.; Rohde, M.; Gerhardt, S.; Einsle, O.
Deposited on : 2015-12-18
Resolution : 2.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8975 (2.20-2.16)
Clashscore	190562	9786 (2.20-2.16)
Ramachandran outliers	187476	9664 (2.20-2.16)
Sidechain outliers	187428	9664 (2.20-2.16)
RSRZ outliers	180081	8979 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	122	6% 86% 9% • •
1	B	122	7% 82% 13% 5%
1	C	122	8% 82% 12% • 5%
1	D	122	17% 84% 10% • •
1	E	122	11% 75% 7% • 16%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4529 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dimeric (2Fe-2S) protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	118	Total	C	N	O	S	0	0	0
			905	567	158	174	6			
1	B	116	Total	C	N	O	S	0	0	0
			885	555	152	172	6			
1	C	116	Total	C	N	O	S	0	0	0
			894	560	156	172	6			
1	D	117	Total	C	N	O	S	0	0	0
			897	561	157	173	6			
1	E	102	Total	C	N	O	S	0	0	0
			776	490	134	147	5			

- Molecule 2 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe₂S₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			4	2	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total 4	Fe 2	S 2	0	0
2	C	1	Total 4	Fe 2	S 2	0	0
2	D	1	Total 4	Fe 2	S 2	0	0
2	E	1	Total 4	Fe 2	S 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	31	Total 31	O 31	0	0
3	B	37	Total 37	O 37	0	0
3	C	16	Total 16	O 16	0	0
3	D	26	Total 26	O 26	0	0
3	E	42	Total 42	O 42	0	0

- Molecule 1: Dimeric (2Fe-2S) protein



-
- | Category | Count |
|----------|-------|
| MET | 1 |
| A2 | 1 |
| L10 | 1 |
| M14 | 1 |
| Q18 | 1 |
| A19 | 1 |
| V20 | 1 |
| A21 | 1 |
| G22 | 1 |
| LYS | 1 |
| ARG | 1 |
| T26 | 1 |
| T27 | 1 |
| F40 | 1 |
| I54 | 1 |
| I62 | 1 |
| K63 | 1 |
| G64 | 1 |
| M65 | 1 |
| K83 | 1 |
| A89 | 1 |
| A90 | 1 |
| V91 | 1 |
| R92 | 1 |
| D93 | 1 |
| L94 | 1 |
| R99 | 1 |
| D108 | 1 |
| P119 | 1 |
| GLY | 1 |
| ALA | 1 |

- | | | | | | | | | | | | |
|-----|----|------|------|------|------|------|------|-----|-----|-----|-----|
| Met | A2 | M10 | M11 | P12 | Q18 | A19 | V20 | A21 | GLY | K23 | L27 | L28 | K37 | F40 | T54 | T55 | I62 | T68 | D69 | K76 | L81 | A89 | L94 | P95 | R99 | I105 | V106 | T107 | D108 | L112 | E118 | PRO | GLY | GLY | ALA |
|-----|----|------|------|------|------|------|------|-----|-----|-----|-----|

-
- | Category | Count |
|----------|-------|
| MET | 1 |
| A2 | 2 |
| L10 | 3 |
| M11 | 2 |
| N14 | 2 |
| A21 | 3 |
| K23 | 3 |
| R24 | 3 |
| L27 | 2 |
| L28 | 2 |
| F40 | 2 |
| I54 | 2 |
| L62 | 3 |
| K65 | 3 |
| L66 | 3 |
| L67 | 3 |
| T68 | 2 |
| D69 | 2 |
| V74 | 2 |
| V78 | 3 |
| G79 | 2 |
| K80 | 3 |
| L81 | 2 |
| S84 | 3 |
| E85 | 3 |
| R88 | 3 |
| K89 | 3 |
| VAL | 2 |
| VAL | 2 |
| R92 | 3 |
| D93 | 3 |
| L94 | 3 |
| P95 | 3 |
| R99 | 2 |
| D108 | 2 |
| G120 | 3 |
| GLY | 2 |
| ALA | 2 |

-

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	133.27Å 134.48Å 36.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	94.66 – 2.17 94.66 – 2.17	Depositor EDS
% Data completeness (in resolution range)	100.0 (94.66-2.17) 100.0 (94.66-2.17)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.16Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.195 , 0.229 0.208 , 0.241	Depositor DCC
R_{free} test set	1687 reflections (4.68%)	wwPDB-VP
Wilson B-factor (Å ²)	39.3	Xtriage
Anisotropy	0.366	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 61.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4529	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/918	1.18	2/1239 (0.2%)
1	B	0.77	0/897	1.21	2/1211 (0.2%)
1	C	0.70	0/905	1.16	0/1219
1	D	0.77	0/909	1.20	1/1224 (0.1%)
1	E	0.85	0/786	1.15	2/1062 (0.2%)
All	All	0.76	0/4415	1.18	7/5955 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	108	ASP	CA-C-N	5.58	128.76	120.95
1	E	108	ASP	C-N-CA	5.58	128.76	120.95
1	B	91	VAL	CA-C-N	5.55	127.65	120.44
1	B	91	VAL	C-N-CA	5.55	127.65	120.44
1	D	95	PRO	N-CA-C	5.51	117.43	110.70
1	A	108	ASP	CA-C-N	5.46	128.51	120.82
1	A	108	ASP	C-N-CA	5.46	128.51	120.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	905	0	925	6	0
1	B	885	0	898	6	0
1	C	894	0	914	7	0
1	D	897	0	913	6	0
1	E	776	0	788	5	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
2	D	4	0	0	0	0
2	E	4	0	0	0	0
3	A	31	0	0	0	0
3	B	37	0	0	0	0
3	C	16	0	0	0	0
3	D	26	0	0	0	0
3	E	42	0	0	0	0
All	All	4529	0	4438	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:38:ILE:HG13	1:E:39:PRO:HD2	1.72	0.70
1:D:54:ILE:HD12	1:D:99:ARG:HG3	1.86	0.57
1:A:94:LEU:HD11	1:B:14:ASN:HD21	1.72	0.53
1:A:14:ASN:HD21	1:B:94:LEU:HD21	1.74	0.52
1:C:76:LYS:HG3	1:C:81:LEU:HD23	1.91	0.51
1:A:28:LEU:HD21	1:A:40:PHE:CD1	2.46	0.50
1:D:28:LEU:HD21	1:D:40:PHE:HB3	1.94	0.49
1:C:54:ILE:HD12	1:C:99:ARG:HG3	1.94	0.49
1:C:28:LEU:HD21	1:C:40:PHE:CD1	2.49	0.48
1:B:54:ILE:HD12	1:B:99:ARG:HG3	1.95	0.48
1:C:94:LEU:HD11	1:D:14:ASN:HD21	1.79	0.47
1:A:54:ILE:HD12	1:A:99:ARG:HG3	1.97	0.46
1:E:27:LEU:HD23	1:E:27:LEU:HA	1.84	0.45
1:A:62:ILE:HD13	1:A:89:ALA:HA	1.99	0.45
1:A:94:LEU:HD23	1:A:95:PRO:HD2	1.99	0.44
1:B:62:ILE:HD13	1:B:89:ALA:HA	2.00	0.44
1:D:10:LEU:HD12	1:D:11:MET:HG3	1.99	0.44
1:E:72:ARG:HG3	1:E:81:LEU:HD21	2.00	0.43
1:E:55:THR:HB	1:E:112:LEU:HB3	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:28:LEU:HD21	1:D:40:PHE:CB	2.49	0.42
1:C:62:ILE:HD13	1:C:89:ALA:HA	2.00	0.42
1:C:94:LEU:HD23	1:C:95:PRO:HD2	2.01	0.42
1:B:40:PHE:CE2	1:E:39:PRO:HD3	2.55	0.42
1:B:92:ARG:H	1:B:92:ARG:HG3	1.66	0.42
1:D:94:LEU:HD12	1:D:95:PRO:HD2	2.02	0.41
1:C:55:THR:HB	1:C:112:LEU:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	116/122 (95%)	115 (99%)	1 (1%)	0	100	100
1	B	112/122 (92%)	111 (99%)	1 (1%)	0	100	100
1	C	112/122 (92%)	110 (98%)	2 (2%)	0	100	100
1	D	113/122 (93%)	111 (98%)	2 (2%)	0	100	100
1	E	96/122 (79%)	93 (97%)	3 (3%)	0	100	100
All	All	549/610 (90%)	540 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/103 (99%)	97 (95%)	5 (5%)	22	26
1	B	100/103 (97%)	93 (93%)	7 (7%)	14	14
1	C	101/103 (98%)	96 (95%)	5 (5%)	22	25
1	D	101/103 (98%)	95 (94%)	6 (6%)	18	19
1	E	88/103 (85%)	83 (94%)	5 (6%)	18	20
All	All	492/515 (96%)	464 (94%)	28 (6%)	18	20

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	65	MET
1	A	83	LYS
1	A	94	LEU
1	A	108	ASP
1	B	18	GLN
1	B	20	VAL
1	B	25	SER
1	B	27	LEU
1	B	65	MET
1	B	83	LYS
1	B	108	ASP
1	C	18	GLN
1	C	27	LEU
1	C	37	LYS
1	C	94	LEU
1	C	108	ASP
1	D	10	LEU
1	D	27	LEU
1	D	65	MET
1	D	69	ASP
1	D	84	SER
1	D	108	ASP
1	E	15	LYS
1	E	23	LYS
1	E	43	GLN
1	E	81	LEU
1	E	108	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	HIS
1	A	14	ASN
1	B	14	ASN
1	B	32	GLN
1	B	43	GLN
1	C	103	GLN
1	D	13	HIS
1	D	14	ASN
1	D	18	GLN
1	D	103	GLN
1	E	14	ASN
1	E	32	GLN
1	E	46	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FES	B	1001	1	0,4,4	-	-	-		
2	FES	E	1001	1	0,4,4	-	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FES	D	1001	1	0,4,4	-	-	-		
2	FES	A	1001	1	0,4,4	-	-	-		
2	FES	C	1001	1	0,4,4	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FES	B	1001	1	-	-	0/1/1/1
2	FES	E	1001	1	-	-	0/1/1/1
2	FES	D	1001	1	-	-	0/1/1/1
2	FES	A	1001	1	-	-	0/1/1/1
2	FES	C	1001	1	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	118/122 (96%)	0.41	7 (5%) 28 27	29, 52, 80, 99	0
1	B	116/122 (95%)	0.43	8 (6%) 23 22	28, 49, 79, 101	0
1	C	116/122 (95%)	0.89	10 (8%) 16 15	37, 73, 111, 135	0
1	D	117/122 (95%)	0.80	21 (17%) 3 3	25, 56, 120, 136	0
1	E	102/122 (83%)	0.61	13 (12%) 8 7	27, 49, 83, 97	0
All	All	569/610 (93%)	0.63	59 (10%) 11 10	25, 56, 105, 136	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	22	GLY	5.9
1	D	94	LEU	4.2
1	C	21	ALA	4.2
1	D	89	ALA	4.1
1	E	82	PRO	4.1
1	E	81	LEU	3.9
1	D	120	GLY	3.8
1	B	40	PHE	3.8
1	E	23	LYS	3.7
1	D	67	LEU	3.6
1	D	78	VAL	3.6
1	E	94	LEU	3.6
1	D	22	GLY	3.6
1	E	62	ILE	3.4
1	E	20	VAL	3.4
1	A	65	MET	3.3
1	D	95	PRO	3.3
1	E	38	ILE	3.3
1	D	65	MET	3.3
1	B	21	ALA	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	40	PHE	3.1
1	E	66	LEU	3.1
1	A	66	LEU	3.0
1	A	119	PRO	3.0
1	C	20	VAL	2.9
1	E	116	THR	2.9
1	C	89	ALA	2.9
1	D	21	ALA	2.9
1	D	10	LEU	2.8
1	B	119	PRO	2.7
1	D	88	ARG	2.7
1	E	22	GLY	2.7
1	A	95	PRO	2.6
1	D	81	LEU	2.6
1	E	39	PRO	2.6
1	C	105	ILE	2.5
1	D	66	LEU	2.5
1	D	40	PHE	2.5
1	D	69	ASP	2.4
1	A	62	ILE	2.4
1	A	78	VAL	2.4
1	B	91	VAL	2.4
1	D	84	SER	2.4
1	D	80	LYS	2.3
1	A	79	GLY	2.3
1	C	10	LEU	2.3
1	C	12	PRO	2.2
1	E	37	LYS	2.2
1	C	68	THR	2.2
1	D	62	ILE	2.1
1	D	85	GLU	2.1
1	C	95	PRO	2.1
1	B	20	VAL	2.1
1	C	106	VAL	2.1
1	D	74	VAL	2.1
1	B	10	LEU	2.1
1	D	24	ARG	2.0
1	C	69	ASP	2.0
1	B	64	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FES	C	1001	4/4	0.98	0.04	80,85,85,88	0
2	FES	B	1001	4/4	0.99	0.03	33,36,37,38	0
2	FES	E	1001	4/4	0.99	0.03	37,39,43,43	0
2	FES	D	1001	4/4	1.00	0.01	29,30,30,31	0
2	FES	A	1001	4/4	1.00	0.02	30,34,34,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.