



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:07 PM UTC

PDB ID : 9FFJ / pdb_00009ffj
Title : Artificial metalloenzyme with a nickel-based 1,10-phenanthroline cofactor and streptavidin N49M-S112V mutant
Authors : Lau, K.; Wang, W.; Pojer, F.; Larabi, A.
Deposited on : 2024-05-23
Resolution : 1.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2836 (1.30-1.26)
Clashscore	190562	2911 (1.30-1.26)
Ramachandran outliers	187476	2841 (1.30-1.26)
Sidechain outliers	187428	2840 (1.30-1.26)
RSRZ outliers	180081	2832 (1.30-1.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 2116 atoms, of which 980 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Streptavidin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	121	Total	C	H	N	O	S	0	20	0
			1937	620	944	171	200	2			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P22629
A	2	ALA	-	expression tag	UNP P22629
A	3	SER	-	expression tag	UNP P22629
A	4	MET	-	expression tag	UNP P22629
A	5	THR	-	expression tag	UNP P22629
A	6	GLY	-	expression tag	UNP P22629
A	7	GLY	-	expression tag	UNP P22629
A	8	GLN	-	expression tag	UNP P22629
A	9	GLN	-	expression tag	UNP P22629
A	10	MET	-	expression tag	UNP P22629
A	11	GLY	-	expression tag	UNP P22629
A	12	ARG	-	expression tag	UNP P22629
A	13	ASP	-	expression tag	UNP P22629
A	14	GLN	-	expression tag	UNP P22629
A	49	MET	ASN	engineered mutation	UNP P22629
A	112	VAL	SER	engineered mutation	UNP P22629

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 3 is N-methyl-N-[(4,4,6,6-tetrahydroxy-4,6-dioxido-1,3,3a,5,6a-tetrahydrothien[3,4-d]imidazol-4-ium-2-yl)methyl]-5-(2,4,4-trihydroxy-2-keto-3,3a,5,6-tetrahydro-1H-thien[3,4-d]imidazol-4-ium-6-yl) (CCD ID: A1ICD) (formula: $C_{24}H_{29}N_5NiO_6S$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	Ni	O	S	0	1
			74	48	10	2	12	2		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			17	4	10	3		
4	A	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is AMMONIUM ION (CCD ID: NH4) (formula: H_4N).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	H	N	0	0
			5	4	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	36	Total	O	0	1
			36	36		

- Molecule 1: Streptavidin



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	57.46Å 57.46Å 172.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	54.51 – 1.27 54.51 – 1.27	Depositor EDS
% Data completeness (in resolution range)	97.6 (54.51-1.27) 93.2 (54.51-1.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 1.28Å)	Xtriage
Refinement program	PHENIX dev_5246	Depositor
R, R_{free}	0.177 , 0.197 0.179 , 0.196	Depositor DCC
R_{free} test set	1859 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	17.0	Xtriage
Anisotropy	0.211	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 44.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2116	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PEG, A1ICD, NH4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1085	0.62	0/1479

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	993	944	874	11	0
2	A	8	12	12	0	0
3	A	74	0	0	0	0
4	A	14	20	20	1	0
5	A	10	0	0	0	0
6	A	1	4	0	0	0
7	A	36	0	0	0	0
All	All	1136	980	906	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103:ARG:HD3	1:A:131:THR:HG22	1.82	0.61
1:A:119:ALA:HB1	4:A:203:PEG:O2	2.04	0.57
1:A:49[A]:MET:HG2	1:A:86:ALA:HB2	1.98	0.45
1:A:51:GLU:OE2	1:A:52[B]:SER:OG	2.35	0.44
1:A:29[A]:PHE:HB3	1:A:43:TYR:CD2	2.53	0.43
1:A:110[A]:LEU:HD23	1:A:110[A]:LEU:C	2.43	0.43
1:A:73:LEU:HD12	1:A:73:LEU:C	2.45	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/159 (86%)	132 (96%)	5 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/115 (92%)	106 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	HIS
1	A	127	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 1 is modelled with single atom - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	A1ICD	A	202[A]	-	42,42,42	7.51	32 (76%)	51,70,70	2.99	23 (45%)
5	SO4	A	207	-	4,4,4	0.67	0	6,6,6	0.08	0
2	EDO	A	201	-	3,3,3	0.25	0	2,2,2	0.32	0
5	SO4	A	206	-	4,4,4	0.66	0	6,6,6	0.09	0
4	PEG	A	204	-	6,6,6	0.25	0	5,5,5	0.25	0
2	EDO	A	205	-	3,3,3	0.24	0	2,2,2	0.29	0
4	PEG	A	203	-	6,6,6	0.28	0	5,5,5	0.46	0
3	A1ICD	A	202[B]	-	42,42,42	8.33	33 (78%)	51,70,70	2.56	17 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1ICD	A	202[A]	-	-	3/15/56/56	0/6/6/6
2	EDO	A	201	-	-	0/1/1/1	-
4	PEG	A	204	-	-	3/4/4/4	-
2	EDO	A	205	-	-	1/1/1/1	-
4	PEG	A	203	-	-	1/4/4/4	-
3	A1ICD	A	202[B]	-	-	6/15/56/56	0/6/6/6

All (65) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	202[B]	A1ICD	C09-S07	-31.26	1.33	1.82
3	A	202[A]	A1ICD	C09-S07	-26.44	1.40	1.82
3	A	202[B]	A1ICD	C14-N15	17.47	1.55	1.34
3	A	202[A]	A1ICD	C14-N15	15.61	1.53	1.34
3	A	202[B]	A1ICD	C04-N03	-14.82	1.18	1.45
3	A	202[B]	A1ICD	C06-C05	-14.42	1.22	1.53
3	A	202[A]	A1ICD	C04-N03	-13.58	1.20	1.45
3	A	202[A]	A1ICD	C06-C05	-13.20	1.25	1.53
3	A	202[B]	A1ICD	C05-N08	-12.99	1.26	1.46
3	A	202[B]	A1ICD	C02-N03	12.76	1.58	1.35
3	A	202[A]	A1ICD	C02-N03	12.75	1.58	1.35
3	A	202[B]	A1ICD	C36-N15	-11.52	1.05	1.45
3	A	202[A]	A1ICD	C36-N15	-11.30	1.06	1.45
3	A	202[A]	A1ICD	C05-N08	-11.02	1.29	1.46
3	A	202[B]	A1ICD	C19-N20	10.42	1.49	1.33
3	A	202[A]	A1ICD	C19-N20	9.95	1.48	1.33
3	A	202[B]	A1ICD	C27-C26	9.35	1.58	1.41
3	A	202[A]	A1ICD	C27-C26	8.57	1.56	1.41
3	A	202[A]	A1ICD	C30-C31	7.66	1.53	1.41
3	A	202[B]	A1ICD	C30-C31	7.56	1.53	1.41
3	A	202[A]	A1ICD	C33-N32	7.35	1.44	1.33
3	A	202[B]	A1ICD	C33-N32	7.20	1.44	1.33
3	A	202[B]	A1ICD	C02-N08	6.92	1.48	1.35
3	A	202[A]	A1ICD	NI21-N32	6.73	2.21	2.07
3	A	202[B]	A1ICD	NI21-N32	6.57	2.20	2.07
3	A	202[A]	A1ICD	C02-N08	6.35	1.46	1.35
3	A	202[B]	A1ICD	C13-C14	6.20	1.64	1.51
3	A	202[B]	A1ICD	C09-C04	-5.63	1.38	1.53
3	A	202[A]	A1ICD	C13-C14	5.58	1.63	1.51
3	A	202[B]	A1ICD	C19-C18	5.34	1.49	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	202[B]	A1ICD	C16-C17	5.08	1.64	1.52
3	A	202[A]	A1ICD	C33-C34	5.08	1.49	1.38
3	A	202[B]	A1ICD	C33-C34	5.02	1.49	1.38
3	A	202[A]	A1ICD	C19-C18	4.90	1.48	1.38
3	A	202[B]	A1ICD	NI21-N20	4.87	2.17	2.07
3	A	202[B]	A1ICD	C05-C04	4.72	1.69	1.55
3	A	202[B]	A1ICD	C29-C30	4.69	1.52	1.42
3	A	202[A]	A1ICD	C09-C04	-4.69	1.41	1.53
3	A	202[A]	A1ICD	C29-C30	4.63	1.52	1.42
3	A	202[B]	A1ICD	C06-S07	-4.58	1.68	1.81
3	A	202[A]	A1ICD	C05-C04	4.56	1.68	1.55
3	A	202[A]	A1ICD	C16-C17	4.28	1.62	1.52
3	A	202[A]	A1ICD	NI21-N20	3.99	2.15	2.07
3	A	202[B]	A1ICD	C18-C17	3.94	1.44	1.37
3	A	202[A]	A1ICD	C18-C17	3.88	1.44	1.37
3	A	202[B]	A1ICD	C16-N15	3.79	1.56	1.46
3	A	202[B]	A1ICD	C34-C35	3.66	1.44	1.36
3	A	202[A]	A1ICD	C34-C35	3.60	1.44	1.36
3	A	202[A]	A1ICD	C06-S07	-3.43	1.71	1.81
3	A	202[A]	A1ICD	C16-N15	3.25	1.55	1.46
3	A	202[B]	A1ICD	C12-C13	3.06	1.63	1.52
3	A	202[B]	A1ICD	C28-C27	2.79	1.47	1.42
3	A	202[A]	A1ICD	C12-C13	2.74	1.62	1.52
3	A	202[A]	A1ICD	C28-C27	2.74	1.47	1.42
3	A	202[B]	A1ICD	C17-C27	2.67	1.48	1.42
3	A	202[B]	A1ICD	C35-C30	2.65	1.47	1.42
3	A	202[A]	A1ICD	C35-C30	2.61	1.47	1.42
3	A	202[A]	A1ICD	C12-C11	2.51	1.64	1.51
3	A	202[B]	A1ICD	C31-N32	2.48	1.40	1.36
3	A	202[B]	A1ICD	C12-C11	2.47	1.64	1.51
3	A	202[B]	A1ICD	C11-C10	2.40	1.62	1.52
3	A	202[A]	A1ICD	C31-C26	-2.25	1.38	1.43
3	A	202[A]	A1ICD	C17-C27	2.12	1.47	1.42
3	A	202[B]	A1ICD	C31-C26	-2.09	1.39	1.43
3	A	202[A]	A1ICD	C11-C10	2.05	1.60	1.52

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	202[A]	A1ICD	C18-C19-N20	-8.32	119.72	123.26
3	A	202[B]	A1ICD	C06-S07-C09	8.03	106.69	89.98
3	A	202[A]	A1ICD	C17-C16-N15	-7.59	103.09	114.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	202[A]	A1ICD	C04-C09-S07	6.90	112.87	105.03
3	A	202[A]	A1ICD	C31-C26-N20	6.25	123.13	117.22
3	A	202[A]	A1ICD	C04-C05-N08	5.80	108.87	102.43
3	A	202[B]	A1ICD	C05-C04-N03	5.40	108.77	102.68
3	A	202[B]	A1ICD	C04-C05-N08	5.29	108.31	102.43
3	A	202[A]	A1ICD	C27-C26-N20	-4.98	118.23	122.80
3	A	202[B]	A1ICD	C17-C27-C26	4.91	120.36	117.56
3	A	202[B]	A1ICD	C18-C19-N20	-4.71	121.25	123.26
3	A	202[B]	A1ICD	C31-C26-N20	4.67	121.64	117.22
3	A	202[B]	A1ICD	C27-C26-N20	-4.58	118.60	122.80
3	A	202[A]	A1ICD	C19-N20-C26	4.53	122.88	118.01
3	A	202[A]	A1ICD	C05-C06-S07	4.43	112.16	106.06
3	A	202[A]	A1ICD	C06-S07-C09	4.04	98.39	89.98
3	A	202[B]	A1ICD	C26-C31-N32	3.93	120.94	117.22
3	A	202[B]	A1ICD	C04-N03-C02	-3.74	107.92	112.56
3	A	202[B]	A1ICD	C05-N08-C02	-3.35	107.46	112.38
3	A	202[A]	A1ICD	C09-C04-C05	-3.25	104.92	108.89
3	A	202[B]	A1ICD	C09-C04-N03	-3.21	109.95	113.34
3	A	202[A]	A1ICD	C26-C31-N32	2.95	120.01	117.22
3	A	202[A]	A1ICD	C05-C04-N03	2.88	105.93	102.68
3	A	202[A]	A1ICD	C05-N08-C02	-2.86	108.17	112.38
3	A	202[B]	A1ICD	C19-N20-C26	2.86	121.08	118.01
3	A	202[A]	A1ICD	C06-C05-C04	-2.85	104.56	109.06
3	A	202[A]	A1ICD	C06-C05-N08	-2.84	109.53	113.18
3	A	202[B]	A1ICD	C17-C16-N15	-2.75	110.36	114.49
3	A	202[A]	A1ICD	C10-C09-C04	-2.57	107.25	114.51
3	A	202[A]	A1ICD	N08-C02-N03	-2.51	105.94	108.85
3	A	202[A]	A1ICD	C12-C13-C14	-2.45	106.32	112.66
3	A	202[A]	A1ICD	C28-C27-C26	2.31	121.16	117.65
3	A	202[B]	A1ICD	C30-C31-N32	-2.30	120.18	122.81
3	A	202[A]	A1ICD	C17-C27-C26	2.27	118.86	117.56
3	A	202[B]	A1ICD	C04-C09-S07	2.19	107.52	105.03
3	A	202[A]	A1ICD	C30-C31-N32	-2.17	120.33	122.81
3	A	202[B]	A1ICD	C09-C04-C05	-2.16	106.25	108.89
3	A	202[A]	A1ICD	C33-N32-C31	2.06	120.22	118.01
3	A	202[B]	A1ICD	C33-N32-C31	2.04	120.20	118.01
3	A	202[A]	A1ICD	C11-C10-C09	-2.01	109.42	114.04

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	202[A]	A1ICD	C04-C09-C10-C11
3	A	202[A]	A1ICD	S07-C09-C10-C11
3	A	202[B]	A1ICD	C04-C09-C10-C11
3	A	202[B]	A1ICD	S07-C09-C10-C11
4	A	203	PEG	O2-C3-C4-O4
3	A	202[B]	A1ICD	C11-C12-C13-C14
4	A	204	PEG	O1-C1-C2-O2
4	A	204	PEG	O2-C3-C4-O4
4	A	204	PEG	C1-C2-O2-C3
3	A	202[B]	A1ICD	N15-C16-C17-C27
2	A	205	EDO	O1-C1-C2-O2
3	A	202[B]	A1ICD	N15-C16-C17-C18
3	A	202[A]	A1ICD	N15-C16-C17-C27
3	A	202[B]	A1ICD	C10-C11-C12-C13


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	203	PEG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand A1ICD A 202 (A)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	121/159 (76%)	0.63	21 (17%) 4 4	9, 23, 51, 67	12 (9%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	49[A]	MET	6.3
1	A	51	GLU	6.3
1	A	64	PRO	4.8
1	A	25	LEU	4.1
1	A	63	ALA	4.1
1	A	47	VAL	4.1
1	A	11	GLY	3.9
1	A	50[A]	ALA	3.9
1	A	46	ALA	3.8
1	A	45	SER	3.7
1	A	68	GLY	3.7
1	A	52[A]	SER	3.5
1	A	84	ARG	3.3
1	A	134	LYS	3.0
1	A	133	VAL	2.9
1	A	100	ALA	2.9
1	A	53[A]	ARG	2.6
1	A	69	SER	2.4
1	A	82	ASN	2.3
1	A	36	ASP	2.2
1	A	12	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SO4	A	207	5/5	0.70	0.24	117,117,118,118	0
5	SO4	A	206	5/5	0.79	0.27	103,103,103,103	5
4	PEG	A	203	7/7	0.80	0.23	48,62,73,74	0
4	PEG	A	204	7/7	0.81	0.23	49,60,71,72	17
2	EDO	A	201	4/4	0.82	0.14	70,84,85,85	10
2	EDO	A	205	4/4	0.84	0.30	67,81,85,87	0
3	A1ICD	A	202[A]	37/37	0.92	0.16	15,40,48,49	37
3	A1ICD	A	202[B]	37/37	0.92	0.16	61,89,107,108	37
6	NH4	A	208	1/1	0.97	0.06	23,27,27,27	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1ICD A 202 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around A1ICD A 202 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.