



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 06:14 PM UTC

PDB ID : 3FHT / pdb_00003fht
Title : Crystal structure of human Dbp5 in complex with AMPPNP and RNA
Authors : von Moeller, H.; Conti, E.
Deposited on : 2008-12-10
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)
RNA backbone	3983	1052 (2.50-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	412	
1	B	412	
2	C	10	
2	D	10	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6815 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent RNA helicase DDX19B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	87	1	0
			3116	1975	546	575	20			
1	B	387	Total	C	N	O	S	41	1	0
			3074	1949	536	569	20			


- Molecule 2 is a RNA chain called RNA (5'-R(*UP*UP*UP*UP*UP*UP*UP*UP*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	6	Total	C	N	O	P	0	0	0
			120	54	12	48	6			
2	D	6	Total	C	N	O	P	0	0	0
			117	54	12	46	5			

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
5	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	142	Total	O	0	0
			142	142		
6	B	147	Total	O	0	0
			147	147		
6	C	7	Total	O	0	0
			7	7		
6	D	4	Total	O	0	0
			4	4		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP-dependent RNA helicase DDX19B




- Molecule 1: ATP-dependent RNA helicase DDX19B



- Molecule 2: RNA (5'-R(*UP*UP*UP*UP*UP*UP*UP*UP*U)-3')



- Molecule 2: RNA (5'-R(*UP*UP*UP*UP*UP*UP*UP*UP*U)-3')





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	41.60Å 79.66Å 124.54Å 90.00° 90.02° 90.00°	Depositor
Resolution (Å)	41.52 – 2.20 41.52 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (41.52-2.20) 99.4 (41.52-2.20)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.32 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0063	Depositor
R, R_{free}	0.168 , 0.225 0.172 , 0.219	Depositor DCC
R_{free} test set	2061 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.181	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 29.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.468 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6815	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.17	5/3170 (0.2%)	1.11	8/4277 (0.2%)
1	B	1.16	5/3127 (0.2%)	1.11	4/4220 (0.1%)
2	C	0.96	0/131	1.40	1/200 (0.5%)
2	D	1.19	0/128	1.54	2/196 (1.0%)
All	All	1.16	10/6556 (0.2%)	1.12	15/8893 (0.2%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	291	ASP	C-O	-11.35	1.18	1.23
1	A	107	VAL	CA-CB	6.64	1.62	1.54
1	A	246	VAL	CA-CB	6.06	1.62	1.54
1	A	433	PHE	C-O	-6.01	1.16	1.23
1	A	388	VAL	CA-CB	6.01	1.61	1.54
1	B	250	THR	CA-CB	5.74	1.62	1.53
1	B	236	ILE	CA-CB	5.72	1.60	1.54
1	A	292	PRO	C-O	-5.71	1.17	1.23
1	B	129	GLU	C-O	-5.51	1.18	1.24
1	B	430	THR	CA-CB	5.46	1.61	1.53

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	226	SER	N-CA-C	7.34	121.96	112.12
1	A	333	ILE	O-C-N	7.05	128.71	121.87
1	B	300	GLU	CB-CG-CD	6.98	124.47	112.60
1	A	226	SER	N-CA-C	6.98	121.47	112.12
1	A	202	LYS	N-CA-C	6.42	119.92	109.59
2	C	5	U	C4'-C3'-O3'	-6.20	103.70	113.00
1	A	208	LYS	N-CA-C	-5.75	100.80	109.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	148	PHE	N-CA-C	5.71	117.58	111.36
2	D	6	U	C2'-C3'-O3'	5.64	117.96	109.50
1	A	146	ALA	N-CA-C	-5.54	105.39	111.82
1	A	225	CYS	N-CA-C	5.46	117.31	111.36
1	B	161	LYS	CA-CB-CG	-5.37	103.35	114.10
1	A	279	ASP	N-CA-C	5.36	117.54	111.11
2	D	4	U	C4'-C3'-C2'	-5.19	97.41	102.60
1	B	225	CYS	N-CA-C	5.18	117.00	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3116	0	3187	51	0
1	B	3074	0	3136	43	0
2	C	120	0	61	3	0
2	D	117	0	62	4	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	6	0	8	2	0
4	B	18	0	24	2	0
5	A	31	0	13	2	0
5	B	31	0	13	1	0
6	A	142	0	0	5	0
6	B	147	0	0	2	0
6	C	7	0	0	0	0
6	D	4	0	0	0	0
All	All	6815	0	6504	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:334:ALA:HB3	1:B:403:SER:OG	1.32	1.22
1:B:227:LYS:HE2	2:D:6:U:O2'	1.57	1.02
1:B:334:ALA:CB	1:B:403:SER:OG	2.19	0.90
1:B:218:PRO:HG3	1:B:247:MET:HE1	1.57	0.87
1:A:110:MET:HE3	1:A:112:PHE:CZ	2.10	0.86
1:A:195:ALA:HB2	1:A:211:GLU:CG	2.07	0.83
1:B:361:ALA:HB2	1:B:384:GLU:HG2	1.61	0.83
1:A:141:GLY:H	5:A:480:ANP:HNB1	1.23	0.83
1:B:313:LEU:HD21	1:B:449:MET:CE	2.09	0.82
1:A:314:CYS:O	1:A:444:ASP:HA	1.81	0.80
1:A:195:ALA:HB2	1:A:211:GLU:HG3	1.63	0.80
1:B:195:ALA:HB2	1:B:211:GLU:HG3	1.64	0.79
1:B:227:LYS:CE	2:D:6:U:O2'	2.31	0.77
1:B:277:PHE:O	4:B:3732:GOL:H31	1.85	0.76
1:A:226:SER:HB3	1:A:263:MET:HE1	1.69	0.73
1:B:152:MET:O	1:B:156:VAL:HG23	1.88	0.73
1:B:141:GLY:H	5:B:2:ANP:HNB1	1.37	0.72
2:C:6:U:O2	2:C:6:U:H2'	1.90	0.71
1:A:413:ASP:OD2	1:A:417:ASN:HB2	1.91	0.71
1:A:334:ALA:HB3	1:A:403:SER:OG	1.90	0.70
1:A:218:PRO:HG3	1:A:247:MET:HE1	1.74	0.70
1:A:310:TYR:CD2	1:A:438:LEU:HD11	2.27	0.69
1:B:256:GLN:NE2	6:B:501:HOH:O	2.26	0.69
1:B:195:ALA:HB2	1:B:211:GLU:CG	2.23	0.68
1:B:414:LYS:HE2	2:D:1:U:H3	1.59	0.68
1:B:78:GLU:OE2	1:B:82:ARG:HD3	1.93	0.68
1:B:180:LYS:HD3	6:B:583:HOH:O	1.95	0.66
1:B:246:VAL:HG21	4:B:3733:GOL:H31	1.78	0.66
1:B:361:ALA:CB	1:B:384:GLU:HG2	2.26	0.65
1:A:209:ILE:HG23	1:A:211:GLU:OE1	1.96	0.64
1:B:313:LEU:HD21	1:B:449:MET:HE2	1.79	0.64
1:A:277:PHE:O	4:A:3731:GOL:H11	1.99	0.62
1:A:328:TYR:O	1:A:332:THR:HG22	2.02	0.60
1:A:277:PHE:C	4:A:3731:GOL:H11	2.27	0.60
1:A:110:MET:HE3	1:A:112:PHE:CE2	2.38	0.59
1:B:334:ALA:HB3	1:B:403:SER:HG	1.60	0.59
1:A:195:ALA:CB	1:A:211:GLU:HG3	2.33	0.58
1:A:209:ILE:HD13	1:A:224:TRP:CZ3	2.39	0.58
1:B:176:LEU:O	1:B:180:LYS:HD2	2.05	0.57
1:A:137:GLN:HA	1:A:274:SER:O	2.05	0.56
1:B:195:ALA:CB	1:B:211:GLU:HG3	2.34	0.56
1:A:310:TYR:CE2	1:A:438:LEU:CD1	2.89	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:334:ALA:HB3	1:B:403:SER:CB	2.33	0.55
1:B:227:LYS:HE2	2:D:6:U:HO2'	1.68	0.55
1:B:181:VAL:O	1:B:185:MET:HG3	2.06	0.55
1:A:209:ILE:CD1	1:A:224:TRP:CZ3	2.90	0.55
1:A:363:LEU:HD23	1:A:393:CYS:SG	2.46	0.55
1:B:334:ALA:CB	1:B:403:SER:CB	2.85	0.54
1:B:137:GLN:HA	1:B:274:SER:O	2.08	0.54
1:B:228:LEU:HD13	1:B:230:PHE:CE1	2.43	0.54
1:B:209:ILE:CD1	1:B:224:TRP:CZ3	2.91	0.53
1:A:233:PRO:HA	1:A:236:ILE:HD12	1.91	0.53
1:A:310:TYR:HD2	1:A:438:LEU:HD11	1.73	0.52
1:A:251:GLN:HG3	6:A:540:HOH:O	2.08	0.52
1:A:456:GLN:HG2	1:A:463:ILE:HG13	1.91	0.52
1:A:181:VAL:O	1:A:185:MET:HG3	2.10	0.52
1:A:310:TYR:CD2	1:A:438:LEU:CD1	2.93	0.51
1:B:107:VAL:HG12	1:B:185:MET:HE1	1.93	0.51
1:A:209:ILE:CD1	1:A:224:TRP:CE3	2.94	0.51
1:B:209:ILE:HD11	1:B:224:TRP:CZ3	2.46	0.50
1:A:195:ALA:HB2	1:A:211:GLU:HG2	1.92	0.49
1:A:195:ALA:N	1:A:211:GLU:HG3	2.27	0.49
1:A:211:GLU:H	1:A:211:GLU:CD	2.20	0.49
1:B:255:ASP:HB3	1:B:259:ARG:NH1	2.28	0.49
1:A:425:HIS:HD2	6:A:49:HOH:O	1.96	0.48
2:C:6:U:O2	2:C:6:U:C2'	2.61	0.48
1:A:413:ASP:OD1	1:A:415:ASP:HB2	2.14	0.47
1:B:164:GLN:HB3	1:B:237:LYS:HE3	1.96	0.47
1:A:110:MET:HE2	1:A:185:MET:SD	2.54	0.47
1:A:331:ILE:HD13	1:A:406:ILE:HD11	1.96	0.47
1:A:199:ARG:HG3	2:C:5:U:OP2	2.15	0.47
1:A:310:TYR:CE2	1:A:438:LEU:HD11	2.50	0.46
1:B:158:PRO:HB2	1:B:191:GLU:OE1	2.16	0.45
1:B:464:GLU:HA	1:B:464:GLU:OE1	2.15	0.45
1:A:282:TRP:O	1:A:286:GLN:HG2	2.17	0.45
1:B:185:MET:HB3	1:B:185:MET:HE2	1.53	0.44
5:A:480:ANP:H8	6:A:30:HOH:O	2.17	0.44
1:A:328:TYR:O	1:A:332:THR:CG2	2.66	0.43
1:B:241:LEU:HD12	1:B:272:LEU:HD22	1.99	0.43
1:A:319:GLU:O	1:A:322:GLN:HB3	2.18	0.43
1:A:321:PHE:CE2	1:A:325:CYS:SG	3.12	0.43
1:A:224:TRP:HB3	1:A:231:ILE:HD12	2.01	0.43
1:A:310:TYR:HE2	1:A:438:LEU:CD1	2.30	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:PRO:CG	1:A:247:MET:HE1	2.47	0.42
1:A:226:SER:HB3	1:A:263:MET:CE	2.45	0.42
1:B:328:TYR:HA	1:B:331:ILE:HG12	2.01	0.42
1:A:276:THR:OG1	1:A:425:HIS:HE1	2.02	0.42
1:A:368:MET:HE3	1:A:371:GLN:OE1	2.20	0.42
1:B:260:ILE:HA	1:B:263:MET:HE2	2.01	0.42
1:B:209:ILE:HD13	1:B:224:TRP:CZ3	2.55	0.42
1:B:282:TRP:CZ2	1:B:296:LYS:HD3	2.55	0.41
1:A:303:THR:HG23	6:A:572:HOH:O	2.21	0.41
1:B:251:GLN:OE1	1:B:414:LYS:HG3	2.21	0.41
1:A:413:ASP:OD1	1:A:415:ASP:N	2.54	0.41
1:B:77:VAL:HG13	1:B:78:GLU:N	2.36	0.41
1:A:234:LYS:HD2	1:A:263:MET:O	2.21	0.40
1:A:199:ARG:NH2	6:A:29:HOH:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	391/412 (95%)	387 (99%)	4 (1%)	0	100	100
1	B	384/412 (93%)	380 (99%)	4 (1%)	0	100	100
All	All	775/824 (94%)	767 (99%)	8 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/363 (95%)	337 (98%)	8 (2%)	44	59
1	B	341/363 (94%)	334 (98%)	7 (2%)	47	63
All	All	686/726 (94%)	671 (98%)	15 (2%)	45	61

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	GLN
1	A	129	GLU
1	A	155	GLN
1	A	180	LYS
1	A	211	GLU
1	A	215	ILE
1	A	414	LYS
1	A	429	ARG
1	B	107	VAL
1	B	110	MET
1	B	193	LYS
1	B	211	GLU
1	B	215	ILE
1	B	429	ARG
1	B	453	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	121	ASN
1	A	201	ASN
1	A	256	GLN
1	A	286	GLN
1	A	322	GLN
1	A	401	GLN
1	A	425	HIS
1	B	113	ASN
1	B	121	ASN
1	B	155	GLN
1	B	256	GLN
1	B	358	HIS
1	B	447	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	5/10 (50%)	1 (20%)	0
2	D	5/10 (50%)	0	0
All	All	10/20 (50%)	1 (10%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	C	6	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ANP	A	480	3	33,33,33	2.24	9 (27%)	45,52,52	2.48	14 (31%)
4	GOL	B	3734	-	5,5,5	0.23	0	5,5,5	1.11	0
4	GOL	B	3732	-	5,5,5	0.29	0	5,5,5	1.29	1 (20%)
5	ANP	B	2	3	33,33,33	2.09	8 (24%)	45,52,52	2.48	13 (28%)
4	GOL	A	3731	-	5,5,5	0.37	0	5,5,5	1.41	1 (20%)
4	GOL	B	3733	-	5,5,5	0.31	0	5,5,5	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ANP	A	480	3	-	2/18/38/38	0/3/3/3
4	GOL	B	3734	-	-	2/4/4/4	-
4	GOL	B	3732	-	-	4/4/4/4	-
5	ANP	B	2	3	-	2/18/38/38	0/3/3/3
4	GOL	A	3731	-	-	2/4/4/4	-
4	GOL	B	3733	-	-	0/4/4/4	-

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	480	ANP	PG-O1G	6.43	1.55	1.46
5	B	2	ANP	C5-C4	5.34	1.48	1.39
5	B	2	ANP	PB-O3A	4.95	1.65	1.59
5	A	480	ANP	PB-O1B	4.90	1.53	1.46
5	A	480	ANP	PG-N3B	4.50	1.75	1.63
5	A	480	ANP	C5-C4	4.28	1.46	1.39
5	B	2	ANP	PG-O1G	4.21	1.52	1.46
5	B	2	ANP	PG-N3B	3.76	1.73	1.63
5	A	480	ANP	PB-O3A	3.70	1.63	1.59
5	B	2	ANP	PB-O2B	-3.56	1.47	1.56
5	A	480	ANP	PB-O2B	-3.14	1.48	1.56
5	A	480	ANP	C8-N7	2.72	1.36	1.31
5	B	2	ANP	PG-O3G	-2.68	1.49	1.56
5	A	480	ANP	PG-O2G	-2.54	1.50	1.56
5	B	2	ANP	PB-N3B	2.33	1.69	1.63
5	A	480	ANP	C5-C6	2.02	1.46	1.41
5	B	2	ANP	C5-N7	-2.00	1.35	1.39

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	480	ANP	O1G-PG-N3B	-8.44	99.34	111.77
5	B	2	ANP	O1G-PG-N3B	-7.74	100.38	111.77
5	B	2	ANP	N3-C2-N1	-6.21	119.19	128.58
5	B	2	ANP	C2-N1-C6	5.77	128.21	118.73
5	A	480	ANP	N3-C2-N1	-5.28	120.59	128.58
5	A	480	ANP	C2-N1-C6	4.52	126.16	118.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	480	ANP	O4'-C1'-C2'	-4.18	97.68	106.62
5	A	480	ANP	C5-C4-N3	-4.12	121.04	126.72
5	A	480	ANP	O4'-C1'-N9	-4.04	100.32	108.09
5	A	480	ANP	O2G-PG-O3G	3.96	118.23	107.59
5	B	2	ANP	O2B-PB-O1B	3.92	118.27	109.87
5	B	2	ANP	N3-C4-N9	3.72	133.49	127.17
5	B	2	ANP	O3'-C3'-C4'	-3.43	101.24	111.08
5	A	480	ANP	C2-N3-C4	3.41	120.17	111.83
5	B	2	ANP	C5-C4-N3	-3.36	122.09	126.72
5	B	2	ANP	C2-N3-C4	3.34	120.00	111.83
5	B	2	ANP	O2G-PG-O3G	3.23	116.28	107.59
5	A	480	ANP	N3-C4-N9	3.23	132.66	127.17
5	B	2	ANP	O4'-C1'-C2'	-3.14	99.89	106.62
5	B	2	ANP	C5'-C4'-C3'	-2.88	104.86	115.21
5	A	480	ANP	C6-C5-N7	2.71	137.31	132.09
4	A	3731	GOL	O3-C3-C2	-2.61	98.63	110.38
5	A	480	ANP	C4-C5-N7	-2.58	107.63	110.58
5	B	2	ANP	C4-N9-C8	2.54	108.40	105.74
5	A	480	ANP	C4-N9-C8	2.32	108.18	105.74
5	B	2	ANP	O3A-PB-N3B	-2.28	100.26	106.59
5	A	480	ANP	O2B-PB-O1B	2.12	114.42	109.87
5	A	480	ANP	C5-N7-C8	2.10	106.74	103.45
4	B	3732	GOL	O2-C2-C3	2.01	117.50	109.18

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	3731	GOL	C1-C2-C3-O3
4	B	3732	GOL	O1-C1-C2-O2
4	B	3732	GOL	O1-C1-C2-C3
4	B	3732	GOL	C1-C2-C3-O3
4	B	3734	GOL	C1-C2-C3-O3
5	A	480	ANP	PA-O3A-PB-O1B
5	A	480	ANP	PA-O3A-PB-O2B
5	B	2	ANP	PA-O3A-PB-O2B
4	A	3731	GOL	O2-C2-C3-O3
4	B	3732	GOL	O2-C2-C3-O3
4	B	3734	GOL	O2-C2-C3-O3
5	B	2	ANP	PA-O3A-PB-O1B


There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	480	ANP	2	0
4	B	3732	GOL	1	0
5	B	2	ANP	1	0
4	A	3731	GOL	2	0
4	B	3733	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	390/412 (94%)	-1.47	0 100 100	6, 18, 33, 40	20 (5%)
1	B	387/412 (93%)	-1.49	0 100 100	6, 18, 33, 42	11 (2%)
2	C	6/10 (60%)	-1.68	0 100 100	22, 25, 52, 55	0
2	D	6/10 (60%)	-1.69	0 100 100	23, 25, 41, 67	0
All	All	789/844 (93%)	-1.48	0 100 100	6, 18, 34, 67	31 (3%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	B	3732	6/6	0.98	0.05	43,46,48,51	0
4	GOL	B	3733	6/6	0.98	0.05	36,41,43,44	0
4	GOL	B	3734	6/6	0.98	0.05	51,54,55,56	0
4	GOL	A	3731	6/6	0.99	0.03	33,36,39,40	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	A	1	1/1	1.00	0.01	15,15,15,15	0
3	MG	B	1	1/1	1.00	0.03	16,16,16,16	0
5	ANP	A	480	31/31	1.00	0.02	7,14,17,20	0
5	ANP	B	2	31/31	1.00	0.01	7,14,19,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.