



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:42 PM UTC

PDB ID : 3FIU / pdb_00003fiu
Title : Structure of NMN synthetase from Francisella tularensis
Authors : Sorci, L.; Martynowski, D.; Eyobo, Y.; Osterman, A.L.; Zhang, H.
Deposited on : 2008-12-12
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	249	4% 80% 14% • •
1	B	249	% 87% 8% •
1	C	249	3% 86% 9% • •
1	D	249	2% 83% 11% • •

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8516 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NH(3)-dependent NAD(+) synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	238	Total	C	N	O	S	0	3	0
			1896	1208	314	365	9			
1	B	238	Total	C	N	O	S	0	0	0
			1885	1203	312	361	9			
1	C	238	Total	C	N	O	S	0	4	0
			1898	1211	314	363	10			
1	D	238	Total	C	N	O	S	0	0	0
			1885	1203	312	361	9			

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		
2	B	2	Total	Mg	0	0
			2	2		
2	C	2	Total	Mg	0	0
			2	2		
2	D	2	Total	Mg	0	0
			2	2		

- Molecule 3 is ADENOSINE MONOPHOSPHATE (CCD ID: AMP) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	D	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 4 is PYROPHOSPHATE 2- (CCD ID: POP) (formula: $\text{H}_2\text{O}_7\text{P}_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 9 7 2	0	0
4	B	1	Total O P 9 7 2	0	0
4	C	1	Total O P 9 7 2	0	0
4	D	1	Total O P 9 7 2	0	0

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Na 1 1	0	0
5	B	1	Total Na 1 1	0	0
5	C	1	Total Na 1 1	0	0
5	D	1	Total Na 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	205	Total O 205 205	0	0
6	B	214	Total O 214 214	0	0
6	C	196	Total O 196 196	0	0
6	D	197	Total O 197 197	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NH(3)-dependent NAD(+) synthetase




- Molecule 1: NH(3)-dependent NAD(+) synthetase




- Molecule 1: NH(3)-dependent NAD(+) synthetase



- Molecule 1: NH(3)-dependent NAD(+) synthetase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.43Å 126.44Å 152.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.85 40.00 – 1.85	Depositor EDS
% Data completeness (in resolution range)	93.9 (40.00-1.85) 93.8 (40.00-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 1.84Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.184 , 0.233 (Not available) , 0.234	Depositor DCC
R_{free} test set	3939 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.195	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8516	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 66.51 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.9771e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, NA, POP, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	0/1953	0.84	0/2645
1	B	0.76	0/1927	0.83	0/2610
1	C	0.78	0/1959	0.81	1/2651 (0.0%)
1	D	0.78	0/1927	0.85	1/2610 (0.0%)
All	All	0.78	0/7766	0.83	2/10516 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	104	PHE	N-CA-C	5.37	119.80	112.88
1	C	7	PHE	N-CA-C	5.09	116.70	108.26

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1896	0	1844	34	0
1	B	1885	0	1836	16	0
1	C	1898	0	1855	21	0
1	D	1885	0	1836	26	0
2	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	23	0	12	0	0
3	B	23	0	12	0	0
3	C	23	0	12	0	0
3	D	23	0	12	0	0
4	A	9	0	0	0	0
4	B	9	0	0	0	0
4	C	9	0	0	0	0
4	D	9	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	A	205	0	0	9	0
6	B	214	0	0	3	0
6	C	196	0	0	3	0
6	D	197	0	0	6	0
All	All	8516	0	7419	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231:ASN:HB2	6:A:824:HOH:O	1.58	1.03
1:C:236:ARG:HB3	6:C:732:HOH:O	1.71	0.90
1:A:63:SER:H	1:A:66:ASN:HD22	1.17	0.89
1:D:68:HIS:CD2	1:D:72:GLN:NE2	2.47	0.82
1:D:63:SER:H	1:D:66:ASN:HD22	1.28	0.81
1:C:115:ILE:HD11	1:D:133:GLN:HB3	1.63	0.80
1:A:134:GLN:HE22	1:B:105:THR:HA	1.51	0.74
1:A:221:GLN:HB2	1:C:71:MET:HE1	1.71	0.73
1:B:112:GLN:HE21	1:B:114:VAL:H	1.39	0.69
1:A:63:SER:H	1:A:66:ASN:ND2	1.89	0.69
1:D:63:SER:H	1:D:66:ASN:ND2	1.91	0.69
1:A:238:HIS:HE1	6:A:326:HOH:O	1.77	0.68
1:A:156:PHE:HA	1:A:160:ALA:HB2	1.76	0.67
1:A:238:HIS:CE1	6:A:326:HOH:O	2.48	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:80:MET:HE1	1:D:182:ASP:O	1.95	0.65
1:D:68:HIS:HD2	1:D:72:GLN:NE2	1.92	0.65
1:C:25[A]:MET:HE2	6:C:394:HOH:O	1.98	0.63
1:D:238:HIS:HB2	6:D:645:HOH:O	1.99	0.62
1:C:134:GLN:HG2	1:D:115:ILE:HD13	1.81	0.62
1:B:225:LYS:HB2	6:B:331:HOH:O	2.00	0.61
1:A:221:GLN:CB	1:C:71:MET:HE1	2.32	0.59
1:D:71:MET:HB2	6:D:282:HOH:O	2.02	0.59
1:A:60:ILE:HG21	1:A:71:MET:HG3	1.85	0.59
1:C:134:GLN:HE22	1:D:105:THR:HA	1.68	0.59
1:A:51:LYS:HE3	6:A:694:HOH:O	2.03	0.58
1:D:156:PHE:HA	1:D:160:ALA:HB2	1.84	0.58
1:D:25:MET:HE3	6:D:392:HOH:O	2.04	0.58
1:A:77:LEU:HG	1:A:81:LEU:HD22	1.86	0.57
1:C:115:ILE:CD1	1:D:133:GLN:HB3	2.32	0.57
1:C:156:PHE:HA	1:C:160:ALA:HB2	1.86	0.57
1:D:68:HIS:CD2	1:D:72:GLN:HE21	2.22	0.57
1:D:78:ILE:HG21	1:D:85:HIS:CG	2.41	0.56
1:A:80:MET:HG3	6:A:582:HOH:O	2.04	0.55
1:A:112:GLN:N	6:A:476:HOH:O	2.40	0.55
1:B:112:GLN:HG3	1:B:115:ILE:H	1.70	0.55
1:A:60:ILE:HG22	1:A:62:PRO:HD3	1.88	0.55
1:C:13[A]:SER:OG	1:C:177:LEU:HD22	2.07	0.54
1:A:65:ASN:HD21	1:A:116:LYS:HE2	1.71	0.54
1:A:133:GLN:HE22	1:B:114:VAL:HG12	1.72	0.54
1:B:60:ILE:HG22	1:B:62:PRO:HD3	1.90	0.54
1:A:134:GLN:NE2	1:B:105:THR:HA	2.23	0.54
1:D:221:GLN:HG3	6:D:666:HOH:O	2.10	0.52
1:B:156:PHE:HA	1:B:160:ALA:HB2	1.92	0.52
1:A:12:TYR:CE2	1:A:16:LEU:HD22	2.45	0.52
1:A:228:GLU:O	6:A:824:HOH:O	2.19	0.51
1:D:60:ILE:HG22	1:D:62:PRO:HD3	1.93	0.51
1:A:50:VAL:HG21	1:A:81:LEU:HG	1.92	0.50
1:A:12:TYR:CE2	1:A:169:LEU:HD21	2.47	0.50
1:A:71:MET:HE2	1:A:72[B]:GLN:HE21	1.75	0.50
1:A:113:LEU:HD12	1:A:196:LEU:HD13	1.93	0.50
1:C:130:ALA:O	1:C:134:GLN:HG3	2.12	0.50
1:C:225[B]:LYS:NZ	6:C:493:HOH:O	2.34	0.50
1:A:78:ILE:HG21	1:A:85:HIS:CG	2.47	0.50
1:A:71:MET:CE	1:A:72[B]:GLN:NE2	2.75	0.49
1:C:133:GLN:HE22	1:D:114:VAL:HG22	1.77	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:227:LEU:HB3	6:A:831:HOH:O	2.12	0.49
1:B:12:TYR:CE2	1:B:169:LEU:HD21	2.48	0.48
1:A:65:ASN:ND2	1:A:116:LYS:HE2	2.29	0.48
1:A:71:MET:CE	1:A:72[B]:GLN:HE21	2.27	0.47
1:D:238:HIS:HB3	6:D:807:HOH:O	2.15	0.46
1:B:80:MET:CE	6:B:606:HOH:O	2.63	0.46
6:A:317:HOH:O	1:B:92:PRO:HB2	2.17	0.45
1:B:134:GLN:NE2	6:B:652:HOH:O	2.50	0.45
1:C:78:ILE:HG21	1:C:85:HIS:CG	2.52	0.45
1:C:134:GLN:HG2	1:D:115:ILE:CD1	2.47	0.45
1:D:63:SER:N	1:D:66:ASN:HD22	2.06	0.45
1:D:12:TYR:CE2	1:D:16:LEU:HD22	2.52	0.44
1:A:221:GLN:HB2	1:C:71:MET:CE	2.46	0.44
1:B:146:CYS:SG	1:B:214:ASP:OD1	2.77	0.43
1:C:71:MET:HE2	1:C:75:LEU:HD11	2.00	0.43
1:A:101:THR:O	1:A:105:THR:HB	2.19	0.42
1:C:126:MET:HE1	1:D:123:LEU:HD13	2.01	0.42
1:B:134:GLN:HE21	1:B:134:GLN:HB3	1.60	0.42
1:B:112:GLN:HB3	1:B:115:ILE:HD12	2.02	0.42
1:A:133:GLN:HE22	1:B:114:VAL:CG1	2.31	0.42
1:A:146:CYS:SG	1:A:214:ASP:OD1	2.78	0.42
1:C:12:TYR:CE2	1:C:169:LEU:HD21	2.55	0.41
1:A:237:SER:HB2	1:A:240:LYS:HD3	2.02	0.41
1:D:16:LEU:HD12	1:D:16:LEU:HA	1.90	0.41
1:D:6:ASP:HB3	6:D:676:HOH:O	2.20	0.40
1:C:6:ASP:HB3	1:C:7:PHE:H	1.54	0.40
1:D:58:ALA:O	1:D:85:HIS:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/249 (95%)	237 (100%)	0	0	100	100
1	B	234/249 (94%)	232 (99%)	1 (0%)	1 (0%)	30	17
1	C	238/249 (96%)	237 (100%)	1 (0%)	0	100	100
1	D	234/249 (94%)	234 (100%)	0	0	100	100
All	All	943/996 (95%)	940 (100%)	2 (0%)	1 (0%)	48	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	7	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/212 (96%)	197 (97%)	7 (3%)	32	17
1	B	201/212 (95%)	196 (98%)	5 (2%)	42	27
1	C	205/212 (97%)	197 (96%)	8 (4%)	28	13
1	D	201/212 (95%)	193 (96%)	8 (4%)	28	13
All	All	811/848 (96%)	783 (96%)	28 (4%)	32	16

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ASP
1	A	8	SER
1	A	16	LEU
1	A	81	LEU
1	A	105	THR
1	A	113	LEU
1	A	205	GLU
1	B	123	LEU
1	B	134	GLN
1	B	196	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	221	GLN
1	B	248	ASN
1	C	6	ASP
1	C	8	SER
1	C	16	LEU
1	C	105	THR
1	C	205	GLU
1	C	225[A]	LYS
1	C	225[B]	LYS
1	C	248	ASN
1	D	16	LEU
1	D	30	GLU
1	D	50	VAL
1	D	75	LEU
1	D	95	GLU
1	D	105	THR
1	D	123	LEU
1	D	225	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (40) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	ASN
1	A	66	ASN
1	A	69	GLN
1	A	102	GLN
1	A	120	GLN
1	A	133	GLN
1	A	134	GLN
1	A	238	HIS
1	A	239	HIS
1	B	26	ASN
1	B	68	HIS
1	B	72	GLN
1	B	85	HIS
1	B	112	GLN
1	B	118	ASN
1	B	136	ASN
1	B	168	ASN
1	B	198	GLN
1	B	221	GLN
1	B	231	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	239	HIS
1	C	26	ASN
1	C	82	ASN
1	C	102	GLN
1	C	120	GLN
1	C	134	GLN
1	C	136	ASN
1	C	186	ASN
1	C	231	ASN
1	C	239	HIS
1	C	248	ASN
1	D	26	ASN
1	D	66	ASN
1	D	68	HIS
1	D	72	GLN
1	D	120	GLN
1	D	136	ASN
1	D	168	ASN
1	D	231	ASN
1	D	239	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 12 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	AMP	A	3001	5,2	25,25,25	1.46	3 (12%)	37,38,38	1.74	10 (27%)
4	POP	D	4001	5,2	6,8,8	0.76	0	12,13,13	0.88	0
4	POP	A	4001	5,2	6,8,8	0.84	0	12,13,13	0.97	1 (8%)
3	AMP	C	3001	5,2	25,25,25	1.53	3 (12%)	37,38,38	2.00	11 (29%)
3	AMP	D	3001	5,2	25,25,25	1.40	5 (20%)	37,38,38	1.79	10 (27%)
4	POP	C	4001	5,2	6,8,8	0.82	0	12,13,13	0.79	0
3	AMP	B	3001	5,2	25,25,25	1.53	4 (16%)	37,38,38	1.72	8 (21%)
4	POP	B	4001	5,2	6,8,8	0.71	0	12,13,13	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AMP	A	3001	5,2	-	0/10/26/26	0/3/3/3
4	POP	D	4001	5,2	-	0/6/6/6	-
4	POP	A	4001	5,2	-	0/6/6/6	-
3	AMP	C	3001	5,2	-	0/10/26/26	0/3/3/3
3	AMP	D	3001	5,2	-	2/10/26/26	0/3/3/3
4	POP	C	4001	5,2	-	0/6/6/6	-
3	AMP	B	3001	5,2	-	0/10/26/26	0/3/3/3
4	POP	B	4001	5,2	-	0/6/6/6	-

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	3001	AMP	C5-C4	5.16	1.48	1.39
3	A	3001	AMP	C5-C4	4.81	1.47	1.39
3	C	3001	AMP	C5-C4	4.70	1.47	1.39
3	D	3001	AMP	C5-C4	4.25	1.46	1.39
3	C	3001	AMP	C8-N7	3.20	1.37	1.31
3	B	3001	AMP	C8-N7	2.92	1.37	1.31
3	C	3001	AMP	C5-C6	2.73	1.48	1.41
3	A	3001	AMP	C5-C6	2.48	1.47	1.41
3	A	3001	AMP	C8-N7	2.44	1.36	1.31
3	D	3001	AMP	C4-N9	-2.44	1.32	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	3001	AMP	C5-N7	-2.39	1.34	1.39
3	D	3001	AMP	C8-N7	2.38	1.36	1.31
3	B	3001	AMP	C5-C6	2.20	1.47	1.41
3	B	3001	AMP	C4-N9	-2.12	1.33	1.37
3	D	3001	AMP	C5-C6	2.02	1.46	1.41

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	3001	AMP	C5-C4-N3	-4.60	120.38	126.72
3	C	3001	AMP	N3-C2-N1	-4.53	121.72	128.58
3	D	3001	AMP	C5-C4-N3	-4.28	120.82	126.72
3	D	3001	AMP	N3-C4-N9	4.03	134.02	127.17
3	B	3001	AMP	C5-C4-N3	-3.93	121.30	126.72
3	C	3001	AMP	C4-N9-C8	3.81	109.74	105.74
3	A	3001	AMP	C5-C4-N3	-3.81	121.47	126.72
3	D	3001	AMP	N3-C2-N1	-3.81	122.81	128.58
3	C	3001	AMP	N3-C4-N9	3.79	133.62	127.17
3	B	3001	AMP	N3-C4-N9	3.77	133.58	127.17
3	B	3001	AMP	C4-N9-C8	3.73	109.66	105.74
3	A	3001	AMP	C2-N1-C6	3.61	124.67	118.73
3	A	3001	AMP	N3-C2-N1	-3.61	123.12	128.58
3	C	3001	AMP	C2-N3-C4	3.60	120.62	111.83
3	B	3001	AMP	N3-C2-N1	-3.53	123.24	128.58
3	A	3001	AMP	N3-C4-N9	3.41	132.97	127.17
3	A	3001	AMP	C4-N9-C8	3.10	108.99	105.74
3	D	3001	AMP	C2-N1-C6	3.00	123.66	118.73
3	B	3001	AMP	C2-N1-C6	2.98	123.62	118.73
3	C	3001	AMP	C2-N1-C6	2.97	123.62	118.73
3	C	3001	AMP	C4-C5-N7	-2.93	107.23	110.58
3	D	3001	AMP	C4-N9-C8	2.93	108.82	105.74
3	D	3001	AMP	C2-N3-C4	2.81	118.71	111.83
3	C	3001	AMP	N9-C8-N7	-2.75	110.03	113.94
3	A	3001	AMP	O2P-P-O5'	-2.70	99.64	106.67
3	B	3001	AMP	C2-N3-C4	2.69	118.40	111.83
3	C	3001	AMP	C6-C5-N7	2.60	137.11	132.09
3	C	3001	AMP	O2P-P-O5'	-2.55	100.03	106.67
4	A	4001	POP	O6-P2-O	2.53	113.13	104.64
3	D	3001	AMP	O2P-P-O5'	-2.52	100.10	106.67
3	A	3001	AMP	C4-C5-N7	-2.44	107.79	110.58
3	A	3001	AMP	C2-N3-C4	2.42	117.74	111.83
3	B	3001	AMP	O2P-P-O5'	-2.30	100.67	106.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	3001	AMP	C2'-C1'-N9	-2.29	107.62	113.30
3	A	3001	AMP	N6-C6-N1	2.29	123.47	118.38
3	D	3001	AMP	N6-C6-N1	2.27	123.42	118.38
3	C	3001	AMP	C5-N7-C8	2.26	107.00	103.45
3	D	3001	AMP	C3'-C2'-C1'	2.26	105.73	101.46
3	B	3001	AMP	N9-C8-N7	-2.04	111.05	113.94
3	A	3001	AMP	O3P-P-O1P	2.01	118.65	110.83

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	3001	AMP	C3'-C4'-C5'-O5'
3	D	3001	AMP	O4'-C4'-C5'-O5'

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand AMP A 3001



Ligand AMP C 3001





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	238/249 (95%)	0.11	11 (4%) 37 40	15, 25, 40, 60	3 (1%)
1	B	238/249 (95%)	-0.12	3 (1%) 75 79	14, 23, 36, 56	0
1	C	238/249 (95%)	0.04	8 (3%) 48 52	15, 23, 37, 59	4 (1%)
1	D	238/249 (95%)	-0.04	6 (2%) 58 62	15, 24, 38, 58	0
All	All	952/996 (95%)	-0.00	28 (2%) 53 57	14, 23, 38, 60	7 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	105	THR	4.0
1	D	105	THR	3.5
1	A	226	ALA	3.2
1	C	6	ASP	3.2
1	C	105	THR	3.1
1	A	148	TRP	2.8
1	A	221	GLN	2.8
1	C	224	ALA	2.8
1	A	248	ASN	2.8
1	C	114	VAL	2.8
1	D	6	ASP	2.7
1	B	105	THR	2.7
1	D	238	HIS	2.7
1	A	6	ASP	2.6
1	C	248	ASN	2.4
1	C	134	GLN	2.4
1	C	112	GLN	2.4
1	A	233	TRP	2.4
1	D	112	GLN	2.3
1	D	68	HIS	2.3
1	A	225	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	204	ASP	2.2
1	A	224	ALA	2.2
1	B	6	ASP	2.1
1	A	238	HIS	2.1
1	C	227	LEU	2.0
1	A	236	ARG	2.0
1	B	235	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NA	A	250	1/1	0.96	0.05	19,19,19,19	0
3	AMP	A	3001	23/23	0.97	0.05	13,21,23,26	0
3	AMP	D	3001	23/23	0.97	0.05	16,19,22,23	0
2	MG	B	5001	1/1	0.97	0.06	19,19,19,19	0
5	NA	D	250	1/1	0.97	0.06	19,19,19,19	0
3	AMP	B	3001	23/23	0.98	0.04	14,18,21,22	0
3	AMP	C	3001	23/23	0.98	0.04	13,18,21,23	0
2	MG	A	5001	1/1	0.98	0.07	20,20,20,20	0
4	POP	A	4001	9/9	0.98	0.04	17,18,19,19	0
2	MG	D	5002	1/1	0.98	0.03	20,20,20,20	0
5	NA	B	250	1/1	0.98	0.05	18,18,18,18	0
5	NA	C	250	1/1	0.98	0.06	18,18,18,18	0
2	MG	A	5002	1/1	0.98	0.03	20,20,20,20	0
4	POP	C	4001	9/9	0.99	0.04	16,17,18,19	0
4	POP	D	4001	9/9	0.99	0.04	15,17,19,20	0
2	MG	B	5002	1/1	0.99	0.03	20,20,20,20	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MG	C	5002	1/1	0.99	0.02	20,20,20,20	0
2	MG	D	5001	1/1	0.99	0.04	16,16,16,16	0
4	POP	B	4001	9/9	0.99	0.03	14,16,19,20	0
2	MG	C	5001	1/1	1.00	0.06	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around AMP A 3001:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AMP D 3001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around AMP B 3001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around AMP C 3001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.