



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:11 AM UTC

PDB ID : 4FIX / pdb_00004fix
Title : Crystal Structure of GlfT2
Authors : Wheatley, R.W.; Zheng, R.B.; Lowary, T.L.; Ng, K.K.S.
Deposited on : 2012-06-11
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	657	 2% 89% 7%
1	B	657	 3% 90% 5%

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 10408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-galactofuranosyl transferase GlfT2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	629	4983	3164	899	898	22	0	0	0
1	B	629	4983	3164	899	898	22	0	0	0

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP O53585
A	-18	GLY	-	expression tag	UNP O53585
A	-17	SER	-	expression tag	UNP O53585
A	-16	SER	-	expression tag	UNP O53585
A	-15	HIS	-	expression tag	UNP O53585
A	-14	HIS	-	expression tag	UNP O53585
A	-13	HIS	-	expression tag	UNP O53585
A	-12	HIS	-	expression tag	UNP O53585
A	-11	HIS	-	expression tag	UNP O53585
A	-10	HIS	-	expression tag	UNP O53585
A	-9	SER	-	expression tag	UNP O53585
A	-8	SER	-	expression tag	UNP O53585
A	-7	GLY	-	expression tag	UNP O53585
A	-6	LEU	-	expression tag	UNP O53585
A	-5	VAL	-	expression tag	UNP O53585
A	-4	PRO	-	expression tag	UNP O53585
A	-3	ARG	-	expression tag	UNP O53585
A	-2	GLY	-	expression tag	UNP O53585
A	-1	SER	-	expression tag	UNP O53585
A	0	HIS	-	expression tag	UNP O53585
B	-19	MET	-	expression tag	UNP O53585
B	-18	GLY	-	expression tag	UNP O53585
B	-17	SER	-	expression tag	UNP O53585
B	-16	SER	-	expression tag	UNP O53585
B	-15	HIS	-	expression tag	UNP O53585

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP O53585
B	-13	HIS	-	expression tag	UNP O53585
B	-12	HIS	-	expression tag	UNP O53585
B	-11	HIS	-	expression tag	UNP O53585
B	-10	HIS	-	expression tag	UNP O53585
B	-9	SER	-	expression tag	UNP O53585
B	-8	SER	-	expression tag	UNP O53585
B	-7	GLY	-	expression tag	UNP O53585
B	-6	LEU	-	expression tag	UNP O53585
B	-5	VAL	-	expression tag	UNP O53585
B	-4	PRO	-	expression tag	UNP O53585
B	-3	ARG	-	expression tag	UNP O53585
B	-2	GLY	-	expression tag	UNP O53585
B	-1	SER	-	expression tag	UNP O53585
B	0	HIS	-	expression tag	UNP O53585

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		
3	B	1	Total	Mn	0	0
			1	1		

- Molecule 4 is BETA-MERCAPTOETHANOL (CCD ID: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	213	Total	O	0	0
			213	213		
5	B	199	Total	O	0	0
			199	199		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UDP-galactofuranosyl transferase GlfT2



- Molecule 1: UDP-galactofuranosyl transferase GlfT2



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	150.79Å 150.79Å 148.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.45 40.00 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.00-2.45) 99.9 (40.00-2.45)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.45Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.185 , 0.222 0.187 , 0.225	Depositor DCC
R_{free} test set	3195 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	47.4	Xtrriage
Anisotropy	0.314	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 35.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.000 for -h,-l,-k 0.000 for l,-k,h	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10408	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.68 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.0304e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MN, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/5118	0.78	1/6976 (0.0%)
1	B	0.50	0/5118	0.79	0/6976
All	All	0.51	0/10236	0.79	1/13952 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	366	LEU	N-CA-C	-5.08	103.00	110.52

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4983	0	4919	33	0
1	B	4983	0	4919	18	0
2	A	18	0	24	2	0
2	B	6	0	8	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	B	4	0	6	0	0
5	A	213	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	199	0	0	0	0
All	All	10408	0	9876	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:425:TRP:HZ2	1:A:431:GLN:HB3	1.40	0.86
1:A:32:THR:HA	2:A:702:GOL:H31	1.58	0.82
1:A:512:MET:HE2	1:A:556:PHE:HD1	1.51	0.75
1:A:425:TRP:CZ2	1:A:431:GLN:HB3	2.23	0.73
1:B:480:LEU:HB3	1:B:481:PRO:HD3	1.81	0.62
1:A:398:ALA:C	1:A:400:SER:H	2.07	0.62
1:A:29:GLU:HG2	5:A:947:HOH:O	1.99	0.62
1:A:512:MET:HE2	1:A:556:PHE:CD1	2.33	0.61
1:A:460:LYS:HE2	1:A:482:GLN:HG3	1.82	0.60
1:A:80:ARG:HG2	1:A:119:GLU:HG2	1.85	0.58
1:A:1:MET:HG3	1:A:2:SER:H	1.68	0.58
1:A:480:LEU:HB3	1:A:481:PRO:HD3	1.86	0.58
1:B:399:TRP:O	1:B:401:ASP:N	2.37	0.57
1:A:233:SER:OG	1:A:366:LEU:O	2.20	0.57
1:A:288:ASN:HD21	1:A:400:SER:HB2	1.70	0.56
1:B:580:ASP:HB3	1:B:583:LYS:HB2	1.89	0.55
1:A:286:MET:HE3	1:A:296:HIS:CD2	2.44	0.53
1:B:233:SER:HB2	1:B:366:LEU:O	2.10	0.53
1:A:233:SER:CB	1:A:366:LEU:O	2.57	0.52
1:A:428:PRO:HD2	1:A:431:GLN:HG3	1.93	0.51
1:B:512:MET:HE2	1:B:556:PHE:HD1	1.75	0.50
1:B:80:ARG:HG2	1:B:119:GLU:HG2	1.95	0.49
1:B:460:LYS:NZ	1:B:482:GLN:HG2	2.28	0.48
1:A:365:PRO:HG3	1:A:473:PHE:HD2	1.77	0.48
1:A:230:LEU:HB2	1:A:234:GLY:HA3	1.96	0.48
1:A:80:ARG:NH1	5:A:856:HOH:O	2.44	0.47
1:B:550:THR:HG21	1:B:569:ALA:HA	1.95	0.47
1:B:307:PHE:HB2	1:B:425:TRP:CZ3	2.48	0.47
1:B:581:ARG:HH11	1:B:581:ARG:HG2	1.80	0.46
1:A:260:ARG:HB2	1:A:395:TRP:HB2	1.97	0.46
1:B:286:MET:HE3	1:B:296:HIS:CD2	2.50	0.46
1:A:307:PHE:HB2	1:A:425:TRP:CZ3	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:448:ALA:HB2	2:A:703:GOL:H11	1.98	0.45
1:B:233:SER:CB	1:B:366:LEU:O	2.64	0.45
1:A:233:SER:HB2	1:A:366:LEU:O	2.15	0.44
1:A:398:ALA:C	1:A:400:SER:N	2.74	0.44
1:A:366:LEU:HB3	1:A:416:ASN:ND2	2.33	0.44
1:A:537:ASN:HA	1:A:538:PRO:HD2	1.86	0.43
1:A:1:MET:CG	1:A:2:SER:H	2.32	0.43
1:B:400:SER:O	1:B:402:LYS:HG2	2.19	0.42
1:B:165:GLY:HA3	1:B:239:VAL:HG11	2.01	0.42
1:B:435:LEU:C	1:B:435:LEU:HD23	2.45	0.41
1:A:245:LYS:HE3	1:A:246:ASN:HD21	1.86	0.41
1:B:512:MET:HE3	1:B:524:ARG:NE	2.35	0.41
1:A:8:LEU:HD21	1:A:11:ARG:HG3	2.02	0.41
1:A:550:THR:HG21	1:A:569:ALA:HA	2.02	0.41
1:A:513:LYS:HA	1:A:514:PRO:HD3	1.94	0.41
1:B:514:PRO:HA	1:B:515:PRO:HD3	1.83	0.41
1:A:54:VAL:HG23	1:A:56:PHE:CE2	2.56	0.41
1:B:156:ALA:HA	1:B:157:PRO:HD3	1.95	0.41
1:A:165:GLY:HA3	1:A:239:VAL:HG11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	627/657 (95%)	606 (97%)	20 (3%)	1 (0%)	43 54
1	B	627/657 (95%)	604 (96%)	21 (3%)	2 (0%)	36 45
All	All	1254/1314 (95%)	1210 (96%)	41 (3%)	3 (0%)	43 54

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	400	SER
1	A	400	SER
1	B	369	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	525/548 (96%)	522 (99%)	3 (1%)	78 86
1	B	525/548 (96%)	520 (99%)	5 (1%)	68 77
All	All	1050/1096 (96%)	1042 (99%)	8 (1%)	73 83

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	368	ILE
1	A	431	GLN
1	A	603	GLU
1	B	7	SER
1	B	368	ILE
1	B	401	ASP
1	B	594	ARG
1	B	599	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	458	GLN
1	A	484	HIS
1	A	539	GLN
1	B	296	HIS
1	B	362	GLN
1	B	482	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	702	-	5,5,5	0.37	0	5,5,5	0.22	0
2	GOL	A	703	-	5,5,5	0.41	0	5,5,5	0.50	0
2	GOL	A	701	-	5,5,5	0.34	0	5,5,5	0.40	0
2	GOL	B	702	-	5,5,5	0.38	0	5,5,5	0.31	0
4	BME	B	701	-	3,3,3	0.42	0	2,2,2	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	702	-	-	2/4/4/4	-
2	GOL	A	703	-	-	2/4/4/4	-
2	GOL	A	701	-	-	0/4/4/4	-
2	GOL	B	702	-	-	2/4/4/4	-
4	BME	B	701	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	703	GOL	C1-C2-C3-O3
2	B	702	GOL	C1-C2-C3-O3
2	B	702	GOL	O2-C2-C3-O3
2	A	702	GOL	C1-C2-C3-O3
2	A	702	GOL	O2-C2-C3-O3
2	A	703	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	702	GOL	1	0
2	A	703	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	629/657 (95%)	-0.18	16 (2%) 58 61	31, 53, 90, 122	0
1	B	629/657 (95%)	-0.14	18 (2%) 53 55	30, 54, 95, 125	0
All	All	1258/1314 (95%)	-0.16	34 (2%) 56 57	30, 53, 92, 125	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	401	ASP	7.2
1	A	400	SER	5.3
1	B	401	ASP	4.9
1	B	629	ALA	4.7
1	A	402	LYS	4.5
1	A	399	TRP	4.0
1	A	405	ALA	3.8
1	B	400	SER	3.7
1	A	629	ALA	3.6
1	A	1	MET	3.5
1	B	399	TRP	3.5
1	B	428	PRO	3.5
1	B	402	LYS	3.3
1	B	1	MET	3.1
1	A	430	ALA	3.1
1	B	425	TRP	3.0
1	B	405	ALA	3.0
1	B	4	LEU	3.0
1	B	403	ASP	2.9
1	A	355	GLN	2.7
1	A	425	TRP	2.6
1	B	530	MET	2.6
1	B	519	LEU	2.6
1	A	518	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	404	ASP	2.5
1	B	2	SER	2.5
1	A	404	ASP	2.4
1	A	428	PRO	2.4
1	B	516	VAL	2.4
1	A	403	ASP	2.3
1	A	429	LYS	2.2
1	B	628	PRO	2.2
1	A	519	LEU	2.0
1	B	398	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	B	702	6/6	0.75	0.15	73,79,81,82	0
2	GOL	A	703	6/6	0.85	0.19	78,82,84,84	0
4	BME	B	701	4/4	0.90	0.13	60,65,66,69	0
2	GOL	A	702	6/6	0.91	0.13	63,64,65,66	0
2	GOL	A	701	6/6	0.96	0.07	46,56,60,63	0
3	MN	B	703	1/1	0.98	0.04	72,72,72,72	0
3	MN	A	704	1/1	0.98	0.04	73,73,73,73	0

6.5 Other polymers [i](#)

There are no such residues in this entry.