



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:36 PM UTC

PDB ID : 6FIF / pdb_00006fif
Title : Crystal structure of the BRI1 Gly644-Asp (bri1-6) mutant from Arabidopsis thaliana.
Authors : Hothorn, M.; Santiago, J.; Hohmann, U.
Deposited on : 2018-01-18
Resolution : 2.54 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	837	 2% 84% 5% 11%
2	B	2	 50% 50%
2	C	2	 100%
3	D	3	 67% 33%
4	E	3	 67% 33%

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 5846 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein BRASSINOSTEROID INSENSITIVE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	744	5626	3559	928	1107	32	0	8	0

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	644	ASP	GLY	conflict	UNP O22476
A	789	LEU	-	expression tag	UNP O22476
A	790	GLU	-	expression tag	UNP O22476
A	791	ASN	-	expression tag	UNP O22476
A	792	LEU	-	expression tag	UNP O22476
A	793	TYR	-	expression tag	UNP O22476
A	794	PHE	-	expression tag	UNP O22476
A	795	GLN	-	expression tag	UNP O22476
A	796	GLY	-	expression tag	UNP O22476
A	797	SER	-	expression tag	UNP O22476
A	798	ALA	-	expression tag	UNP O22476
A	799	TRP	-	expression tag	UNP O22476
A	800	SER	-	expression tag	UNP O22476
A	801	HIS	-	expression tag	UNP O22476
A	802	PRO	-	expression tag	UNP O22476
A	803	GLN	-	expression tag	UNP O22476
A	804	PHE	-	expression tag	UNP O22476
A	805	GLU	-	expression tag	UNP O22476
A	806	LYS	-	expression tag	UNP O22476
A	807	GLY	-	expression tag	UNP O22476
A	808	GLY	-	expression tag	UNP O22476
A	809	GLY	-	expression tag	UNP O22476
A	810	SER	-	expression tag	UNP O22476
A	811	GLY	-	expression tag	UNP O22476
A	812	GLY	-	expression tag	UNP O22476
A	813	GLY	-	expression tag	UNP O22476
A	814	SER	-	expression tag	UNP O22476

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	815	GLY	-	expression tag	UNP O22476
A	816	GLY	-	expression tag	UNP O22476
A	817	SER	-	expression tag	UNP O22476
A	818	ALA	-	expression tag	UNP O22476
A	819	TRP	-	expression tag	UNP O22476
A	820	SER	-	expression tag	UNP O22476
A	821	HIS	-	expression tag	UNP O22476
A	822	PRO	-	expression tag	UNP O22476
A	823	GLN	-	expression tag	UNP O22476
A	824	PHE	-	expression tag	UNP O22476
A	825	GLU	-	expression tag	UNP O22476
A	826	LYS	-	expression tag	UNP O22476
A	827	GLY	-	expression tag	UNP O22476
A	828	ALA	-	expression tag	UNP O22476
A	829	HIS	-	expression tag	UNP O22476
A	830	HIS	-	expression tag	UNP O22476
A	831	HIS	-	expression tag	UNP O22476
A	832	HIS	-	expression tag	UNP O22476
A	833	HIS	-	expression tag	UNP O22476
A	834	HIS	-	expression tag	UNP O22476
A	835	HIS	-	expression tag	UNP O22476
A	836	HIS	-	expression tag	UNP O22476
A	837	HIS	-	expression tag	UNP O22476

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	28	16	2	10	0	0	0
2	C	2	28	16	2	10	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	3	38	22	2	14	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	E	3	39	22	2	15	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
5	A	1	14	8	1	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	27	Total	O	0	0
			27	27		



- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  67% 33%



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	119.68Å 67.57Å 151.85Å 90.00° 100.65° 90.00°	Depositor
Resolution (Å)	19.89 – 2.54 19.89 – 2.54	Depositor EDS
% Data completeness (in resolution range)	98.7 (19.89-2.54) 98.7 (19.89-2.54)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 2.53Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.205 , 0.254 0.208 , 0.252	Depositor DCC
R_{free} test set	1898 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	65.6	Xtrriage
Anisotropy	0.659	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 45.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5846	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, EDO, BMA, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.31	12/5757 (0.2%)	1.10	6/7820 (0.1%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	551	ASN	CG-ND2	9.45	1.53	1.33
1	A	718	ILE	C-O	-8.34	1.14	1.24
1	A	392	LEU	N-CA	7.38	1.55	1.46
1	A	289	PRO	C-O	7.15	1.29	1.24
1	A	235	SER	N-CA	5.92	1.52	1.46
1	A	154	ASN	CG-ND2	5.78	1.45	1.33
1	A	428	PHE	C-O	-5.55	1.17	1.23
1	A	49	LYS	N-CA	5.55	1.53	1.46
1	A	221	CYS	C-O	-5.48	1.17	1.23
1	A	515	ASN	C-O	-5.42	1.16	1.24
1	A	344	SER	N-CA	5.22	1.52	1.46
1	A	249	HIS	N-CA	5.02	1.52	1.46

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	644	ASP	N-CA-C	6.54	120.47	111.71
1	A	650	PHE	N-CA-C	6.14	118.83	110.35
1	A	649	THR	N-CA-C	5.96	117.86	111.36
1	A	645	HIS	N-CA-C	5.71	119.27	111.39
1	A	151	LYS	CB-CG-CD	5.29	123.46	111.30
1	A	244	CYS	N-CA-C	-5.12	104.83	111.74

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5626	0	5536	17	0
2	B	28	0	25	1	0
2	C	28	0	25	0	0
3	D	38	0	33	0	0
4	E	39	0	34	0	0
5	A	56	0	52	0	0
6	A	4	0	6	0	0
7	A	27	0	0	0	0
All	All	5846	0	5711	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:595:LYS:O	1:A:646:THR:HG23	1.62	0.99
1:A:595:LYS:O	1:A:646:THR:CG2	2.40	0.68
1:A:597:TYR:HD1	1:A:646:THR:H	1.40	0.66
1:A:601:LYS:HB3	1:A:640:ARG:CB	2.33	0.58
1:A:155:VAL:HG23	1:A:180:LEU:HD23	1.90	0.53
1:A:155:VAL:CG2	1:A:180:LEU:HD23	2.38	0.53
1:A:295:TYR:CE2	2:B:1:NAG:H82	2.46	0.51
1:A:643:GLY:C	1:A:644:ASP:OD1	2.56	0.48
1:A:650:PHE:CE1	1:A:654:GLY:HA3	2.49	0.47
1:A:207:ILE:HG23	1:A:207:ILE:O	2.13	0.47
1:A:609:CYS:HB2	1:A:635:CYS:HA	1.99	0.44
1:A:336:PHE:HB3	1:A:364:MET:HE2	2.00	0.43
1:A:515:ASN:HA	1:A:538:LEU:HA	2.01	0.43
1:A:567:LEU:HD12	1:A:661:MET:HE2	2.00	0.43
1:A:644:ASP:OD1	1:A:644:ASP:N	2.53	0.41
1:A:765:TYR:CG	1:A:766:PRO:HA	2.55	0.41
1:A:324:SER:HA	1:A:348:SER:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	750/837 (90%)	696 (93%)	53 (7%)	1 (0%)	48 61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	647	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	648/733 (88%)	643 (99%)	5 (1%)	73 84

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119	LYS
1	A	501	THR
1	A	644	ASP
1	A	653	ASN
1	A	769	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	110	HIS
1	A	204	HIS
1	A	249	HIS
1	A	280	GLN
1	A	583	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.51	0	17,19,21	1.43	3 (17%)
2	NAG	B	2	2	14,14,15	0.85	1 (7%)	17,19,21	0.87	0
2	NAG	C	1	1,2	14,14,15	0.63	0	17,19,21	1.19	2 (11%)
2	NAG	C	2	2	14,14,15	0.59	0	17,19,21	1.22	1 (5%)
3	NAG	D	1	1,3	14,14,15	0.46	0	17,19,21	0.74	0
3	NAG	D	2	3	14,14,15	1.23	2 (14%)	17,19,21	4.11	6 (35%)
3	FUC	D	3	3	10,10,11	0.48	0	14,14,16	0.65	0
4	NAG	E	1	4,1	14,14,15	1.63	1 (7%)	17,19,21	1.03	1 (5%)
4	NAG	E	2	4	14,14,15	0.38	0	17,19,21	0.61	0
4	BMA	E	3	4	11,11,12	0.42	0	15,15,17	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	1/6/23/26	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	3/6/23/26	0/1/1/1
3	FUC	D	3	3	-	-	0/1/1/1
4	NAG	E	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	E	2	4	-	0/6/23/26	0/1/1/1
4	BMA	E	3	4	-	0/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	1	NAG	C1-C2	5.53	1.59	1.52
3	D	2	NAG	C2-N2	-3.00	1.41	1.46
2	B	2	NAG	C1-C2	2.38	1.55	1.52
3	D	2	NAG	C7-N2	-2.07	1.27	1.34

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	2	NAG	C8-C7-N2	-11.53	96.99	116.12
3	D	2	NAG	C2-N2-C7	8.78	134.66	122.90
3	D	2	NAG	O7-C7-C8	7.32	135.08	122.05
2	C	2	NAG	C1-O5-C5	4.09	117.67	112.19
4	E	1	NAG	C1-C2-N2	-3.37	105.11	110.43
3	D	2	NAG	O7-C7-N2	3.05	127.38	121.98
2	B	1	NAG	O7-C7-N2	2.82	126.97	121.98
2	B	1	NAG	O5-C1-C2	-2.69	107.13	111.29
2	C	1	NAG	C3-C4-C5	2.37	114.52	110.23
3	D	2	NAG	O3-C3-C2	2.10	113.76	109.40
2	B	1	NAG	C1-C2-N2	2.03	113.63	110.43
3	D	2	NAG	C1-O5-C5	2.03	114.90	112.19
2	C	1	NAG	O5-C5-C4	-2.01	105.93	110.83

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	2	NAG	C1-C2-N2-C7
2	C	1	NAG	C4-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
3	D	2	NAG	O7-C7-N2-C2
2	B	2	NAG	O5-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6
3	D	2	NAG	C8-C7-N2-C2


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	A	903	1	14,14,15	0.61	0	17,19,21	1.25	3 (17%)
6	EDO	A	915	-	3,3,3	0.77	0	2,2,2	0.42	0
5	NAG	A	906	1	14,14,15	0.62	0	17,19,21	1.09	1 (5%)
5	NAG	A	911	1	14,14,15	0.61	0	17,19,21	1.18	1 (5%)
5	NAG	A	910	1	14,14,15	0.78	0	17,19,21	1.54	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	903	1	-	0/6/23/26	0/1/1/1
6	EDO	A	915	-	-	1/1/1/1	-
5	NAG	A	906	1	-	0/6/23/26	0/1/1/1
5	NAG	A	911	1	-	0/6/23/26	0/1/1/1
5	NAG	A	910	1	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	910	NAG	C4-C3-C2	4.06	116.97	111.02
5	A	903	NAG	C1-C2-N2	-2.70	106.18	110.43
5	A	910	NAG	O7-C7-C8	-2.66	117.31	122.05
5	A	911	NAG	C1-O5-C5	2.57	115.63	112.19
5	A	903	NAG	C1-O5-C5	2.46	115.49	112.19
5	A	906	NAG	C2-N2-C7	2.19	125.83	122.90
5	A	903	NAG	C8-C7-N2	-2.03	112.74	116.12

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	910	NAG	O5-C5-C6-O6
6	A	915	EDO	O1-C1-C2-O2
5	A	910	NAG	C3-C2-N2-C7
5	A	910	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	744/837 (88%)	0.06	20 (2%) 56 57	32, 84, 141, 199	8 (1%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	589	ALA	6.4
1	A	592	ILE	6.3
1	A	636	ASN	5.0
1	A	637	ILE	4.3
1	A	618	PHE	4.2
1	A	648	PRO	4.1
1	A	591	PHE	4.1
1	A	588	ALA	3.6
1	A	413	ASN	3.3
1	A	359	ASP	3.0
1	A	340	SER	2.8
1	A	49	LYS	2.7
1	A	649	THR	2.7
1	A	604	GLY	2.7
1	A	410	LEU	2.7
1	A	401[A]	ASN	2.6
1	A	744	GLU	2.6
1	A	414	PRO	2.1
1	A	166	VAL	2.1
1	A	415	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BMA	E	3	11/12	0.46	0.14	122,140,146,156	0
3	NAG	D	2	14/15	0.53	0.12	120,189,214,217	0
2	NAG	C	2	14/15	0.55	0.13	141,164,182,187	0
2	NAG	C	1	14/15	0.58	0.14	93,113,137,158	0
3	FUC	D	3	10/11	0.61	0.14	106,153,170,175	0
2	NAG	B	2	14/15	0.79	0.11	74,104,124,128	0
3	NAG	D	1	14/15	0.82	0.10	111,147,176,203	0
4	NAG	E	2	14/15	0.85	0.10	90,100,106,109	0
2	NAG	B	1	14/15	0.91	0.09	62,75,85,87	0
4	NAG	E	1	14/15	0.93	0.10	74,90,93,98	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NAG	A	910	14/15	0.62	0.14	130,148,167,187	0
5	NAG	A	903	14/15	0.64	0.14	92,118,136,140	0
5	NAG	A	906	14/15	0.69	0.12	105,116,130,133	0
5	NAG	A	911	14/15	0.73	0.13	136,152,162,163	0
6	EDO	A	915	4/4	0.88	0.11	63,70,71,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.