



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 10:34 AM UTC

PDB ID : 8FIA / pdb_00008fia
Title : The structure of fly Teneurin self assembly
Authors : Bandekar, S.J.; Li, J.; Arac, D.
Deposited on : 2022-12-15
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1962	7% 78% 12% 10%
2	B	6	67% 33%
3	C	3	33% 67%
4	D	6	50% 33% 17%
5	E	3	33% 67%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	2	 50%50%

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 14844 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Teneurin-m.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1772	14201	8978	2494	2676	53	0	2	0

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	776	ALA	-	expression tag	UNP O61307
A	777	ASP	-	expression tag	UNP O61307
A	778	PRO	-	expression tag	UNP O61307
A	2732	HIS	-	expression tag	UNP O61307
A	2733	HIS	-	expression tag	UNP O61307
A	2734	HIS	-	expression tag	UNP O61307
A	2735	HIS	-	expression tag	UNP O61307
A	2736	HIS	-	expression tag	UNP O61307
A	2737	HIS	-	expression tag	UNP O61307

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	6	72	40	2	30	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	6	Total	C	N	O	0	0	0
			71	40	2	29			

- Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	E	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	F	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		
8	A	1	Total	C	O	0	0
			6	3	3		
8	A	1	Total	C	O	0	0
			6	3	3		
8	A	1	Total	C	O	0	0
			6	3	3		
8	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			7	4	3		
9	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	309	Total	O	0	0
			309	309		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Teneurin-m





- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B: 67% 33%




- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 33% 67%



- Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 50% 33% 17%



- Molecule 5: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 33% 67%



- Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	95.20Å 129.86Å 188.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.57 – 2.40 48.57 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.57-2.40) 87.6 (48.57-2.40)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.27 (at 2.27Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.216 , 0.243 0.216 , 0.239	Depositor DCC
R_{free} test set	2000 reflections (1.85%)	wwPDB-VP
Wilson B-factor (Å ²)	40.4	Xtriage
Anisotropy	0.337	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	14844	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, NAG, GOL, BMA, FUC, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/14525	0.91	9/19683 (0.0%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	978	GLY	CA-C-O	-7.19	117.28	122.45
1	A	2285	PHE	CB-CA-C	6.16	115.99	110.44
1	A	2072[A]	GLN	CA-C-O	6.13	127.02	119.79
1	A	2072[B]	GLN	CA-C-O	6.13	127.02	119.79
1	A	850	TYR	N-CA-C	6.01	117.70	111.03

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	14201	0	13874	149	0
2	B	72	0	61	0	0
3	C	38	0	34	1	0
4	D	71	0	61	2	0
5	E	39	0	34	1	0
6	F	28	0	25	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	42	0	39	1	0
8	A	30	0	40	0	0
9	A	14	0	20	1	0
10	A	309	0	0	2	0
All	All	14844	0	14188	150	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 150 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1267:LEU:HB3	1:A:1270:ASP:HB3	1.59	0.83
1:A:1940:SER:HB3	1:A:1943:ASN:HD22	1.49	0.76
1:A:1805:VAL:HG21	5:E:1:NAG:H82	1.70	0.74
1:A:1271:GLU:HB3	1:A:1274:CYS:SG	2.30	0.72
1:A:1604:THR:CG2	1:A:1608:TYR:HB3	2.21	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1750/1962 (89%)	1663 (95%)	82 (5%)	5 (0%)	36 50

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1272	ALA
1	A	2636	LYS
1	A	1811	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1271	GLU
1	A	1109	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1167/1708 (68%)	1150 (98%)	17 (2%)	57 77

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2633	HIS
1	A	2691	LEU
1	A	1949	ILE
1	A	2148	ARG
1	A	2175	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2366	HIS
1	A	2374	GLN
1	A	2628	HIS
1	A	2453	GLN
1	A	2170	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

20 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	B	1	1,2	14,14,15	0.41	0	17,19,21	0.88	1 (5%)
2	NAG	B	2	2	14,14,15	0.43	0	17,19,21	0.72	0
2	BMA	B	3	2	11,11,12	0.37	0	15,15,17	0.92	1 (6%)
2	MAN	B	4	2	11,11,12	0.21	0	15,15,17	0.66	0
2	MAN	B	5	2	11,11,12	0.19	0	15,15,17	0.60	0
2	MAN	B	6	2	11,11,12	0.28	0	15,15,17	0.65	0
3	NAG	C	1	3,1	14,14,15	0.39	0	17,19,21	1.42	3 (17%)
3	NAG	C	2	3	14,14,15	0.33	0	17,19,21	0.71	0
3	FUC	C	3	3	10,10,11	0.26	0	14,14,16	0.63	0
4	NAG	D	1	1,4	14,14,15	0.41	0	17,19,21	1.10	1 (5%)
4	NAG	D	2	4	14,14,15	0.42	0	17,19,21	0.86	1 (5%)
4	BMA	D	3	4	11,11,12	0.36	0	15,15,17	0.65	0
4	MAN	D	4	4	11,11,12	0.25	0	15,15,17	0.58	0
4	MAN	D	5	4	11,11,12	0.25	0	15,15,17	0.60	0
4	FUC	D	6	4	10,10,11	0.29	0	14,14,16	0.55	0
5	NAG	E	1	1,5	14,14,15	0.46	0	17,19,21	0.66	0
5	NAG	E	2	5	14,14,15	0.37	0	17,19,21	0.86	1 (5%)
5	BMA	E	3	5	11,11,12	0.29	0	15,15,17	0.67	0
6	NAG	F	1	1,6	14,14,15	0.38	0	17,19,21	0.89	1 (5%)
6	NAG	F	2	6	14,14,15	0.38	0	17,19,21	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	1/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	MAN	B	5	2	-	0/2/19/22	0/1/1/1
2	MAN	B	6	2	-	0/2/19/22	0/1/1/1
3	NAG	C	1	3,1	-	4/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	FUC	C	3	3	-	-	0/1/1/1
4	NAG	D	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	-	1/6/23/26	0/1/1/1
4	BMA	D	3	4	-	0/2/19/22	0/1/1/1
4	MAN	D	4	4	-	0/2/19/22	0/1/1/1
4	MAN	D	5	4	-	0/2/19/22	0/1/1/1
4	FUC	D	6	4	-	-	0/1/1/1
5	NAG	E	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	E	2	5	-	2/6/23/26	0/1/1/1
5	BMA	E	3	5	-	0/2/19/22	0/1/1/1
6	NAG	F	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	F	2	6	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C2-N2-C7	-3.46	118.27	122.90
3	C	1	NAG	O5-C1-C2	-3.17	106.39	111.29
4	D	1	NAG	O5-C1-C2	-2.82	106.93	111.29
2	B	3	BMA	C1-O5-C5	2.50	115.53	112.19
2	B	1	NAG	O5-C1-C2	-2.27	107.78	111.29

There are no chirality outliers.

5 of 15 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2
4	D	1	NAG	C8-C7-N2-C2
4	D	1	NAG	O7-C7-N2-C2
2	B	1	NAG	C8-C7-N2-C2


There are no ring outliers.



6 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	6	FUC	1	0
3	C	3	FUC	1	0
5	E	1	NAG	1	0
6	F	1	NAG	1	0
6	F	2	NAG	1	0
4	D	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	NAG	A	2810	1	14,14,15	0.30	0	17,19,21	0.89	0
9	PEG	A	2806	-	6,6,6	0.11	0	5,5,5	0.07	0
7	NAG	A	2801	1	14,14,15	0.37	0	17,19,21	1.01	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	PEG	A	2805	-	6,6,6	0.10	0	5,5,5	0.11	0
8	GOL	A	2808	-	5,5,5	0.08	0	5,5,5	0.34	0
8	GOL	A	2807	-	5,5,5	0.08	0	5,5,5	0.31	0
8	GOL	A	2804	-	5,5,5	0.08	0	5,5,5	0.29	0
7	NAG	A	2802	1	14,14,15	0.30	0	17,19,21	0.84	0
8	GOL	A	2803	-	5,5,5	0.08	0	5,5,5	0.30	0
8	GOL	A	2809	-	5,5,5	0.08	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	A	2810	1	-	5/6/23/26	0/1/1/1
9	PEG	A	2806	-	-	1/4/4/4	-
7	NAG	A	2801	1	-	3/6/23/26	0/1/1/1
9	PEG	A	2805	-	-	0/4/4/4	-
8	GOL	A	2808	-	-	0/4/4/4	-
8	GOL	A	2807	-	-	0/4/4/4	-
8	GOL	A	2804	-	-	0/4/4/4	-
7	NAG	A	2802	1	-	4/6/23/26	0/1/1/1
8	GOL	A	2803	-	-	0/4/4/4	-
8	GOL	A	2809	-	-	0/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	2801	NAG	C2-N2-C7	-2.60	119.42	122.90
7	A	2801	NAG	O5-C1-C2	2.10	114.54	111.29

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	2801	NAG	C8-C7-N2-C2
7	A	2801	NAG	O7-C7-N2-C2
7	A	2802	NAG	C3-C2-N2-C7
7	A	2802	NAG	C8-C7-N2-C2
7	A	2802	NAG	O7-C7-N2-C2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	2810	NAG	1	0
9	A	2806	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1772/1962 (90%)	0.56	135 (7%) 20 16	36, 63, 114, 166	2 (0%)

The worst 5 of 135 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2461	LEU	6.2
1	A	2638	ALA	5.3
1	A	850	TYR	5.2
1	A	851	ALA	5.1
1	A	1268	PRO	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	BMA	E	3	11/12	0.26	0.18	114,116,119,120	0
2	MAN	B	6	11/12	0.39	0.21	103,109,117,119	0
3	NAG	C	2	14/15	0.49	0.18	99,114,123,123	0
4	MAN	D	5	11/12	0.55	0.17	98,107,120,125	0
3	NAG	C	1	14/15	0.57	0.17	84,106,112,115	0
6	NAG	F	2	14/15	0.57	0.17	101,112,114,114	0
3	FUC	C	3	10/11	0.58	0.19	94,108,111,111	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	D	2	14/15	0.70	0.15	90,98,103,105	0
2	MAN	B	5	11/12	0.71	0.13	80,92,94,96	0
5	NAG	E	2	14/15	0.73	0.15	82,97,109,110	0
4	MAN	D	4	11/12	0.77	0.14	83,90,95,97	0
4	FUC	D	6	10/11	0.78	0.17	81,91,95,98	0
6	NAG	F	1	14/15	0.79	0.14	94,104,110,111	0
4	BMA	D	3	11/12	0.81	0.12	88,95,97,101	0
4	NAG	D	1	14/15	0.82	0.12	74,81,90,90	0
2	BMA	B	3	11/12	0.84	0.12	66,70,81,98	0
2	MAN	B	4	11/12	0.87	0.11	65,76,80,87	0
5	NAG	E	1	14/15	0.90	0.11	59,73,82,84	0
2	NAG	B	2	14/15	0.92	0.09	45,57,64,67	0
2	NAG	B	1	14/15	0.93	0.10	50,59,71,72	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain E:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	NAG	A	2810	14/15	0.45	0.19	96,116,120,121	0
7	NAG	A	2801	14/15	0.46	0.20	91,99,107,107	0
7	NAG	A	2802	14/15	0.50	0.17	111,117,123,124	0
8	GOL	A	2808	6/6	0.66	0.15	78,88,91,93	0
9	PEG	A	2806	7/7	0.77	0.20	61,63,70,81	0
9	PEG	A	2805	7/7	0.79	0.17	63,74,81,82	0
8	GOL	A	2809	6/6	0.83	0.18	53,67,70,74	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	GOL	A	2803	6/6	0.90	0.15	51,58,60,66	0
8	GOL	A	2804	6/6	0.90	0.18	51,55,60,64	0
8	GOL	A	2807	6/6	0.92	0.11	42,52,56,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.